

# Latest Charm Semileptonic Decay Results from CLEO-c



## OUTLINE

*CLEO-c in the context of testing the Standard Model with precision quark flavor physics.*

*Decay constants*

*Form Factors*


*CKM matrix elements*



$$\psi(3770) \rightarrow D^0 \bar{D}^0$$

$$\bar{D}^0 \rightarrow K^+ \pi^-, D^0 \rightarrow K^- e^+ \nu$$

Ian Shipsey, Purdue University  
CLEO-c Collaboration



# Big Questions in Flavor Physics

Dynamics of flavor?

Why generations?  
Why a hierarchy of masses  
& mixings?


Origin of Baryogenesis?


Sakharov's criteria: Baryon number violation  
CP violation Non-equilibrium

3 examples: Universe, kaons, beauty but Standard Model CP violation too small, need additional sources of CP violation


Connection between flavor physics & electroweak symmetry breaking?

Extensions of the Standard Model (ex: SUSY) contain flavor & CP violating couplings that should show up at some level in flavor physics, but *precision* measurements and *precision* theory are required to detect the new physics







# Precision Quark Flavor Physics




The discovery potential of B physics is limited by systematic errors from QCD:




$$\propto \left[ f^{B \rightarrow \pi}(q) \right]^2 |V_{ub}|^2$$



$$\propto \left[ f_{B_d} \right]^2 |V_{td}|^2$$



# Precision Quark Flavor Physics



The discovery potential of B physics is limited by systematic errors from QCD:

$B \rightarrow \ell^- \bar{\nu} \pi$   
 $\propto [f^{B \rightarrow \pi}(q)]^2 |V_{ub}|^2$

$B_d \rightarrow \ell^- \bar{\nu} \bar{B}_d$   
 $\propto [f_{Bd}]^2 |V_{td}|^2$


D system- CKM elements known to < 1% by unitarity

$D \rightarrow \ell^- \bar{\nu} \pi$   
 $\propto [f^{D \rightarrow \pi}(q)]^2 |V_{cd}|^2$


$D \rightarrow \ell^- \bar{\nu}$   
 $\propto [f_{D^+}]^2 |V_{cd}|^2$

→ *measurements of absolute rates for D semileptonic & leptonic decays* yield decay constants & form factors to *test* and hone QCD techniques into a *precision theory* which can be applied to the B system enabling improved determination of the apex  $(\rho, \eta)$


+  $\text{Br}(B \rightarrow D) \sim 100\%$  *absolute D hadronic rates* normalize B physics important for  $V_{cb}$  (scale of triangle) - also normalize D physics




# Precision theory + charm = large impact



Theoretical errors dominate width of bands





# Precision theory + charm = large impact



Theoretical errors dominate width of bands


*Few % precision QCD Calculations tested with few % precision charm data*  
 → theory errors of a few % on B system decay constants & semileptonic form factors






# Precision theory? Lattice QCD

BEFORE  
Quenched  
10-15%  
precision



- $f_\pi$
- $f_K$
- $3m_\Xi - m_N$
- $m_\Omega$
- $\psi(1P-1S)$
- $2m_{B_{s,av}} - m_Y$
- $Y(3S-1S)$
- $Y(2P-1S)$
- $Y(1P-1S)$
- $Y(1D-1S)$


$\frac{\text{theory-expt}}{\text{expt}}$




# Precision theory? In 2003 a breakthrough in Lattice QCD

Recent revolutionary progress in algorithms allows inclusion of QCD vacuum polarization. LQCD demonstrated it can reproduce a wide range of mass differences & decay constants. *These were postdictions*

**BEFORE**  
Quenched  
10-15%  
precision



$\frac{\text{theory-expt}}{\text{expt}}$




$\frac{\text{theory-expt}}{\text{expt}}$

This dramatic improvement needs validation of predictions.  $m(B_c)$  successful.


*Charm* decay constants  $f_{D^+}$  &  $f_{D_s}$

*Charm* semileptonic form factors

Understanding strongly coupled systems is important beyond flavor physics. LHC might discover new strongly interacting physics








# Precision theory? In 2003 a breakthrough in Lattice QCD

Recent revolutionary progress in algorithms allows inclusion of QCD vacuum polarization. LQCD demonstrated it can reproduce a wide range of mass differences & decay constants. *These were postdictions*


**BEFORE**  
Quenched  
10-15%  
precision



$\frac{\text{theory-expt}}{\text{expt}}$

- $f_\pi$
- $f_K$
- $m_\Omega$
- $3m_\Xi - m_N$
- $m_{D_s}$
- $m_D$
- $m_{D_s^*} - m_{D_s}$
- $m_\psi - m_{\eta_c}$
- $\psi(1P-1S)$
- $2m_{B_{s,2v}} - m_Y$
- $m_{B_c}$
- $Y(3S-1S)$
- $Y(2P-1S)$
- $Y(1P-1S)$
- $Y(1D-1S)$

**More**  
Quantities  
added  
2007




$\frac{\text{theory-expt}}{\text{expt}}$


This dramatic improvement needs validation of predictions.  $m(B_c)$  successful.

*Charm* decay constants  $f_{D^+}$  &  $f_{D_s}$

*Charm* semileptonic form factors

Understanding strongly coupled systems is important beyond flavor physics. LHC might discover new strongly interacting physics





# Precision Experiment for charm?

Circa 2004 (pre-CLEO-c)

Key leptonic, semileptonic & hadronic modes:

Experiment : Theory


Poorly known

$$\frac{Br}{\tau} = \Gamma$$

Measured very precisely  
0.4-0.8%

$$\frac{\delta B}{B}(D \rightarrow \pi e^+ \nu) = 45\%$$

$$\frac{\delta B}{B}(D^+ \rightarrow \mu^+ \nu) = 100\%$$




Before CLEO-c precise measurements of charm decay constants and form factors did not exist, because at Tevatron/FT/ B factories:

$$Br(D \rightarrow X) = \frac{\#X \text{ Observed}}{\text{efficiency} \times \#D\text{'s produced}}$$


Backgrounds are large.

#D's produced is usually not well known.



# CLEO-c: World's largest data sets at charm threshold

CLEO-c: Oct. 2003 – March 2008, **CESR (10GeV) → CESR-c at 4GeV**  
**CLEO III detector → CLEO-c**



$\sqrt{s}$  (MeV) Ldt (pb<sup>-1</sup>)


3686      54       $N(\psi(2S)) \approx 27M$

3773      818       $\psi(3770) \rightarrow D\bar{D} \approx 5 \times 10^6 D\bar{D}$       ←

4170      600       $D_{(s)}^{(*)} \bar{D}_{(s)}^{(*)} \approx 6 \times 10^5 D_s^* \bar{D}_s$       ←


X86 MARK III  
X25 BES II

First sample at this energy



# $\psi(3770)$ Analysis Strategy

$$e^+e^- \rightarrow \psi(3770) \rightarrow D\bar{D}$$




$\psi(3770)$  is to charm what Y(4S) is to beauty

- ❑ Pure DD, no additional particles ( $E_D = E_{\text{beam}}$ ).
- ❑  $\sigma(DD) = 6.4 \text{ nb}$  (Y(4S)  $\rightarrow$  BB  $\sim 1 \text{ nb}$ )
- ❑ Low multiplicity  $\sim 5\text{-}6$  charged particles/event


$\rightarrow$  high tag efficiency:  $\sim 20\%$  of events  
 Compared to  $\sim 0.1\%$  of B's at the Y(4S)

A little luminosity goes a long way:  
 Tagging ability:  
 # D tags in  $800 \text{ pb}^{-1}$  @ charm factory  
 $\sim$  # B tags in  $1300 \text{ fb}^{-1}$  @ Y(4S)



$$\psi(3770) \rightarrow D^+ D^-$$

$$D^+ \rightarrow K^- \pi^+ \pi^+, \quad D^- \rightarrow K^+ \pi^- \pi^-$$




# Absolute Charm Branching Ratios at Threshold


281/pb

$$E_D \Rightarrow E_{beam} : \Delta E = E_D - E_{beam} \quad M_{BC} = \sqrt{E_{beam}^2 - |\vec{p}_D|^2}$$

## 1 D reconstructed (a tag)




## 1D+ & 1D- reconstructed in same event



Independent of L and cross section



$$B(D^- \rightarrow K^+ \pi^- \pi^-) = \frac{\#(K^+ \pi^- \pi^-) \text{ Observed in tagged events}}{\text{detection efficiency for } (K^+ \pi^- \pi^-) \bullet \#D \text{ tags}}$$



# $B(D^0 \rightarrow K^- \pi^+)$

Sets scale of bd triangle

BABAR




Phys. Rev. Lett. 100, 051802 (2008)

$\mathcal{B}$ (%)	Error(%)	Source
$3.80 \pm 0.09$	2.4	PDG04
$3.891 \pm 0.035 \pm 0.069$	2.0	CLEO-c 281/pb
$4.007 \pm 0.037 \pm 0.072$	2.0	BABAR

# $B(D^+ \rightarrow K^- \pi^+ \pi^+)$

Previous best:




measure:

$$\frac{B(D^{*+} \rightarrow D^0 \pi^+) B(D^0 \rightarrow K^- \pi^+)}{B(D^{*+} \rightarrow D^+ \pi^0) B(D^+ \rightarrow K^- \pi^+ \pi^+)}$$

$B(D^+ \rightarrow K^- \pi^+ \pi^+)$   
dependent on  
 $B(D^0 \rightarrow K^- \pi^+)$

$\mathcal{B}$ (%)	Error(%)	Source
$9.3 \pm 0.6 \pm 0.8$	10.8	CLEO
$9.1 \pm 1.3 \pm 0.4$	14.9	MKIII
$9.1 \pm 0.7$	7.7	PDG04
$9.14 \pm 0.10 \pm 0.17$	1.9	CLEO-c 281/pb




Syst. limited: 2%

CLEO-c & BABAR agree vastly superior S/N at CLEO-c

*charm hadronic scale is finally on a SECURE FOUNDATION*



now:  $B(D^+ \rightarrow K^- \pi^+ \pi^+)$  independently measured

CLEO-c x 3.5 More precise than PDG






# f<sub>D+</sub> from Absolute Br(D<sup>+</sup> → μ<sup>+</sup>ν) at ψ(3770)




$$MM^2 = (E_D - E_\mu)^2 - (\vec{P}_D - \vec{P}_\mu)^2$$

where  $E_D = E_{beam}$ ,  $\vec{P}_D = -\vec{P}_{Dtag}$

	$B(D^+ \rightarrow \mu\nu) \times 10^{-4}$	$f_D$ MeV
MkIII	< 7.2	< 290
BESII	$12.2_{-5.3}^{+11.1} \pm 0.11$	$371_{-119}^{+129} \pm 25$








$f_{D^+}$  from  $D^+ \rightarrow \mu^+ \nu$

$f_{D_s}$  from  $D_s \rightarrow \mu^+ \nu$  &  $\tau^+ (\pi^+ \nu) \nu$


818 pb<sup>-1</sup> at  $\psi(3770)$

1630608-019



PRD 78,052003 (2008)

600 pb<sup>-1</sup> at 4170 MeV




PRD 79,052001(2009)  $D_s \rightarrow 3\pi / K^0 \pi / \eta \pi$

$f_{D_s}$  Method 2:  $D_s \rightarrow \tau^+ \nu, \tau^+ \rightarrow e^+ \nu \nu$


PRD 79:052002,2009





$f_D$

$f_{D_s}$



Belle

BaBar

PRL 98, 141801(2008)

CLEO-c

PRD 79,052001(2009)

PRD 79:052002,2009

Lattice(HPQCD+UKQCD)

PRL100, 062002(2008)

Lattice(FNAL+MILC+HPQCD)

PRL95, 122002(2005)

# Comparison to LQCD


CLEO  $f_D$  consistent with calculations  
(4% test of lattice) 😊

CLEO  $f_{D_s}$  (and Belle & BABAR) higher  
than most theoretical expectations

CLEO  $f_{D_s}$  is  $\sim 2.3\sigma$  above the most recent  
& precise LQCD calculations

*Ds leptonic decay width could be  
modified by new physics example:  
Dobrescu and Kronfeld arXiv:0803.0512*

**The difference between experiment HPQCD+UKQCD could be due to new physics, unlikely statistical fluctuations in experiment or lattice calculations or systematic uncertainties which are not understood in the lattice calculation or experiment. *BES III* measurements are eagerly awaited.**



# Importance of Charm Semileptonic Decays

$$\frac{d\Gamma}{dq^2} \propto |V_{cs(d)}|^2 |f_+^{D \rightarrow (K)\pi}(q^2)|^2$$

- 1 Assuming theoretical form factors  $\Rightarrow V_{cs}$  and  $V_{cd}$
- 2 Assuming  $V_{cs}$  and  $V_{cd}$  known, we can check theoretical calculations of the form factors
- 3 *Potentially* useful input to  $V_{ub}$  from exclusive B semileptonic decays


$Br(B \rightarrow \pi l \nu) \sim 6\%$  precision  
BABAR/Belle/CLEO(HFAG)

(summer 2008)


Expt. 5%  
 $q^2 > 16 \text{ GeV}^2$

$$|V_{ub}| = (3.62 \pm 0.22 \pm_{-0.41}^{+0.63}) \times 10^{-3}$$

$\pm \text{exp} \pm \text{LQCD}$




~11- 17% e.g.  
HPQCD & FNAL



$$\propto [f^{B \rightarrow \pi}(q)]^2 |V_{ub}|^2$$

$$\propto [f^{D \rightarrow \pi}(q)]^2 |V_{cd}|^2$$

Related at  
same invariant  
4 velocity





# Absolute Semileptonic Branching Fractions


The neutrino direction is determined to  $1^\circ$

no kinematics ambiguity

$$U \equiv E_{miss} - |\vec{p}_{miss}| = 0$$




Tagging creates a single D beam of known 4-momentum



$$U = E_{miss} - |\mathbf{P}_{miss}| \text{ (GeV)}$$



$$\mathcal{B}(D \rightarrow K e \nu) = \frac{N(D \rightarrow K e \nu)}{\text{Efficiency} \times N_{\text{tags}}}$$



arXiv:0906.2983  
(accepted PRD Jul 22 2009)

CLEOIII 10 GeV

CLEO-c




Compare to:  
state of the  
art measurement  
at 10 GeV (CLEO III)  
PRL 94, 11802 (2004)

Tag with  $D^{*+} \rightarrow D^0 \pi_s$   
 $D^0 \rightarrow \pi^- \ell^+ \nu$   
observable:  
 $\Delta m = m(\pi_s \pi \ell \nu) - m(\pi \ell \nu)$

Only other high statistics measurement is from Belle  
282/fb (x350 CLEO-c)  $222 \pm 17$  events S/N 4/1


Note:  
kinematic  
separation.




# CLEO-c semileptonic tagging analysis technique: big impact

## 1<sup>st</sup> Observations:


$$D^0 \rightarrow \rho^- e^+ \nu_e$$




$$D^+ \rightarrow \eta e^+ \nu_e$$



$$D^+ \rightarrow \omega e^+ \nu_e$$



$$D^0 \rightarrow K^- \pi^+ \pi^- e^+ \nu_e$$




$$+ D^+ / D^0 \rightarrow X e^+ \nu_e$$

$$D \rightarrow K^* e^+ \nu_e$$

form factors

## Precision Measurements:



Note: use PDG2004, as PDG2006 & PDG2008 are dominated by CLEO-c measurements

PRL, 97, 251801 (2006);  
 arXiv:0906.2983(accepted PRD);  
 PRL, 100, 251802(2008);  
 PRD, 77, 112005(2008);  
 PRD, 74, 052001(2006);  
 PRL, 102, 081801(2009);  
 PRL, 99, 191801 (2007);


\*  $D \rightarrow K^* e^+ \nu$  branching fractions are for 56/pb  
 $D \rightarrow \pi^- / K^- / \pi^0 / \bar{K}^0 e^+ \nu$  branching fractions are for 818/pb

CLEO's measurements most precise for ALL modes; *4 modes* observed for the first time



# D → K, π eν Branching Fractions


**D → K e<sup>+</sup> ν**



$B(D^0 \rightarrow K^- e^+ \nu)$   
 3.50(3)(4)%  
 (CLEO-c 818 pb<sup>-1</sup>)


$\sigma(B(Ke\nu)) / B(Ke\nu) \sim 1.4\%$   
 $\sigma(B(\pi e\nu)) / B(\pi e\nu) \sim 3.0\%$

**D → π e<sup>+</sup> ν**



$B(D^0 \rightarrow \pi^- e^+ \nu)$   
 0.288(8)(3)%  
 arXiv:0906.2983  
 (accepted PRD Jul 22 2009)

Precision measurements from BABAR/Belle/CLEO-c.  
 CLEO-c most precise. Theoretical precision lags experiment.





# Measuring the form factor in $D \rightarrow K/\pi e \nu$

$$\frac{d\Gamma(D \rightarrow K e \nu)}{dq^2} = \frac{G_F^2}{24\pi^3} P_K^3 |f_+(q^2)|^2 |V_{cs}|^2$$

$$q^2 = m_W^2 = (P_e + P_\nu)^2$$

Form factor  $\rightarrow$  probability hadron forms as a function of  $q^2$



□ We perform binned likelihood fits to  $U$  distributions in each  $q^2$  bin and each tag mode


□ Signal shapes are taken from signal MC, smeared with double Gaussians

□ Background shapes are taken from MC with all  $\overline{D}D$  and non- $\overline{D}D$  decays

S/N	~300/1
Signal events	~14000
U resolution	~10 MeV
$q^2$ resolution	~0.008 $\text{GeV}^2/c^4$

arXiv:0906.2983 (accepted PRD Jul 22 2009)






# D → K/π e<sup>+</sup> ν : Fits to the dΓ/dq<sup>2</sup> Distributions

$$\frac{d\Gamma}{dq^2} = \frac{G_F^2}{24\pi^3} P_{K/\pi}^3 |f_+(q^2)|^2 |V_{cx}|^2$$

Fit to Becher-Hill Series


$$f_+(q^2) = \frac{1}{P(q^2)\phi(q^2, 0)} \left[ \sum_k a_k z^k(q^2, 0) \right]$$



Other form factor parameterizations exist, but are only used as functional forms as their physical pictures are not supported by the data

Simultaneous fits to isospin conjugate modes are also performed


Experimentally measured decay rates  $\Gamma_i^{measured}$



Theoretically predicted decay rates

$$\Gamma_i^{predicted} = \int d\Gamma = \frac{G_F^2 |V_{Qq'}|^2}{24\pi^3} \int |f_+(q^2)|^2 p_P^3 dq^2$$







# $D \rightarrow Ke^+ \nu$ Form Factor: test of LQCD

$$\frac{d\Gamma}{dq^2} = \frac{G_F^2}{24\pi^3} P_K^3 |f_+(q^2)|^2 |V_{cs}|^2$$


Form factor measures probability hadron will be formed



## Shape: $\alpha(K e \nu)$




## Normalization: $f_+^K(0)$



Modified pole model used as example

$$f_+(q^2) = \frac{f_+(0)}{(1 - q^2/m_{pole}^2)(1 - \alpha q^2/m_{pole}^2)}$$


Normalization: experiments (1.2%) consistent with LQCD (10%). *Theoretical precision lags.*  
 CLEO-c prefers smaller value for shape parameter,  $\alpha$




# $D \rightarrow Ke^+ \nu$ Form Factor: test of LQCD

$$\frac{d\Gamma}{dq^2} = \frac{G_F^2}{24\pi^3} P_K^3 |f_+(q^2)|^2 |V_{cs}|^2$$


Form factor measures probability hadron will be formed




Shape:  $\alpha(K_{e\nu})$



Normalization:  $f_+^K(0)$




CLEO-c prefers smaller value for shape parameter,  $\alpha$   
 Normalization: experiments (1.2%) consistent with  
 LQCD (10%). *Theoretical precision lags.*




# $D^0 \rightarrow \pi^- e^+ \nu$ Form Factor: test of LQCD


$$\frac{d\Gamma}{dq^2} = \frac{G_F^2}{24\pi^3} P_\pi^3 |f_+(q^2)|^2 |V_{cd}|^2$$



shape:  $\alpha(\pi e \nu)$




Normalization:  $f_+^\pi(0)$



Modified pole model used as example


$$f_+(q^2) = \frac{f_+(0)}{(1 - q^2/m_{pole}^2)(1 - \alpha q^2/m_{pole}^2)}$$

Normalization experiments (2%) consistent with LQCD (10%). CLEO-c is most precise. *Theoretical precision lags.*



# $D^0 \rightarrow \pi^- e^+ \nu$ Form Factor: test of LQCD

$$\frac{d\Gamma}{dq^2} = \frac{G_F^2}{24\pi^3} P_\pi^3 |f_+(q^2)|^2 |V_{cd}|^2$$



$\pi$  fast

$\pi$  at rest



Normalization:  $f_+^\pi(0)$


Modified pole model used as example

$$f_+(q^2) = \frac{f_+(0)}{(1 - q^2/m_{pole}^2)(1 - \alpha q^2/m_{pole}^2)}$$

Normalization experiments (2%) consistent with LQCD (10%). CLEO-c is most precise. *Theoretical precision lags.*

The data determines  $|V_{cd}|f_+(q^2)$ . To extract  $|V_{cd}|$  we fit to  $|V_{cd}|f_+(q^2)$  using Becher-Hill  $z$ -expansion to determine  $|V_{cd}|f_+(0)$  & use  $f_+(0)$  from LQCD (FNAL-MILC-HPQCD.) Same for  $|V_{cs}|$





# $|V_{cs}|$ & $|V_{cd}|$ Results

arXiv:0906.2983  
(accepted PRD Jul 22 2009)



CLEO-c: the most precise *direct* determination of  $V_{cs}$

$$\sigma(|V_{cs}|)/|V_{cs}| \sim 1.1\%(\text{expt}) \oplus 10\%(\text{theory})$$

CLEO - c	$ V_{cs} $		
(818 pb <sup>-1</sup> )	0.985	± 0.009	± 0.006 ± 0.103
	stat	syst	theory


CLEO-c:  $\sigma(|V_{cd}|)/|V_{cd}| \sim 3.1\%(\text{expt}) \oplus 10\%(\text{theory})$   
 $\nu N$  remains most precise determination (*for now*)

CLEO - c	$ V_{cd} $		
(818 pb <sup>-1</sup> )	0.234	± 0.007	± 0.002 ± 0.025
	stat	syst	theory



\* PDG2002

Fits use Becher-Hill z-expansion



# $|V_{cs}|$ & $|V_{cd}|$ Results


arXiv:0906.2983  
(accepted PRD Jul 22 2009)

THEORY UNCERTAINTY REMOVED

CLEO-c: the most precise *direct* determination of  $V_{cs}$


$$\sigma(|V_{cs}|)/|V_{cs}| \sim 1.1\%(\text{expt}) \oplus 10\%(\text{theory})$$

CLEO - c	$ V_{cs} $		
(818 pb <sup>-1</sup> )	0.985	± 0.009	± 0.006 ± 0.103
	stat	syst	theory




CLEO-c:  $\sigma(|V_{cd}|)/|V_{cd}| \sim 3.1\%(\text{expt}) \oplus 10\%(\text{theory})$   
 $\nu N$  remains most precise determination (*for now*)

CLEO - c	$ V_{cd} $		
(818 pb <sup>-1</sup> )	0.234	± 0.007	± 0.002 ± 0.025
	stat	syst	theory



LQCD form factors with improved precision are eagerly awaited

\* PDG2002



# Unitarity Test: Compatibility of charm & beauty sectors of CKM matrix?

arXiv:0906.2983  
(accepted PRD Jul 22 2009)

$|V_{cd}|$  &  $|V_{cs}|$  indirect

1) K & nucleon

$$|V_{ud}| \approx |V_{cs}| \quad \& \quad |V_{cd}| \approx |V_{us}|$$

2) B physics

Indirect = global CKM fit = 1+2

$|V_{cd}|$  &  $|V_{cs}|$  direct

(D semileptonic decays CLEO)


CLEO-c full data set

$$\sigma(|V_{cd}|) / |V_{cd}| \sim 3.1\% \oplus \text{theory}$$

$$\sigma(|V_{cs}|) / |V_{cs}| \sim 1.1\% \oplus \text{theory}$$

D semileptonic decays with comparable theory and experimental uncertainty may lead to interesting competition between direct and indirect constraints  
We eagerly await new precise lattice calculations

CLEO-c full data set + 3-4% theory uncertainties



Plots by Sebastien Descotes-Genon & Ian Shipsey

See also talk by Descotes-Genon at joint BABAR-Belle-BESIII-CLEO-c Workshop 11/07, Beijing




# Observe 6 Exclusive $D_s$ Semileptonic Decays

- Similar to  $D_s \rightarrow \mu\nu$  analysis: tag; reconstruct visible parts of signal; plot  $MM^2$
- First absolute branching fraction measurements for  $D_s$  SL decays
- Total width of these exclusive modes is 16% lower than the  $D^0/D^+$  semileptonic widths.
- Shed light on  $\eta$ - $\eta'$ -glueball mixing
- Observation of a semileptonic decay including a scalar meson in the final state.

arXiv:0903:0601


Signal Mode	$\mathcal{B}(\%)$
$D_s^+ \rightarrow \phi e^+ \nu_e$	$2.29 \pm 0.37 \pm 0.11$
$D_s^+ \rightarrow \eta e^+ \nu_e$	$2.48 \pm 0.29 \pm 0.13$
$D_s^+ \rightarrow \eta' e^+ \nu_e$	$0.91 \pm 0.33 \pm 0.05$
$D_s^+ \rightarrow K^0 e^+ \nu_e$	$0.37 \pm 0.10 \pm 0.02$
$D_s^+ \rightarrow K^{*0} e^+ \nu_e$	$0.18 \pm 0.07 \pm 0.01$
$D_s^+ \rightarrow f_0 e^+ \nu_e$	$0.13 \pm 0.04 \pm 0.01$

$$B(D_s^+ \rightarrow f_0(980)e^+\nu) \times B(f_0 \rightarrow \pi^+\pi^-) \rightarrow$$




310 pb<sup>-1</sup> @4170 (Half of full dataset)





# $D_s^+ \rightarrow f_0(980)e^+\nu$

□  $D_s$  semileptonic decays provide a very clean environment to study the properties of the  $f_0(980)$  meson



600 pb<sup>-1</sup> @4170

(CLEO-c full dataset)

□ It is suggested that  $B_s \rightarrow J/\Psi f_0$  can be an alternative to  $B_s \rightarrow J/\Psi \Phi$  to measure CP Violation in the  $B_s$  system  
Stone & Zhang [PRD79, 074024]

arXiv: 0907.3201

(submitted to PRD Jul 18 2009)


□ Many interesting results:


- ✓  $B(D_s^+ \rightarrow f_0(980)e^+\nu, f_0 \rightarrow \pi^+\pi^-) = (0.20 \pm 0.03 \pm 0.01)\%$
- ✓  $B(D_s^+ \rightarrow \phi e^+\nu) = (2.36 \pm 0.23 \pm 0.13)\%$

$$\frac{\Gamma(D_s^+ \rightarrow f_0(980)e^+\nu, f_0 \rightarrow \pi^+\pi^-)}{\Gamma(D_s^+ \rightarrow \phi e^+\nu, \phi \rightarrow K^+K^-)} \Bigg|_{q^2=0} = (42 \pm 11)\%$$

$$\left[ \text{Predicted to equal } \frac{\Gamma(B_s \rightarrow J/\Psi f_0(980), f_0 \rightarrow \pi^+\pi^-)}{\Gamma(B_s \rightarrow J/\Psi \phi, \phi \rightarrow K^+K^-)} \right]$$

- ✓  $M_{f_0(980)} = (977_{-9}^{+11} \pm 1) \text{ MeV}, \Gamma_{f_0(980)} = (91_{-22}^{+30} \pm 3) \text{ MeV}$
- ✓ Simple pole model  $M_{\text{pole}} = (1.7_{-0.7}^{+4.5} \pm 0.2) \text{ GeV}$








# Inclusive Semileptonic Decays of $D^0, D^+,$ and $D_s$

NEW

PRELIMINARY




	$D^0 \rightarrow X e^+ \nu$	$D^+ \rightarrow X e^+ \nu$	$D_s \rightarrow X e^+ \nu$
Inclusive $\mathcal{B}$ (%)	$6.55 \pm 0.10 \pm 0.09$	$16.36 \pm 0.11 \pm 0.29$	$6.49 \pm 0.40 \pm 0.18$
Sum of exclusive $\mathcal{B}$ (%)	$6.1 \pm 0.2 \pm 0.2$	$15.1 \pm 0.5 \pm 0.5$	$6.47 \pm 0.60$

□ Use knowledge of D semileptonic decay to extrapolate below the momentum cutoff (200 MeV/c)

Any additional exclusive modes will have small branching ratios

$$\Gamma_{D^+}^{SL} / \Gamma_{D^0}^{SL} = 0.99 \pm 0.02 \pm 0.02 \quad \text{Isospin symmetry}$$

$$\Gamma_{D_s^+}^{SL} / \Gamma_{D^0}^{SL} = 0.81 \pm 0.05 \pm 0.03 \quad \text{SU(3) is broken}$$



## Summary of CLEO-c CKM physics motivated measurements

CLEO-c hadronic  $D^0$ ,  $D^+$  and  $D_s$  branching fractions more precise than PDG averages: (for  $D^0$ ,  $D^+$  2% precision is syst.limited) CLEO establishes charm hadronic scale

Most precise:  $f_{D^+} = (205.8 \pm 8.5 \pm 2.5)$  MeV consistent with LQCD  $\rightarrow$  1% (2 MeV) full data

Most precise:  $f_{D_s} = (259.5 \pm 6.6 \pm 3.1)$  MeV  $\sim 2.3\sigma$  higher than LQCD.

To interpret as "prosaic" or "exciting": requires more data (BES III)

Most precise measurement of form factors magnitudes in  $D \rightarrow K/\pi e \nu$

Most precise  $|V_{cs}| = 0.985 \pm 0.009 \pm 0.006 \pm 0.103_{\text{theory}}$


$|V_{cd}| = 0.234 \pm 0.007 \pm 0.002 \pm 0.025_{\text{theory}}$

Most precise determination from semileptonic decay


6 exclusive  $D_s$  semileptonic decays & measurement of form factor in  $D_s \rightarrow f_0(980)e^+ \nu$ .  $SU(3)$  is broken in  $D \rightarrow X e \nu$  decays

$\sim 90$  CLEO-c papers now published or submitted, & many more analyses to come.

Longer term the charm factory mantle passes to BES III




# Precision theory + charm = large impact



- \* CLEO-c: a major contribution to the goal the lower plot represents
- \* LQCD has been validated at the 4% level ( $f_{D^+}$ )
- \* A triumph for theory & experiment!

More precise LQCD form factor calculations needed  
more data → BESIII



*Few % precision QCD*  
 Calculations tested with few % precision charm data  
 → theory errors of a few % on B system decay constants & semileptonic form factors