



Search for $K_L \rightarrow \pi^0\pi^0\mu^+\mu^-$ with KTeV data



*American Physical Society
Division of Particles & Fields
Detroit 27 - 31 July 2009*

*Leo Bellantoni
Fermi National Accelerator Lab
For the KTeV Collaboration*

The HyperCP Result

Park, et.al. (HyperCP), Phys.Rev.Lett. 94,021801 (2005)


- $Br(\Sigma^+ \rightarrow p\mu^+\mu^-) = (8.6^{+6.6}_{-5.4\text{STAT}} \pm 5.5_{\text{SYST}}) \times 10^{-8}$ (3 events)
- Expectation is $\sim 0.1 \times 10^{-8}$
- All 3 events at the same mass: 214.3 ± 0.5 MeV
- C.L. for this in S.M. is 0.8%




Resonance with width = detector resolution

The HyperCP Result

Maybe it isn't just



Maybe there is also



That would change the detection efficiency


$$Br(\Sigma^+ \rightarrow pP^0, P^0 \rightarrow \mu^+\mu^-) = [3.1_{-1.9}^{+2.4} \pm 1.5] \times 10^{-8}$$

\Rightarrow Partial width of

$$\Gamma(\Sigma^+ \rightarrow pP^0, P^0 \rightarrow \mu^+\mu^-) = 2.5 \times 10^{-19} \text{ MeV}$$

Should appear at other s - d vertices

In Kaon Decays



The existing measurement of
 $Br(K^+ \rightarrow \pi^+ \mu^+ \mu^-) = 8.1 \times 10^{-8}$
 corresponds to a $\Gamma = 4.3 \times 10^{-21}$ MeV
 Almost 2 orders of magnitude
 below the *Hyper CP* result

Maybe $K \rightarrow \pi \mu^+ \mu^-$ is parity suppressed(?) - look for $K_L \rightarrow \pi \pi \mu^+ \mu^-$

In fact, look for $\pi^0 \pi^0$ not $\pi^+ \pi^-$:

$$m(K_L) - 2[m(\pi^0) + m(\mu^\pm)] = 16.3 \text{ MeV}$$

$$m(K_L) - 2[m(\pi^\pm) + m(\mu^\pm)] = 7.2 \text{ MeV}$$

} Factor of
 10 in phase
 space!

The CsI calorimeter of KTeV is very good at finding π^0 s

Flurry Of Excitement

Gorbunov & Rubakov, Phys.Rev.D 73,035002 (2006)

It might be Sgoldstino!

He, Tandeen & Valencia, Phys.Rev.D 72,074003 (2005)

Reevaluate the standard model branching ratio - the Br measurement is consistent with it

He, Tandeen & Valencia, Phys.Lett.B 631,100 (2005)

Can't be scalar or vector couplings

If pseudoscalar, $Br(K_L \rightarrow \pi^0 \pi^0 X^0 \rightarrow \pi^0 \pi^0 \mu^+ \mu^-) \sim 8.3 \times 10^{-9}$

If axial-vector, $Br(K_L \rightarrow \pi^0 \pi^0 X^0 \rightarrow \pi^0 \pi^0 \mu^+ \mu^-) \sim 1.0 \times 10^{-10}$

He, Tandeen & Valencia, Phys.Rev.Lett, 98,081812 (2007)

It could be the pseudoscalar a of the NMSSM




Leading to a searches by CLEO, BaBar, D0

Deshpande, Eilam & Jiang, Phys.Lett.B 632,212 (2006)


If a pseudoscalar, $Br(K_L \rightarrow \pi^0 \pi^0 X^0 \rightarrow \pi^0 \pi^0 \mu^+ \mu^-) = 8.02 \times 10^{-9}$

E391a set a 90% C.L. limit on $Br(K_L \rightarrow \pi^0 \pi^0 X^0; X^0 \rightarrow \gamma\gamma) < 2.4 \times 10^{-7}$

The KTeV Detector



The KTeV Detector



E.T.Worcester
Thesis

Pure CsI calorimeter: Energy resolution $< 1\%$ for $> 10 \text{ GeV } \gamma$
Position resolution $\sim 1 \text{ mm}$ for e^\pm

Muon system: Total thickness 5.1m of steel $\sim 31 \lambda_I$
Efficiency $> 98\%$ for $> 10 \text{ GeV } \mu^\pm$
 π^\pm punchthrough probability $\sim (1.69 + 0.17P [\text{GeV}]) / 1000$

The KTeV Detector

KTeV actually was 2 experiments:

E832 - precision measurement of $\Re(\epsilon'/\epsilon)$

E799 - kaon rare decay program

Data taken 1996-1997, 1999-2000

E799 configuration collected 733 $\times 10^9$ K_L decays

Previous KTeV Dimuon Results:

$Br(K_L \rightarrow \mu^+ \mu^- \gamma) = (3.62 \pm 0.04_{\text{STAT}} \pm 0.08_{\text{SYST}}) \times 10^{-7}$ 9327 events

PRL 87, 071801 (2001)

$Br(K_L \rightarrow e^+ e^- \mu^+ \mu^-) = (2.69 \pm 0.24_{\text{STAT}} \pm 0.12_{\text{SYST}}) \times 10^{-9}$ 132 events


PRL 90, 141801 (2003)

$Br(K_L \rightarrow \pi^0 \mu^+ \mu^-) < 3.8 \times 10^{-10}$ 2 events

PRL 84, 5279 (2000)

733 x10⁹ K_L decays

We measure the number of $K_L \rightarrow \pi^0 \pi^0 \pi^0$; $\pi^0 \rightarrow \gamma e^+ e^-$
A mode of similar signature and the Brs are known



1997 $\pi^0 \pi^0 \pi^0$ Inv. Mass

1997 $\pi^0 \pi^0 \pi^0$ P_T^2

● = Data
□ = MC

Signal Characteristics

$\pi^0 \rightarrow \gamma\gamma$

4 CsI clusters without matching tracks

$\mu + \text{matter} \rightarrow \text{ions}$

2 opposite-sign tracks with hits in μ system and $> 7 \text{ GeV}$

$\mu + \text{CsI} \rightarrow \text{not much}$

CsI cluster matching track has $E < 1 \text{ GeV}$

All decay products reconstructed

Sum of momenta \perp to K line of flight $< \sqrt{0.001} \text{ GeV}$

$m(K_L) - 2m(\pi^0) > m(\mu\mu)$

$m(\mu\mu) < 232 \text{ MeV}$



π^0 and μ pair originate at K_L decay point

$m(\pi^0)$ within 9 MeV

Signal Boxes

Two boxes to open... one for the K_L and one for the X^0

Σp_{\perp}^2



m_{RECO}

Simulated signal

Signal Sensitivity

1997 Acceptance ($K_L \rightarrow \pi^0 \pi^0 \mu^+ \mu^-$) = $(3.14 \pm 0.004_{stat.})\%$
[flat phase space]

1997 Acceptance ($K_L \rightarrow \pi^0 \pi^0 X^0 \rightarrow \pi^0 \pi^0 \mu^+ \mu^-$) = $(2.80 \pm 0.004_{stat.})\%$

1997 Acceptance ($K_L \rightarrow \pi^0 \pi^0 \pi^0_D$) = $(5.94 \pm 0.02_{stat.}) \times 10^{-5}$

1999 Acceptance ($K_L \rightarrow \pi^0 \pi^0 \mu^+ \mu^-$) = $(4.03 \pm 0.005_{stat.})\%$

1999 Acceptance ($K_L \rightarrow \pi^0 \pi^0 X^0 \rightarrow \pi^0 \pi^0 \mu^+ \mu^-$) = $(3.74 \pm 0.004_{stat.})\%$

1999 Acceptance ($K_L \rightarrow \pi^0 \pi^0 \pi^0_D$) = $(1.29 \pm 0.003_{stat.}) \times 10^{-4}$

A single detected signal event corresponds
to a Br of 4.1×10^{-11}

Backgrounds


Decay Mode	# '97 MC events generated	# '99 MC events generated
$K_{\mu 3}^0$ (punch through)	~ 2.6 Billion (0.039 f)	1,752,020,868 (0.027 f)
$K_{\mu 3}^0$ (pion decay = $\pi^{+-} \rightarrow \mu^{+-} \nu_{\mu}$)	244,692,689 (0.0037 f)	421,656,663 (0.0064 f)
$K_{\mu 4}^0$ (punch through)	120,066,571 (8.38 f)	96,372,292 (6.72 f)
$K_{\mu 4}^0$ (pion decay) *	93,373,819 (6.51 f)	109,831,267 (7.66 f)
$K_L \rightarrow \pi^+ \pi^- \pi^0$ (2x punch through)	1,848,796,492 (0.060 f)	1,062,004,339 (0.035 f)
$K_L \rightarrow \pi^+ \pi^- \pi^0$ (2x pion decay)	85,552,978 (0.0028 f)	106,912,811 (0.0035 f)
$K_L \rightarrow \pi^+ \pi^- \pi^0$ (punch & decay)	455,374,316 (0.015 f)	456,480,690 (0.015 f)
$K_L \rightarrow \pi^+ \pi^- \gamma$ (2x punch through)	15,034,557 (1.41 f)	21,646,250 (2.03 f)
$K_L \rightarrow \pi^+ \pi^- \gamma$ (2x pion decay)	20,304,857 (1.90 f)	16,311,114 (1.53 f)
$K_L \rightarrow \pi^+ \pi^- \gamma$ (punch & decay)	14,249,908 (1.34 f)	14,495,323 (1.36 f)
$K_L \rightarrow \pi^+ \pi^-$ (2x punch through)	683,676,428 (1.35 f)	671,923,195 (1.32 f)
$K_L \rightarrow \pi^+ \pi^-$ (2x pion decay)	8,529,573 (0.017 f)	21,840,183 (0.044 f)
$K_L \rightarrow \pi^+ \pi^-$ (punch & decay)	50,306,906 (0.100 f)	26,557,616 (0.053 f)
$K_L \rightarrow \mu^+ \mu^-$	1,183,635 (670.0 f)	5,240,705 (2967 f)
$K_L \rightarrow \mu^+ \mu^- \gamma$	9,582,978 (109.8 f)	119,650,358 (1372 f)
$K_L \rightarrow \mu^+ \mu^- \gamma \gamma$	10,869,003 (4473 f)	48,801,465 (20084 f)
$K_L \rightarrow \pi^0 \mu^+ \mu^-$	11,042,193	13,008,645

Not one of these events passed event selection

Backgrounds

Not a lot of background in the data either


3 events that all look like accidentals



Backgrounds

Assume no background at all

not subtracting background is more conservative than doing the subtraction



Systematic Uncertainties

Source of Systematic Error	$\frac{\Delta F_{Norm,1997}}{F_{Norm,1997}}$	$\frac{\Delta F_{Norm,1999}}{F_{Norm,1999}}$
$(473 \mp 1) \text{ MeV} \leq M_{\text{rec.}\pi^0} \leq (523 \pm 1) \text{ MeV}$	+0.04% -0.05%	+0.05% -0.06%
$ M_{\text{rec.}\pi^0} - M_{\pi^0} \leq (14 \pm 1) \text{ MeV}$	+0.02% -0.03%	+0.02% +0.01%
$(94.0 \mp 1.0) \text{ m} \leq Z_{\text{VTX}} \leq (158.0 \pm 1.0) \text{ m}$	+0.16% +0.02%	+0.20% -0.10%
$P_T^2 \leq (1.0 \pm 0.1) * 10^{-3} \text{ GeV}^2$	+0.11% +0.02%	+0.06% -0.08%
$(0.95 \mp 0.1) \leq E_{\text{cl}}(\text{track}) / p_{\text{track}} \leq (1.05 \pm 0.1)$	+1.24% -2.41%	+2.23% -4.05%
P_z Weighting	-----	1.87%
Cracks in μ Counting Planes	0.50%	0.50%
Energy Loss in μ Filters	0.40%	0.40%
$\text{Br}(K_L \rightarrow \pi^0 \pi^0 \pi^0)$	0.61%	0.61%
Total Systematic Error from Flux	+1.54% - 2.57%	+3.05% - 4.55%

Preliminary!

Result

There were no events in the signal boxes when opened

$$\text{At 90\% C.L. } Br < 2.30 \left(1 + \frac{2.30}{2} \left[\frac{\sigma_R (SES^{-1})}{SES^{-1}} \right]^2 \right) \cdot SES$$

$$Br(K_L \rightarrow \pi^0 \pi^0 \mu^+ \mu^-) < 8.63 \times 10^{-11}$$

$$Br(K_L \rightarrow \pi^0 \pi^0 X^0 \rightarrow \pi^0 \pi^0 \mu^+ \mu^-) < 9.44 \times 10^{-11}$$

Preliminary!!!

Conclusions

The 90% C.L. upper limit on $Br(K_L \rightarrow \pi^0 \pi^0 X^0 \rightarrow \pi^0 \pi^0 \mu^+ \mu^-)$ is about 80 times the Br that would obtain if the *HyperCP* result were due to new pseudoscalar couplings

The 90% C.L. upper limit on $Br(K_L \rightarrow \pi^0 \pi^0 X^0 \rightarrow \pi^0 \pi^0 \mu^+ \mu^-)$ is only a little below the Br that would obtain if the *HyperCP* result were due to new axial-vector couplings
This must still be regarded as possible

The possibility that the *HyperCP* result is due to new tensor couplings remains to be examined

The transition $D \rightarrow D^* \mu^+ \mu^-$ → Dr. David Phillips
has also been observed