

MC-TESTER:

a universal tool for comparisons
of Monte Carlo predictions
in High Energy Physics

Piotr Golonka

Piotr.Golonka@CERN.CH


CERN EP/ATR, AGH/INP Kraków

Zbigniew Waś (CERN/TH, INP Kraków)

Tomasz Pierzchała (UŚI Katowice)

LCG Generator Services monthly meeting, 16 October 2003

MC-TESTER: why?




- Comparisons of physics content
- Debugging, development
- Library versions, maintainance
- Automated tests of installations of large MC systems

Areas of use:

- **Exploration tool**
 - for new (versions/implementations of) generators
- **Comparisons of physics in generators:**
 - Particles' decays
 - Parton-level ($2 \rightarrow n$ processes)
- **Debugging:**
 - *Generator versions (“flavours”)*
 - Porting the code (F77 \rightarrow C++ transition)
 - Phase-space inefficiencies
 - *Event record formats*


MC-TESTER analysis:

- *Generation step:*



- For every event produced by tested generator, certain (decay) processes are searched for and identified

- For identified process, all possible *invariant masses* are histogrammed




- At the end of the run, histogrammes are stored in root file

- *Analysis step:*

- Creates a table of all processes, matches the corresponding processes from two runs
- Calculates branching ratios (and statistical errors)
- Performs statistical analysis all histograms from two generated root files, produces EPS files (and root file)
- Produces a complete booklet (.ps)


MC-TESTER analysis:

GENERATION STEP




MC-TESTER
libraries

ROOT
FILE




-decay channels
-histogrammes



MC-TESTER
libraries

ROOT
FILE



ANALYSIS STEP




- list of decay channels
- branching ratios

Tex
FILE

analyzed
histogrammes



BOOKLET (.ps)



ROOT
FILE

EPS
FILES

Features:

- **Written completely in C++**
(compiled code and ROOT macros)
- **Directly usable from F77 code**
- **Simple: two F77 routines** to be called in the code
- **Support for HEPEVT,LUJETS,PYJETS**
- **Runtime parameters specified by root macros**
(flexibility! e.g. read parameters from external file!)
- **Easy to integrate/use with existing MC installations** (2 .so/.a libraries + root libs)
- **Examples of use provided** (TAUOLA, PYTHIA)
- **Complete analysis environment**

Table of decay modes:

Found decay modes:

Decay channel	Branching Ratio \pm Rough Errors		Max. shape dif. param.
	Generator #1	Generator #2	
$\tau^- \rightarrow \nu_\tau \pi^0 \pi^-$	25.3683 \pm 0.0159%	25.3085 \pm 0.0159%	0.04375
$\tau^- \rightarrow e^- \tilde{\nu}_e \nu_\tau$	17.8479 \pm 0.0134%	18.1093 \pm 0.0135%	0.00000
$\tau^- \rightarrow \mu^- \tilde{\nu}_\mu \nu_\tau$	17.3866 \pm 0.0132%	17.6326 \pm 0.0133%	0.00000
$\tau^- \rightarrow \nu_\tau \pi^-$	11.0768 \pm 0.0105%	11.1765 \pm 0.0106%	0.00000
$\tau^- \rightarrow \nu_\tau \pi^0 \pi^0 \pi^-$	9.1865 \pm 0.0096%	9.1171 \pm 0.0095%	0.09413
$\tau^- \rightarrow \nu_\tau \pi^+ \pi^- \pi^-$	8.9837 \pm 0.0095%	8.8828 \pm 0.0094%	0.09368
$\tau^- \rightarrow \nu_\tau \pi^0 \pi^+ \pi^- \pi^-$	4.2973 \pm 0.0066%	4.5319 \pm 0.0067%	0.30310
$\tau^- \rightarrow \nu_\tau \pi^0 \pi^0 \pi^0 \pi^-$	1.0765 \pm 0.0033%	1.0090 \pm 0.0032%	0.00724
$\tau^- \rightarrow \nu_\tau K^-$	0.7202 \pm 0.0027%	0.7138 \pm 0.0027%	0.00000
$\tau^- \rightarrow \nu_\tau \pi^0 \pi^0 \pi^+ \pi^- \pi^-$	0.4990 \pm 0.0022%	0.0897 \pm 0.0009%	0.00000
$\tau^- \rightarrow \nu_\tau \pi^0 K^-$	0.4785 \pm 0.0022%	0.4611 \pm 0.0021%	0.00000
$\tau^- \rightarrow \nu_\tau K_L^0 \pi^-$	0.4624 \pm 0.0022%	0.4444 \pm 0.0021%	0.00000
$\tau^- \rightarrow \nu_\tau \pi^- K_S^0$	0.4610 \pm 0.0021%	0.4449 \pm 0.0021%	0.00000
$\tau^- \rightarrow \nu_\tau \pi^+ \pi^- K^-$	0.3902 \pm 0.0020%	0.5051 \pm 0.0022%	0.52330
$\tau^- \rightarrow \nu_\tau \pi^0 \pi^- \eta$	0.1707 \pm 0.0013%	0.1696 \pm 0.0013%	0.00000
$\tau^- \rightarrow \nu_\tau \pi^- K^+ K^-$	0.1704 \pm 0.0013%	0.1509 \pm 0.0012%	0.07360
$\tau^- \rightarrow \nu_\tau \pi^0 K_L^0 \pi^-$	0.1605 \pm 0.0013%	0.2745 \pm 0.0017%	0.92850
$\tau^- \rightarrow \nu_\tau \pi^0 \pi^- K_S^0$	0.1592 \pm 0.0013%	0.2734 \pm 0.0017%	0.93657
$\tau^- \rightarrow \nu_\tau \gamma \pi^0 \pi^-$	0.1559 \pm 0.0012%	0.1303 \pm 0.0011%	0.00000
$\tau^- \rightarrow \nu_\tau K_L^0 \pi^- K_S^0$	0.1510 \pm 0.0012%	0.0763 \pm 0.0009%	0.00270
$\tau^- \rightarrow \nu_\tau K_L^0 K^-$	0.1289 \pm 0.0011%	0.0508 \pm 0.0007%	0.00000
$\tau^- \rightarrow \nu_\tau K_S^0 K^-$	0.1287 \pm 0.0011%	0.0507 \pm 0.0007%	0.00000
$\tau^- \rightarrow \nu_\tau \pi^0 \pi^0 \pi^0 \pi^+ \pi^- \pi^-$	0.1094 \pm 0.0010%	0.0506 \pm 0.0007%	0.00000
$\tau^- \rightarrow \nu_\tau \pi^+ \pi^+ \pi^- \pi^- \pi^-$	0.0803 \pm 0.0009%	0.0401 \pm 0.0006%	0.00000
$\tau^- \rightarrow \nu_\tau \pi^0 \pi^0 K^-$	0.0792 \pm 0.0009%	0.0504 \pm 0.0007%	0.2919
$\tau^- \rightarrow \nu_\tau K_L^0 K_L^0 \pi^-$	0.0760 \pm 0.0009%	0.0372 \pm 0.0006%	0.00854
$\tau^- \rightarrow \nu_\tau \pi^- K_S^0 K_S^0$	0.0756 \pm 0.0009%	0.0378 \pm 0.0006%	0.01189
$\tau^- \rightarrow \nu_\tau \pi^0 K_L^0 K^-$	0.0507 \pm 0.0007%	0.0763 \pm 0.0009%	0.85321
$\tau^- \rightarrow \nu_\tau \pi^0 K_S^0 K^-$	0.0498 \pm 0.0007%	0.0746 \pm 0.0009%	0.87506
$\tau^- \rightarrow \nu_\tau \pi^0 \pi^+ \pi^+ \pi^- \pi^- \pi^-$	0.0186 \pm 0.0004%	0.0293 \pm 0.0005%	0.00000

Similarity coefficients: T1=1.881148, T2=4.510389

• Process (decay channel)

• Branching ratio for generator #1 and #2


• Rough statistical errors of branching ratios

• Maximal “Shape Difference Parameter”

• Similarity Coefficients (combined: for all decay modes)

Example of histogrammes:

Comparison of Mass(1) of nu_tau pi- pi- in channel tau- => nu_tau pi0 pi+ pi- pi-




- Histogrammes of invariant mass from generator

#1 and #2

- Ratio of the two histogrammes

- **Shape Difference Parameter** value

HEPEvent library: a unified interface to event record formats



MC-TESTER as librarian tool:

- Automated comparison tests for changing versions of code
- Differences (e.g. new features) visible at a glance!
- Verification of compatibility with other version
- Example: **TAUOLA** generator
 - 3 versions of code to be managed:
CPC, CLEO, ALEPH

MC-TESTER as debugging tool:


- Detection of integrity errors in generator systems:
 - Event gramatics problems
 - cannot interpret data – reports the cause in debug mode
 - Configuration clashes and overwrites of private data:
 - Detects places where one generators “overwrite” the internal data of the other -> differences in distributions (example: BaBar)
 - Invariant mass distribution – good variable for finding phase-space inefficiencies

Event Record debugging:

- Support for various formats of event record:
HEPEvent library: easily extensible!
- Compare the same events stored in various formats of event records
- Debug event persistency methods and event record I/O

Event records over pipes...

In general, Monte Carlo modules coded in different languages (F77,F90,C++, Java,Perl...)



- Hard to link single main program (needs wrappers)
- One may export an event to a file in one module (executable), then read it in another
- Possibly large executable files
 - why not use **UNIX FIFO** queues instead?
- Synchronized parallelism!
- **MC-TESTER** example: reading output text files from MadGraph ME generator

Requirements:

- We support **Linux** (RedHat 6/7/8/9/10beta) on i386
(Tested also on AMD64)
- gcc 2.95/2.96/3.x: gcc,g++,g77,make
- root 3.X
- latex, (+dvips,gv)
- Unweighted events to be stored in one of supported event records (or text file)
- Work in progress:
 - Support for weighted events
 - support for HEPMC C++ event record
 - Support for MacOS X

Status and availability:

Project's homepage:

`http://cern.ch/Piotr.Golonka/MC/MC-TESTER`

- Current version: 1.112, 1.1p1 (src on the web)
- Documentation: (on the web)
CERN, LANL, LC preprints, to be published in CPC
- Contact:

`Piotr.Golonka@CERN.CH`

`Zbigniew.Was@CERN.CH`

`tomek@friend.phys.us.edu.pl`

We are open to discussions about:

- possible extensions and areas of use
- event record formats to be supported
(F77,C++,...)
- hardware/software platforms to be supported

Please, contact the authors.