

Peer review in the era of LHC experiments

Experimental particle physics as a
Big Science paradigm

QuickTime™ and a
TIFF (Uncompressed) decompressor
are needed to see this picture.

Rüdiger Voss
Physics Department
CERN, Geneva, Switzerland

OAI3@CERN, 12-14th February 2004

QuickTime™ and a
TIFF (Uncompressed) decompressor
are needed to see this picture.



Charting experimental particle physics

- Strongly centralised in a few big accelerator labs (CERN, Fermilab, DESY, SLAC, KEK ...)
- Increasingly concentrating on few very big projects:
 - CERN: LHC (4 experiments, under construction)
 - Fermilab: Tevatron (2 experiments)
 - DESY: HERA (4 experiments) (until end-2006?)
 - SLAC: PEP-II (BaBar)
 - KEK: KEK-B (BELLE)
- N.B. Astroparticle physics not discussed here



Charting.... (continued)

- The community:
 - \approx 10 000 scientists worldwide
 - \approx 50% Europe
 - \approx 50% rest of the world (US, Russia, Japan)
- \approx 80-90% work on the 'big' projects
- Corollary: CERN has a base of 6500 registered scientific users




The LHC project





LHC experiments in ex-LEP tunnel






The ATLAS example

- Multi-purpose detector for the LHC
- 2000 physicists, 150 institutes, 34 countries
- 500 MCHF investment
- Preparation since \approx 1990, operation starts 2007
- Estimated lifetime: 10 – 20 years
- Unprecedented technical complexity of
 - Hardware
 - Software
 - Data analysis



The ATLAS detector





Q: Can traditional peer review cope?

- Since era of LEP/HERA/Tevatron detectors (≈ 15 years ago), experiments have grown too complex to be mastered by the single scientist
- The LHC example: ATLAS/CMS will each generate a raw data flow \approx today's world throughput in telecommunications:
 - Imagine the real-time data processing challenge
 - Difficult not to make mistakes....
- Technical correctness of design, operation and analysis difficult (impossible?) to assess by classical peer review



A: Internal review by collaboration

- Strictly regulated multi-step process:
 1. Papers written by (small) 'Editorial Board'
 2. Review by 'Publication Committee' (PC) (non-anonymous)
 3. Draft made 'public' inside collaboration for comments
- Iterate steps 1-3 (sometimes restrictively) until PC decrees convergence
- Based on 'open archive local to collaboration'
 - Essential for efficient and transparent management of authoring and refereeing process
- 'Formal peer review with subsequent commentary'
- Successfully implemented by LEP and other major, non-CERN collaborations



Role of traditional peer review?

- Final publication still mostly in ‘conventional’ refereed journals
- Difficult to scratch deeper than the surface
- Largely reduced to rubber-stamping exercise, but still important and useful:
 - Formal/editorial aspects
 - Phrasing (conclusions!)
 - Interpretation & integration of final results in wider scientific context
- Minor revisions (at most)
- Strong self-selection of journals by authors
- ~0% rejection rate



Benefits of two-stage refereeing

- No wrong results known to data
 - but don't confuse with publication of 'effects' or 'particles' from statistical fluctuations
- Strong protection against scientific fraud!
 - Not due to formal review of publications only – large dispersed collaborations with flat hierarchies exercise informal but efficient self-control at many levels and all stages of the experiment



'Organized redundancy'

- Ultimately, quality assurance in particle physics is enforced by 'organized redundancy': build 2, 3, ... detectors to pursue same/similar scientific goals with
 - Different/complementary technologies
 - Different people
- First large-scale policy implementation with UA1/UA2 experiments at CERN (\approx 1975!)
- Climax in LEP programme (1989-2000): 4 detectors
- So far... redundant???
- Don't confuse with (friendly) scientific competition



Any useful conclusions?

- Experimental particle physicists have successfully implemented and operated for 15 years now a two-stage scheme of peer review that works
- It even works well! (judging by the results)
- Can it be mapped onto other disciplines?
 - Critical mass (> 100 scientists?)
 - Common project/facility
 - Flat hierarchies – scientific independence of sub-groups and individuals