

# Comparisons of hadronic shower packages

G.Mavromanolakis, University of Cambridge



---

## Outline

- ▶ **General**
- ▶ **Brief description**
- ▶ **Results**
- ▶ **Conclusions**

# General

- ▶ . simulation studies focused on CALICE ECAL-HCAL prototypes, to support and guide the testbeam program
  
- ▶ . survey of hadronic models in **GEANT3** and **GEANT4**
  - ▷ **GEANT3.21** : GHEISHA : GHEISHA SLAC  
: FLUKA + GHEISHA : GCALOR  
: FLUKA + MICAP
  
  - ▷ **GEANT4.6.0\*\*\*** : LHEP : QGSP  
: LHEP-BERT : QGSP-BERT  
: LHEP-BIC : QGSP-BIC  
: LHEP-GN : QGSC  
: LHEP-HP : FTFP

\*\*\* with hadronic physics list PACK 2.3

---


<b>model tag</b>	<b>brief description</b>
<b>G3-GHEISHA</b>	: <b>GHEISHA</b>
<b>G3-FLUKA+GH</b>	: <b>FLUKA, for neutrons with <math>E &lt; 20</math> MeV GHEISHA</b>
<b>G3-FLUKA+MI</b>	: <b>FLUKA, for neutrons with <math>E &lt; 20</math> MeV MICAP</b>
<b>G3-GH SLAC</b>	: <b>GHEISHA with some bug fixes from SLAC</b>
<b>G3-GCALOR</b>	: <b><math>E &lt; 3</math> GeV Bertini cascade, <math>3 &lt; E &lt; 10</math> GeV hybrid Bertini, FLUKA, <math>E &gt; 10</math> GeV FLUKA for neutrons with <math>E &lt; 20</math> MeV MICAP</b>
<b>G4-LHEP</b>	: <b>GHEISHA ported from GEANT3</b>
<b>G4-LHEP-BERT</b>	: <b><math>E &lt; 3</math> GeV Bertini cascade, <math>E &gt; 3</math> GeV GHEISHA</b>
<b>G4-LHEP-BIC</b>	: <b><math>E &lt; 3</math> GeV Binary cascade, <math>E &gt; 3</math> GeV GHEISHA</b>
<b>G4-LHEP-GN</b>	: <b>GHEISHA + gamma nuclear processes</b>
<b>G4-LHEP-HP</b>	: <b>as G4-LHEP, for neutrons with <math>E &lt; 20</math> MeV use evaluated cross-section data</b>
<b>G4-QGSP</b>	: <b><math>E &lt; 25</math> GeV GHEISHA, <math>E &gt; 25</math> GeV quark-gluon string model</b>
<b>G4-QGSP-BERT</b>	: <b><math>E &lt; 3</math> GeV Bertini cascade, <math>3 &lt; E &lt; 25</math> GeV GHEISHA, <math>E &gt; 25</math> GeV quark-gluon string model</b>
<b>G4-QGSP-BIC</b>	: <b><math>E &lt; 3</math> GeV Binary cascade, <math>3 &lt; E &lt; 25</math> GeV GHEISHA, <math>E &gt; 25</math> GeV quark-gluon string model</b>
<b>G4-FTFP</b>	: <b><math>E &lt; 25</math> GeV GHEISHA, <math>E &gt; 25</math> GeV quark-gluon string model with fragmentation ala FRITJOF</b>
<b>G4-QGSC</b>	: <b><math>E &lt; 25</math> GeV GHEISHA, <math>E &gt; 25</math> GeV quark-gluon string model</b>

---


# Brief description

- ▶ . run under Mokka(v2.2) and Brahms(v3.01) frameworks (GEANT4.6.0 and GEANT3.21 based, respectively)
- ▶ . study with W/Si ECAL+Fe/Scint or RPC HCAL CALICE prototypes porting detector geometry from GEANT4 to GEANT3
- ▶ . cutoffs : GEANT3 energy cutoffs EM = 10 keV, HAD = 0.1 MeV  
: GEANT4 range cutoff = 5  $\mu\text{m}$   
: ECAL, HCAL cellsize  $1 \times 1 \text{ cm}^2$ , threshold = 0.5 mip
- ▶ . samples of 10000 events, results normalised to G4-LHEP case (Mokka default), shown  $\pm 10\%$  and  $\pm 20\%$  bands wrt 1 to guide the eye

## GEANT3



## GEANT4



**ECAL** 30 layers  $\times$  50 cm  $\times$  38 cm interleaved with 0.5 mm Si pads

- ▷ W absorber, 10+10+10 layers, 1.4 mm:2.8 mm:4.2 mm thick per respective layer
- ▷ readout by 1 cm<sup>2</sup> cells





**HCAL** 40 layers  $\times$  100 cm  $\times$  100 cm interleaved with 6.5 mm scintillator  
or 1.2 mm RPCgas (digital HCAL)

- ▷ Fe absorber, 18 mm thick per layer
- ▷ readout by 1 cm<sup>2</sup> cells

# "response" vs model, $\pi^-$ 10 GeV/c

N cells hit

E deposited







△ different models predict different calorimeter response

△ HCAL more sensitive than ECAL

# "response" vs model, $\pi^-$ 1 GeV/c

N cells hit

E deposited







△ same pattern as at 10 GeV case, even more pronounced

△ ECAL standalone may have some discriminating power

# "response" rms/mean vs model, $\pi^-$ 10 GeV/c

## rms N/mean N




## rms Edep/mean Edep



$\blacktriangledown$  differences are still present but they are smaller

# shower transverse width vs model, $\pi^-$ 10 GeV/c

weighted by Edep per cell



△ models agree within  $\pm 5\%$  for mean energy deposited/cell

△ different calorimeter response per model is largely because of predicting different shower size



# Conclusions

- ▶ .
  - : all GHEISHA based models agree within  $\pm 20\%$  at 10 GeV,  
 $\pm 30\%$  at 1 GeV
  - : G3-FLUKA based models are definitely different
  - : low energy neutrons are important especially for the HCAL scint  
(e.g. compare G3-FLUKA+GH, G3-FLUKA+MI, G3-GCALOR)
  - : intranuclear cascade models are also important  
(e.g. compare Bertini or Binary cascade models with the rest)
- ▶ .
  - : ECAL standalone may have some discriminating power at low energies
- ▶ .
  - : different models predict different calorimeter response
    - ▷ mainly as a consequence of predicting different shower size
  - thus, different models predict different optimum calorimeter granularity

# Reminder

- ▶ : LHC experiments conducted a wide program of testbeams to validate hadronic shower simulation
  - : in general, ... the level of agreement between simulation and data is close to requirements in some cases (e.g.  $e/\pi$  ratio) whereas more work is needed for other cases (e.g. shower profiles)
- ▶ . **please note**
  - : LHC testbeams at  $E_{hadron} > 20$  GeV
  - : LC's region of interest is mainly in  $E_{hadron} < 20$  GeV
- ▶ . **so**
  - : suitable data desperately needed
  - : for the time being predictions are expected to have low reliability and low persistency