


# Top-Higgs Yukawa coupling measurement at TESLA

A. Gay, A. Besson, M. Winter

- Principle of the coupling measurement
- Background and tools
- Channels studied and results
- 6 fermions background
- Extensions of the analysis

# Principle of the coupling measurement

## t-tbar-H: diagrams, $\sigma$ and Br




$$\Rightarrow \sigma_{t\bar{t}H} \propto g_{t\bar{t}H}^2$$

## Uncertainty

$$\left(\frac{\Delta g_{t\bar{t}H}}{g_{t\bar{t}H}}\right)_{stat} \approx \frac{1}{S_{stat}(g_{t\bar{t}H}^2)\sqrt{\epsilon\rho L}} \text{ with } S_{stat}(g_{t\bar{t}H}^2) = \frac{1}{\sqrt{\sigma_{t\bar{t}H}}} \left| \frac{d\sigma_{t\bar{t}H}}{dg_{t\bar{t}H}^2} \right| \approx \frac{\sqrt{\sigma_{t\bar{t}H}}}{g_{t\bar{t}H}^2}$$

$$\left(\frac{\Delta g_{t\bar{t}H}}{g_{t\bar{t}H}}\right)_{syst} \approx \frac{1}{S_{syst}(g_{t\bar{t}H}^2)} \frac{1-\rho}{\rho} \frac{\Delta\sigma_{BG}^{ff}}{\sigma_{BG}^{ff}} \text{ with } S_{syst}(g_{t\bar{t}H}^2) = \frac{1}{\sigma_{t\bar{t}H}} \left| \frac{d\sigma_{t\bar{t}H}}{dg_{t\bar{t}H}^2} \right| \approx \frac{1}{g_{t\bar{t}H}^2}$$



# Background and tools

## Background:

Final State	$\sigma$ (fb)
q-qbar (u,d,s,c,b)	$\sim 1600$
t-tbar	$\sim 300$
W+W-	$\sim 4300$
ZZ	$\sim 240$
t-tbar-Z	$\sim 4.3$

## Generation/simulation

- t-tbar-H and t-tbar-Z generated with COMPHEP v41.10 + Pythia 6.158
- Other Bckgds generated with Pythia
- Simulated with SIMDET v4.

## B-tag

- NN based, including Zvtop, impact parameter joint probability tag, mass and momentum of the vertex, etc.
  - Implemented in SIMDET by T.Kuhl

## $t\bar{t}H; (H \rightarrow b\bar{b})$

– Semi-leptonic channel  $\rightarrow 4b2qlv$

➤  $BR(tt\bar{t}H \rightarrow 4b2qlv) \approx 43.9\% \times BR(H \rightarrow b\bar{b})$

– Hadronic channel  $\rightarrow 4b4q$

➤  $BR(tt\bar{t}H \rightarrow 4b4q) \approx 45.6\% \times BR(H \rightarrow b\bar{b})$

## $t\bar{t}H; (H \rightarrow W^+W^-)$

– 2 like sign leptons + 6 jets channel  $\rightarrow 2l^{+/-}2\nu4q2b$



➤  $BR(tt\bar{t}H \rightarrow 2l^{+/-}2\nu4q2b) \approx 10\% \times BR(H \rightarrow W^+W^-)$

– 1 lepton + 8 jets channel  $\rightarrow lv6q2b$

➤  $BR(tt\bar{t}H \rightarrow lv6q2b) \approx 40\% \times BR(H \rightarrow W^+W^-)$


## Neural Net selection

➤ Thrust, number of jets, Fox-Wolfram M., b-tag prob., etc.




# Selection ( $h \rightarrow b \bar{b}$ , semi leptonic channel)

before preselection




Fox-Wolfram moment  $h_{10}$


Fox-Wolfram moment  $h_{20}$




Fox-Wolfram moment  $h_{30}$



Fox-Wolfram moment  $h_{40}$





Minimum jet multiplicity




$\sum_{i=1}^4 p_{Ti}^2$

N.N. output



# Results

- **H → bb semlep;  $\Delta\sigma_{BG}^{eff}/\sigma_{BG}^{eff} = 5\%$**
- **H → bb semlep;  $\Delta\sigma_{BG}^{eff}/\sigma_{BG}^{eff} = 10\%$**
- **H → bb hadro;  $\Delta\sigma_{BG}^{eff}/\sigma_{BG}^{eff} = 5\%$**
- **H → bb hadro;  $\Delta\sigma_{BG}^{eff}/\sigma_{BG}^{eff} = 10\%$**
- **H → WW 2 like sign lep;  $\Delta\sigma_{BG}^{eff}/\sigma_{BG}^{eff} = 5\%$**
- **H → WW 2 like sign lep;  $\Delta\sigma_{BG}^{eff}/\sigma_{BG}^{eff} = 10\%$**
- **H → WW 1 lep;  $\Delta\sigma_{BG}^{eff}/\sigma_{BG}^{eff} = 5\%$**
- **H → WW 1 lep;  $\Delta\sigma_{BG}^{eff}/\sigma_{BG}^{eff} = 10\%$**
- **4 channels combined;  $\Delta\sigma_{BG}^{eff}/\sigma_{BG}^{eff} = 5\%$**
- **4 channels combined;  $\Delta\sigma_{BG}^{eff}/\sigma_{BG}^{eff} = 10\%$**



relative uncertainty on the residual background normalisation : 5% or 10 %

# 6 fermions background (1)

Previous results don't include 6 fermions processes

Cross-sections / generation with WHIZARD

- At the partonic level
- Include ISR and beamstrahlung
- Hadronization and FSR done by PYTHIA

Some processes receive contributions of resonant diagrams already taken into account with

- $t\text{-}\bar{t}$ ,  $W+W^-$ ,  $ZZ$ ,  $t\text{-}\bar{t}\text{-}Z$ ,  $t\text{-}\bar{t}\text{-}H$ 
  - In this case these contributions are subtracted from the total cross-section to avoid double counting

Cross-section calculation and generation

- Difficult and time consuming
- Ambiguities arise when parton pairs are defined for fragmentation
  - Loss of resolution on the  $g_{t\bar{t}H}$  measurement due to 6 f. background is an estimate

# 6 fermions background (2)

Hadronic channel ( $m_H = 120 \text{ GeV}/c^2$ )

Final state	$\sigma(fb)$	$\epsilon_{sel}$
$\bar{b}\bar{b}\bar{b}\bar{b}\bar{b}\bar{b}$	$6.4 \cdot 10^{-3}$	5.8%
$\bar{b}\bar{b}\bar{b}\bar{b}q\bar{q}$ ( $q = u, d$ )	$1.2 \cdot 10^{-1}$	1.6%
$\bar{b}\bar{b}\bar{b}\bar{s}\bar{s}$	$6.3 \cdot 10^{-2}$	1.8%
$\bar{b}\bar{b}\bar{b}\bar{c}\bar{c}$	$5.1 \cdot 10^{-2}$	2.0%
$\bar{b}\bar{b}\bar{b}\bar{t}\bar{t}$	$6.9 \cdot 10^{-3}$	31.8%
$\bar{b}\bar{b}q\bar{q}\bar{t}\bar{t}$ ( $q = u, d, s$ )	$8.9 \cdot 10^{-3}$	12.5%
$\bar{b}\bar{b}\bar{c}\bar{c}\bar{t}\bar{t}$	$3.5 \cdot 10^{-3}$	17.2%
$\bar{b}\bar{b}\bar{t}\bar{b}\bar{d}\bar{u}$ *	$\sim 1 \cdot 10^{-2}$	14.0%
$\bar{b}\bar{b}\bar{t}\bar{b}\bar{\mu}\bar{\nu}$ *	$\sim 3 \cdot 10^{-3}$	1.2%

$$\Delta\sigma_{BG}^{\text{eff}} / \sigma_{BG}^{\text{eff}} = 5 \%$$



# 6 fermions background (3)

Loss of resolution in the hadronic channel ( $m_H = 120 \text{ GeV}/c^2$ )

$\frac{\Delta\sigma_{BG}^{eff}}{\sigma_{BG}^{eff}}$	$\frac{\Delta g_{ttH}}{g_{ttH}}$ (without 6f)	$\frac{\Delta g_{ttH}}{g_{ttH}}$ (with 6f)
5%	8.3%	$\sim 8.5\%$
10%	10.1%	$\sim 10.5\%$

Loss of resolution in the semileptonic channel

$\frac{\Delta\sigma_{BG}^{eff}}{\sigma_{BG}^{eff}}$	$\frac{\Delta g_{ttH}}{g_{ttH}}$ (without 6f)	$\frac{\Delta g_{ttH}}{g_{ttH}}$ (with 6f)
5%	9.1%	$\sim 9.3\%$
10%	11.7%	$\sim 12.1\%$

➤ No significant effect


## New top mass from TeVatron:

–  $m_t = 178.0 \pm 4.3 \text{ GeV}/c^2$

- Changes the cross-sections to less than  $\sim 1\%$
- Higgs mass limit  $\sim 250/340 \text{ GeV}/c^2$  @ 95/99% c.l.
- Extension of the analysis to  $m_H \simeq 300 \text{ GeV}/c^2$

## Include the ( $H \rightarrow ZZ$ ) channel

- $26\% \leq \text{Br} \leq 30\%$  for  $(200 \leq m_H \leq 300 \text{ GeV}/c^2)$
- 2 leptons + 8 jets channel  $\rightarrow l^+l^-6q2b$



# Conclusion

@ the LC, the precision of the measurement of the top Yukawa Coupling will be better than  $\sim 10\%$

if  $m_H \leq 190 \text{ GeV}/c^2$

For a light Higgs ( $m_H \simeq 120 \text{ GeV}/c^2$ )

– Precision  $\sim 5\text{-}6\%$

First 6-fermions background simulations affect only marginally the precision of the measurement

The extension of the analysis is in progress (high masses, HZZ channel)

# 6 fermions background (4)

2 like sign lep. + 6 jets channel ( $m_H = 150 \text{ GeV}/c^2$ )

Final state	$\sigma(fb)$	$\epsilon_{sel}$
$b\bar{b}u\bar{d}e^-\bar{\nu}_e$ *	$\sim 1.5$	$8.4 \cdot 10^{-3}\%$
$b\bar{b}t\bar{b}d\bar{u}$ *	$\sim 6 \cdot 10^{-3}$	$< 0.2\%$
$b\bar{b}t\bar{b}\mu\bar{\nu}$ *	$\sim 1.5 \cdot 10^{-3}$	$0.4\%$
$t\bar{t}q\bar{q}q'\bar{q}'$ ( $q, q' = u, d, s$ )	$1.2 \cdot 10^{-2}$	$0.2\%$
$q\bar{q}t\bar{b}l\bar{\nu}_l$ ( $q = u, d, s, c; l = e^-, \mu^-$ ) *	$\sim 7.6 \cdot 10^{-2}$	$0.2\%$
$q\bar{q}t\bar{b}d\bar{u}$ ( $q = u, d, s, c$ ) *	$\sim 1.3 \cdot 10^{-1}$	$0.05\%$

\* Contribution of resonant reactions was subtracted

The selection efficiency is shown for  $\frac{\Delta\sigma_{BG}^{cff}}{\sigma_{BG}^{cff}} = 5\%$

$\frac{\Delta\sigma_{BG}^{cff}}{\sigma_{BG}^{cff}}$	$\frac{\Delta g_{ttH}}{g_{ttH}}$ (without 6f)	$\frac{\Delta g_{ttH}}{g_{ttH}}$ (with 6f)
5%	13.6%	$\sim 13.9\%$
10%	14.2%	$\sim 14.5\%$

# 6 fermions background (5)

1 lep. + 8jets channel ( $m_H = 150 \text{ GeV}/c^2$ )

Final state	$\sigma(fb)$	$\epsilon_{sel}$
$b\bar{b}u\bar{d}e^-\bar{\nu}_e$ *	$\sim 1.5$	0.04%
$b\bar{b}t\bar{b}d\bar{u}$ *	$\sim 6 \cdot 10^{-3}$	2.4%
$b\bar{b}t\bar{b}\mu\bar{\nu}$ *	$\sim 1.5 \cdot 10^{-3}$	3.4%
$t\bar{t}q\bar{q}q'\bar{q}'$ ( $q, q' = u, d, s$ )	$1.2 \cdot 10^{-2}$	7.6%
$q\bar{q}t\bar{b}l\bar{\nu}$ ( $q = u, d, s, c; l = e^-, \mu^-$ ) *	$\sim 7.6 \cdot 10^{-2}$	1.7%
$q\bar{q}t\bar{b}d\bar{u}$ ( $q = u, d, s, c$ ) *	$\sim 1.3 \cdot 10^{-1}$	1.2%

\* Contribution of resonant reactions was subtracted

The selection efficiency is shown for  $\frac{\Delta\sigma_{BG}^{eff}}{\sigma_{BG}^{eff}} = 5\%$

$\frac{\Delta\sigma_{BG}^{eff}}{\sigma_{BG}^{eff}}$	$\frac{\Delta g_{ttH}}{g_{ttH}}$ (without 6f)	$\frac{\Delta g_{ttH}}{g_{ttH}}$ (with 6f)
5%	13.5%	$\sim 14\%$
10%	16.8%	$\sim 17.5\%$