

Astroparticle Physics (3/3)

Nathalie PALANQUE-DELABROUILLE
CEA-Saclay

CERN Summer Student Lectures, August 2004

- 1) What is Astroparticle Physics ?
Big Bang Nucleosynthesis
Cosmic Microwave Background
- 2) Dark matter, dark energy
- 3) High energy astrophysics
Cosmic rays
Gamma rays
Neutrino astronomy



Lecture outline

- 1) What is Astroparticle Physics ?
Big Bang Nucleosynthesis
Cosmic Microwave Background
- 2) Dark matter, dark energy
- 3) High energy astrophysics
Cosmic rays
Gamma rays
Neutrino astronomy



Brief Cosmic Ray history

1912 Hess discovers cosmic rays




1925 Quasi-isotropy

1938 Auger discovered extensive air showers
($E = 10^{15}$ eV!)


1946 First air shower experiment



Energy spectrum



Structure in cosmic ray spectrum





$$\frac{dN}{dE} = E^{-\alpha} \quad \text{above } 10 \text{ GeV}$$

Source acceleration
 $2.0 - 2.2$

Propagation
 ~ 0.6

GZK (Greisen Zatsepin Kuzmin) cut-off



When process energetically allowed ($>5 \times 10^{19}$ eV), space becomes **opaque** to CR


Sources with $E > E_{\text{GZK}}$ must be at $d < 100$ Mpc (local cluster)

(no known acceleration sites...)

Acceleration mechanisms

1949 : Fermi acceleration

Stochastic acceleration of particles
on magnetic inhomogeneities



Head-on collisions \Rightarrow Energy gain
Tail-end collisions \Rightarrow Energy loss
On average, head-on more probable
 \Rightarrow Energy gain over many collisions


$$\Delta E/E \propto \beta^2 \quad \beta = v/c \diamond 10^{-4}$$

Slow and inefficient

“ Second order ”

First order Fermi acceleration

1970's : First order Fermi acceleration
 Acceleration in strong shock waves



Conservation of nb of particles :

$$\rho_1 v_1 = \rho_2 v_2$$

Strong shock : $\rho_2/\rho_1 = (\gamma+1)/(\gamma-1)$

Fully ionized plasma (\Leftrightarrow ideal gas)

$$\gamma = 5/3 \text{ and } v_1/v_2 = 4$$

\Rightarrow Rapid gain in energy as particles repeatedly cross shock front


$\Delta E/E \propto \beta (\sim 10^{-1})$ and E^{-2} spectrum

" First order "

Powerful shocks?



Supernovae !



Low mass star

High mass star

Supernova

(too short) life and
(extremely violent) death
of massive stars



SN 1987A
(SN II)



Crab supernova
remnant

1 SN II / 50 years in our galaxy

Energy limitation

Natural limit : containment of particles in acceleration (shock) region

$$E_{\max} \sim Z e B R c$$

(no energy losses)


→ Need high B, large R

Supernova remnants :

$$\rightarrow E_{\max} \sim 10^{15} \text{ eV (knee)}$$



Cosmic rays in $10^{15} - 10^{20}$ eV region ?

→ Relativistic motions (Γ)




Active Galactic Nuclei

- AGN : galaxy with $10^8 - 10^9 M_{\odot}$ central black hole
- 10% - radio jets (relativistic ejection of plasma)
 - 1% - blazars (all EGRET AGNs !)



Cosmic ray detectors



$E < 10^{14}$ eV
 Flux high enough for detection of primary particle
 (AMS on ISS)


$E > 10^{14}$ eV
 Use atmosphere to create extensive air showers
 (AGASA, Fly's eye)

Counting particles: AGASA

Trajectory determined from arrival time of shower front on ground detectors

Cerenkov detectors measure height of shower maximum width (X_{\max}) related to primary energy

130 km west of Tokio




Air fluorescence: Fly's Eye

Spherical mirrors viewed by
PMT's at the focal plane

Dual setup allows accurate
trajectory **reconstruction**


Amount of light (with $1/r^2$
correction for geometry)
→ shower profile
→ shower maximum X_{max}
→ **primary energy**



Can only operate on clear
and moonless nights


13 km apart in
Utah desert

Ultra High Energy Cosmic Rays



AGASA: 17 events above 6×10^{19} eV
 HiRes : 2 events (~ 20 expected)

Need cross calibration !



$E_{\text{max}} = 3.2 \cdot 10^{20} \text{ eV} = 50 \text{ J} !$


Puzzling facts

Ultra high energy protons are
not confined in galaxy

+
Isotropy



Extra-galactic sources



No counterpart (any wavelength)
Cosmological sources

→ Invisible source ?
→ No GZK cut-off ?

Possible suspects : | Local GRB's
| Exotica
| ...

Need more events

Future with AUGER and EUSO


AUGER

Air fluorescence + ground arrays
2 sites (Argentina, USA):
1600 detectors + 4 telescopes, 3000 km²
First results (though not all detectors)



EUSO

Air fluorescence from space
Expect 10^3 CR yr⁻¹ above GZK
Launch: 2010 (for 3 years)



Lecture outline

- 1) What is Astroparticle Physics ?
Big Bang Nucleosynthesis
Cosmic Microwave Background
- 2) Dark matter, dark energy
- 3) High energy astrophysics
Cosmic rays
Gamma rays
Neutrino astronomy




Gamma ray satellites

1972 ● SAS-2




1975 ● COS-B
70 MeV - 5 GeV

1991 ● G.R.O.-EGRET
50 MeV - 30 GeV


2006 ● GLAST
20 MeV - 300 GeV




21st century



EGRET ($E > 100 \text{ MeV}$)



Galactic diffuse interstellar emission from interaction with cosmic rays



Point sources

- Jets from active galactic nuclei
- Galactic sources in star-forming sites : pulsars, binaries, supernova remnants ...
- Unidentified sources (170/270)


Blazars

Low energy emission (X-ray) :
Synchrotron emission of e^- in jet

VARIABILITY !




Size $\sim \Gamma c t_{\text{var}}$
($\Gamma > 10$)



High energy emission (γ -ray):


- self-compton (electro-magnetic) ?
- π^0 decay (hadronic) ?

Markarian 421 : closest blazar




Quasars and Microquasars

QUASAR 3C 223




MICROQUASAR 1E1740.7-2942



QUASAR

MICROQUASAR



$$R \propto M_{BH}$$


$$T \propto M_{BH}^{-1/4}$$

Mirabel & Rodriguez

Gamma ray bursts (GRB)

1967 Chance discovery of prompt emission by VELA (16 events), published in 1973

1991 Observation with the satellites C.G.R.O (EGRET, BATSE...) & BeppoSAX





brightest objects in the universe, emitting mostly at high E
→ emission collimated ?
wide variety of time profiles, Δt from 10ms to 1000s
→ compact region, Lorentz boost ($\Gamma \sim 100$)


2002 (>2000 bursts) still very poorly understood ...

Burst location


Isotropy
+
Optical counterparts
↓
Cosmological phenomena!
($z = 0.43$ to 4.50)




Afterglow in optical



TeV sky



High energy cut-off !



GZK cutoff
Main explanation
for lack of
TeV sources

GZK cutoff for γ
but not for UHECR ?

Lecture outline

- 1) What is Astroparticle Physics ?
Big Bang Nucleosynthesis
Cosmic Microwave Background
- 2) Dark matter, dark energy
- 3) High energy astrophysics
Cosmic rays
Gamma rays
Neutrino astronomy



Other messengers ?

Photons: absorbed (GZK)

Neutrons: $\tau \sim 15 \text{ mn}$
 $d_{\text{max}} = 10 \text{ kpc}$ ($E = 10^{18} \text{ eV}$)


Protons: absorbed (GZK)
& deviated ($E < 10^{18} \text{ eV}$)

Neutrinos: no charge, "no"
interaction with matter
nor radiation

High energy sources

High energy emission (γ -ray):

- ~~self-compton (electro-magnetic) ?~~
- ~~π^0 decay (hadronic) ?~~



High energy
 ν sources !

Experimental challenge

Low fluxes @ high E
Low cross-sections




Large volume of detector
(lake, sea, polar ice)

High background
(atmospheric μ & ν)




Good shielding
(> 1000m water eq.)
Search for upgoing ν 's

Detection principles



- Cosmic ν (> 1 TeV)
- $\chi\chi \rightarrow \nu$ (10-1000 GeV)
- Atm. ν (10-100 GeV)
- Atm. μ

$\nu \rightarrow \mu \rightarrow \text{Cerenkov light}$




HE neutrino experiments



Detectors

Strings with optical modules (PMT in glass sphere)





- d_{OM-OM} : E threshold
- # of OM: E resolution
- $d_{string-string}$: effective volume, E limit

Sky coverage

ANTARES (43° North)

AMANDA (South pole)



ANTARES/AMANDA: 0.6π sr overlap

Conclusions

Cosmic Ray physics

Existence or not of post GZK cut-off events ?

Gamma Ray physics

Study of high energy sources (AGNs, blazars)

GRB mystery

Neutrino physics

Complementary to photon astrophysics
(models confrontations)

Indirect dark matter searches

New look on the Universe → room for unexpected discoveries