

HERA/LHC Workshop  
CERN, 11th October 2004


# Diffractive dijet photoproduction at ZEUS

ALESSIA BRUNI, INFN BOLOGNA


# Factorization tests in $\gamma p$

Real photon ( $Q^2 \simeq 0$ ) can develop hadronic structure  
⇒ study fact. breaking at HERA

direct  $\gamma$  ( $x_\gamma \simeq 1$ )




resolved  $\gamma$  (hadron like, at low  $x_\gamma$ )



in NLO, suppression of resolved contribution by a factor 3 expected  
(Klasen & Kramer)

# Kinematics



Photoproduction:  $Q^2 \simeq 0$

- $y$  fraction of longitudinal momentum of  $e$  taken by  $\gamma$
- $x_\gamma$  fraction of longitudinal momentum of  $\gamma$  in hard scattering
- $M_X$  hadronic mass from  $\gamma$  dissociation
- $z_{IP}$  fraction of longitudinal momentum of  $IP$  taken by parton
- $x_{IP}$  fraction of longitudinal momentum of  $p$  in diffr. exchange


# Event selection


- Data sample 99-00 data ( $E_e = 27.5 \text{ GeV}$ ,  $E_p = 920 \text{ GeV}$ )
- Integrated luminosity  $77.6 \text{ pb}^{-1}$

PHP selection	no $e$ in detector $0.20 < y_{JB} < 0.85$
Diffr. selection	rapidity gap of at least 3 units $x_{IP} < 0.035$
dijet selection	$\geq 2$ jets with $k_T$ algorithm in lab frame $E_T^{jet1(2)} > 7.5(6.5) \text{ GeV}$ , $-1.5 < \eta^{jet1,2} < 2.0$
$\Rightarrow 10673$ events are selected	

Background: p-diss events  $(16 \pm 4)\%$  subtracted  
non-diffractive events 10% non subtracted

# Event display: dijet in diffractive $\gamma p$






## Data vs MC LO

- MC RAPGAP (direct + resolved) describes shape of  $y, x_{IP}, E_T^{jet}, \eta_T^{jet}$

RAPGAP v3.00  
structure functions:  $p$  MRSG,  
 $\gamma$  GRV-G-LO,  $IP$  H1 fit2




## Data vs MC LO

- MC RAPGAP (direct + resolved) describes shape of hadronic mass  $M_x, z_{IP}, x_\gamma$
- ratio data/MC: no suppression of resolved component as described by RAPGAP


## Comparison with NLO predictions

- NLO calculations by Klasen and Kramer for partonic level
- corrections to hadron level estimated using RAPGAP MC




## Comparison with NLO predictions

- corrections to hadron level estimated using RAPGAP MC
- corrections in  $x_\gamma$  are not large



## Data vs NLO

- $y, x_{IP}, z_{IP}, E_T^{jet}, \eta_T^{jet}$
- shape described
- normalization not described, a suppression factor is needed



## Data vs NLO - $x_\gamma$

- normalization not described, a suppression factor is needed
- the same suppression factor seems to be applied to both resolved and direct components

# Diffractive Dijet in Photoproduction - Summary

- Cross sections measured by ZEUS
- Data described by LO MC RAPGAP
- NLO describe shape of data
- NLO need a suppression factor  $\sim 0.5$
- Data favour a suppression also of the direct  $\gamma$  component

Next steps:

- double differential cross sections
- ratio of diffractive/total cross section