

# Discussion Benchmark cross sections

# Motivation

Different theoretical approaches for heavy flavour cross-sections  
 each approach typically available both for HERA and LHC

An incomplete set of those with full differential prediction reported here:

Model	HERA $\gamma p$	HERA DIS	LHC
LO(massless)+PS	Pythia	Pythia, Lepto	Pythia
LO+PS+MI	Pythia/JIMMY		Pythia/JIMMY
LO(massive)+PS	Pythia	Rapgap, Aroma	Pythia
CCFM MC (Cascade)	X	X	X
NLO	FMNR	HVQDIS	HVQMNR
FONLL	X		X
NLO(massive)+PS	(MCaNLO ?)		MCaNLO
Massless (KK)	X		X
$K_T$ factorization			X
Enhancement at low-x		?	X
Saturation at Low-x Kutak			X

## Motivation 2

It would be nice to have in the workshop proceedings prediction from all these different models for a given set of observables:

- Direct way to compare features of different models and show regions of (un)compatibility
- show in a single place how the same approach works for HERA and LHC
- are two models equivalent for LHC but not for HERA and vice-versa ?
- Collect all available prediction in a “reference” book

To do this we need to:

- Define a standard set of observables, the “benchmark cross sections”, typically differential cross sections after some cut
- Ask theorists or in general who has access to the programs to calculate prediction for the different models
- predictions should have uncertainty bands or at least provide a set of curves for different (reasonable) parameter choices

# Which cross sections ?

Not too far from experimental observables

Not too complicate

i.e don't repeat exactly expt. cuts (as in JetWeb)

Basic observables such as  $d\sigma/dp_T$  in a given  $y$  range

Interesting observables to spot particular regions where models may differ

the number of cross sections should not be too large (  $< 10$  )

Quark or hadron cross sections or both ? (I think hadron)

# Cross Sections for LHC

Differential cross sections ( $d\sigma/dX$ , where X:)

Basic:

1.1)  $p_T$  central rapidity  $|y| < 2.5$  (ATLAS/CMS)

1.2)  $p_T$  large rapidity  $2.5 < |y| < 4.5$  (LHC-B)

1.3)  $y$  for  $p_T > 2$  GeV ? (LHC-B/Alic<sup>e</sup>)

need special Alic<sup>e</sup> range ?

different cuts for charm and b ?

Correlations:

observables sensitive to intrinsic  $k_T$  and g splitting:

1.4)  $p_T(Q + \bar{Q})$

1.5)  $\Delta\phi(Q - \bar{Q})$

observables sensitive to 4b production:

1.6)  $\Delta y(QQ)$



# Cross Sections for HERA ( $\gamma p$ )

Basic cross sections:

2.1)  $p_T$  for  $|y| < 1.5$ ,  $130 < W < 300\text{GeV}$ ,  $Q^2 < 1\text{GeV}^2$  (range of existing  $D^*$  measurements)

2.2)  $y$  for  $p_T > 2\text{ GeV}$ ,  $130 < W < 280\text{GeV}$ ,  $Q^2 < 1\text{GeV}^2$

More complex vars:

$x_\gamma$  is interesting but beyond a very simple implementation of Heavy Hadron cross sections

Correlations:

2.3)  $p_T(Q + \bar{Q})$  and/or

2.4)  $\Delta\phi(Q - \bar{Q})$

(in which range ?)

...

# Cross Sections for HERA (DIS)

Here it's more complex: many variables ( $x, Q^2$  and  $y, p_T, \phi$  of the HQ)

I see two possible choices:

a) plot  $y, p_T$  for ranges in  $x, Q^2$  ?

b) fix a range in  $y, p_T$  and plot  $x$ , for different  $Q^2$  ?

Option b) is probably better:

1.1-4)  $x$  for  $Q^2 = 1, 10, 100, 1000 \text{ GeV}^2, p_T > 1.5, |y| < 1.5$   
similar to “visible”  $F_2^c$

with or without EW corrections ? (without)



## Let's discuss and see ...

$14 \times 2(\mathbf{b+c})$  cross section proposed, too many ?

remove some, propose some new

fix standard PDFs,  $\alpha_S$ ,  $m_b$ ,  $m_c$  ?

find who does what !