




# Medium Effects on Jets and their Energy Dependence

**Henner Büsching**


**Brookhaven National Laboratory  
for the PHENIX Collaboration**

**Hard Probes - Ericeira , Nov. 8 2004**

# Outline



**p+p 200 GeV**  
Initial conditions



**Au+Au 200 GeV**  
Probe hot and dense nuclear  
matter with high  $p_T$  partons

**d+Au**  
**200 GeV**

**Au+Au**  
**62.4 GeV**


**Particle**  
**species**

# PHENIX at RHIC

2 central spectrometers

2 forward spectrometers



3 global detectors





*Initial Conditions*  
 *$p+p$  at 200 GeV*

# Do we understand our p+p data?



$$E \frac{d^3 \sigma}{d^3 p} = \frac{1}{p_T^n} F(x_T) = \frac{1}{\sqrt{s}^n} G(x_T)$$


$$x_T = 2 p_T / \sqrt{s}$$

$x_T$  scaling at high  $p_T$

# pQCD works – $\pi^0$

- Good agreement with NLO pQCD
  - Factorization theorem:

$$\sigma_{AB \Rightarrow hX} \propto f_{a/A}(x_a, Q_a^2) \otimes f_{b/B}(x_b, Q_b^2) \otimes \sigma_{ab \Rightarrow cd} \otimes D_{h/c}(z_c, Q_c^2)$$





- Constrains Fragmentation Function  $D(\text{Gluon-}\pi)$
- Reference for Au+Au spectra
- Especially good reference if measured together with heavy ion data : reduced syst. errors

Run3 : preliminary

Run2:  
Phys. Rev. Let 91, 241803 (2003)

# pQCD works – direct photons



# Nuclear Modification Factor $R_{AA}$

- **Hard processes**
  - yield scales with  $N_{coll}$
  - reason:
    - small cross section
    - incoherent superposition
- **Nuclear Modification Factor  $R_{AA}$**

$$R_{AB} = \frac{\left(1/N_{AB}^{evt}\right) d^2N_{AB}/dydp_T}{\langle T_{AB} \rangle d^2\sigma_{pp}/dydp_T}$$



$$\langle N_{coll} \rangle / \sigma_{NN}$$

**Geometrical factor from nuclear overlap**

**Nuclear overlap function**

$$T_{AB} = \int d^2\mathbf{r} T_A(\mathbf{r}) T_B(\mathbf{b} - \mathbf{r})$$

**Nuclear thickness function**

$$T_A(b) = \int dz \rho_A(b, z)$$

---

**from Glauber model**




# Nuclear Modification Factor $R_{AA}$

- **Hard processes**
  - yield scales with  $N_{\text{coll}}$
  - reason:
    - small cross section
    - incoherent superposition
- **Nuclear Modification Factor  $R_{AA}$**

$$R_{AB} = \frac{\left(1/N_{AB}^{\text{evt}}\right) d^2N_{AB}/dydp_T}{\langle T_{AB} \rangle d^2\sigma_{pp}/dydp_T}$$

# Nuclear Modification Factor $R_{AA}$

- **Hard processes**
  - yield scales with  $N_{\text{coll}}$
  - reason:
    - small cross section
    - incoherent superposition
- **Nuclear Modification Factor  $R_{AA}$**



$$R_{AB} = \frac{\left(1/N_{AB}^{\text{evt}}\right) d^2N_{AB}/dydp_T}{\langle T_{AB} \rangle d^2\sigma_{pp}/dydp_T}$$


- **In the absence of nuclear effects:  $R_{AA}=1$  at high  $p_T$**

# Nuclear Modification Factor $R_{AA}$

- **Hard processes**
  - yield scales with  $N_{\text{coll}}$
  - reason:
    - small cross section
    - incoherent superposition
- **Nuclear Modification Factor  $R_{AA}$**

$$R_{AB} = \frac{\left(1/N_{AB}^{\text{evt}}\right) d^2N_{AB}/dydp_T}{\langle T_{AB} \rangle d^2\sigma_{pp}/dydp_T}$$

**7.2 GeV muons  
on various targets  
scale as  $\alpha=1.0$  in  $\mu+A$**




**Phys. Rev. Lett. 35, 407 (1975)**

# Nuclear Modification Factor $R_{AA}$

- **Hard processes**
  - yield scales with  $N_{coll}$
  - reason:
    - small cross section
    - incoherent superposition
- **Nuclear Modification Factor  $R_{AA}$**

$$R_{AB} = \frac{\left(1/N_{AB}^{evt}\right) d^2N_{AB}/dydp_T}{\langle T_{AB} \rangle d^2\sigma_{pp}/dydp_T}$$



**Cronin effect:**


$$\alpha(p_T) > 1$$

**Multiple scattering of incident partons**



*AU+AU*  
*200 GeV*

# Binary scaling - direct photons





- Important test of QCD and initial state
  - Only interact electromagnetically
  - No interaction with medium
- Observation:
  - Pure  $N_{\text{Coll}}$  scaling relative to pQCD calculation



Talk by K. Reygers


# $\pi^0$ -Production



$N_{coll}$ -scaling works in peripheral Au+Au, but strong suppression in central Au+Au

Phys. Rev. Lett. 91, 072301 (2003)

# Nuclear Modification Factor $R_{AA}$



- RAA at higher  $p_T$  ?
- Run4 higher statistics



$$R_{AB} = \frac{(1/N_{AB}^{\text{evt}}) d^2 N_{AB}/dydp_T}{\langle T_{AB} \rangle d^2 \sigma_{pp}/dydp_T}$$

Phys. Rev. Lett. 91, 072301 (2003)




# Nuclear Modification Factor $R_{AA}$

- RAA at higher  $p_T$  ?
- Run4 higher statistics
- New run3 pp reference
- Reduced stat. errors
- Reach to higher  $p_T$




$$R_{AB} = \frac{\left(1/N_{AB}^{\text{evt}}\right) d^2N_{AB}/dydp_T}{\langle T_{AB} \rangle d^2\sigma_{pp}/dydp_T}$$

# $\eta$ Production




- $\pi^0$  and  $\eta$  suppression consistent in magnitude and  $p_T$  dependence
- Suppression of factor 5 in central collisions
- Agreement with parton energy loss predictions up to highest  $p_T$  measured so far



GLV  $R_{AA}$   
 $dN^g/dy = 1100$

charged hadrons less suppressed at medium  $p_T$

later in talk 



**C**  
**e**  
**n**  
**t**  
**r**  
**a**  
**i**  
**t**  
**y**




**D**  
**e**  
**p**  
**e**  
**n**  
**d**  
**e**  
**n**  
**c**  
**e**

Phys. Rev. Lett. 91,  
072301 (2003)


Phys. Rev. C 69,  
034910 (2004)

# Jet Quenching?

Comparison with model calculations with and without parton energy loss:




- $\pi^0$  suppression described by models with parton energy loss
- Additional nuclear effects needed to describe  $p_T$  dependence
- Other explanations not ruled out at this stage



# $x_T$ scaling in Au+Au

$$\sqrt{s}^n \times E \frac{d^3\sigma}{d^3p} = G(x_T)$$



- **If** high  $p_T$  particle production in Au+Au result of hard scattering:
  - $x_T$  scaling should work
- **Assumption: structure and fragmentation functions should scale**
- $n(x_T, \sqrt{s_{NN}})$  in AuAu:
  - Compare 130 GeV and 200 GeV data
  - Central and peripheral
- $\pi^0$  and peripheral charged particles scale
- Central charged particles do not scale





**Follow pQCD  
as in pp**



*d+Au*

*200 GeV*

# The Control Experiment: d+Au




Kharzeev, Levin, McLerran,  
hep-ph/0210332

- No suppression in d+Au
- Initial-state effects ruled out as explanation for suppression in Au+Au at mid rapidity

Phys. Rev. Lett. 91, 072303 (2003)

# $\pi^0$ 's in d+Au: Data vs. pQCD



- Data well reproduced by
  - NLO pQCD calculation, plus
  - Phenomenological model of Cronin-Effect, plus
  - Shadowing



Levai et al., nucl-th/0306019

Kopeliovich et al., Phys. Rev. Lett. 88, 232303 (2002)





# $R_{dAu}$ for different centralities

$\langle k_T \rangle = 0.52 \text{ GeV}^2$







Prediction for gluon saturation



Kharzeev, Levin, McLerran,  
hep-ph/0210332

$(N_{\text{part, Au}})^{1/2}$  scaling  
 $R_{AA} \approx 0.7$  in central d+Au


# $\eta$ Production in dAu




PHENIX  
Preliminary

# Centrality Dependence


Au + Au Experiment



d + Au Control Experiment



charged



- Different behavior with increasing centrality in Au+Au and d+Au
  - $R_{AuAu}$  decreases
  - $R_{dAu}$  charged increases
  - $R_{dAu} \pi^0$  increases slightly




*AU+AU*

---

*62.4 GeV*

# 62.4 GeV p+p $\pi^0$ Reference



- $\pi^0$  data from ISR
- Corrected for (when necessary)
  - hadronic decay
  - direct photons
- Global fit:


$$f(p_T) = A / (e^{a \cdot x^2 + b \cdot x} + x/p_0)^n$$

- Fit and data cross-checked by NLO calculation
- $\pm 25\%$  systematic uncertainty



Talk by D. d'Enterria

# $R_{AA}$ for $\pi^0$ at 62.4 GeV




$N_{coll}$  for peripheral events not yet understood


- Ratios close to unity at  $p_T \sim 2.5$  GeV/c, then decreasing
- Same tendency as 200 GeV Au+Au
- Less suppression at intermediate  $p_T$  (compared to 200 GeV)

# $R_{AA}$ for $\pi^0$ at 62.4 GeV: Predictions

I. Vitev nucl-th/0404052




Adil & Gyulassy nucl-th/0405036



$R_{AA}(\pi^0) \sim 0.5 - 0.3$

# $R_{AA}$ for $\pi^0$ at 62.4 GeV

Vitev  
nucl-th/0404052



- Reasonably good agreement (esp. high  $p_T$ ) within uncertainties
- Uncertainty in the  $p_T$  shape (esp. low  $p_T$ ) of p+p reference ...





# $R_{AA}$ for Charged Hadrons


- $\pi^0$  yield is multiplied by 1.6
- Discrepancy between charged and  $\pi^0$
- Pions more suppressed than  $h^{+/-}$  at intermediate  $p_T$
- Large proton contribution up to at least 4 GeV/c

62.4 GeV

charged reference  
is used here



# $\pi^0 R_{AA}$ , Central Events, Different $\sqrt{s}$



WA98, EPJ C 23, 225 (2002)

[ new reference compiled by D.d'Enterria nucl-ex/0403055]

PHENIX, PRL 88 022301 (2002)


- Suppression at 17 GeV and 62.4 GeV similar at medium  $p_T$
- 62.4 GeV expected lower
- Better reference needed!



# *Particle ratios*

---


# $\eta$ to $\pi^0$ Ratio



world average



- Flat  $\eta/\pi^0$  ratio as function of  $p_T$
- No centrality dependence
- $\eta/\pi^0$  ratio in AuAu consistent with world average

# Proton-Scaling




- Proton/anti-proton-yield **scales with  $N_{coll}$**  in the range  $2 \text{ GeV} < p_T < 3 \text{ GeV}$
- Why are protons/anti-protons not suppressed?

# $\rho/\pi$ – Ratio




- Expectation for particle production from jet-fragmentation:  $\rho/\pi$  less than  $\approx 0.25$  at high  $p_T$
- $\rho/\pi^+ > 1$  at intermediate  $p_T$
- less for antiproton ( $\bar{p}/\pi^- \sim 0.7$ ).
- Weaker centrality dependence than at 200 GeV

# Charged Hadron / $\pi^0$



- Unidentified charged hadrons can be measured up to  $p_T = 9$  GeV/c
- *Ratio above 5 GeV/c similar for*
  - *central Au+Au*
  - *peripheral Au+Au*
  - *p+p*
- *This implies that  $p/\pi$  ratio goes back to “normal” value at high  $p_T$*

# A Possible Explanation: Quark Recombination



Fries, et al, nucl-th/0301087

also, Greco, Ko, Levai, nucl-th/0301093

- Two competing processes for hadron production
  - Jet-Fragmentation in the vacuum
  - Recombination of 3 quarks or a quark/anti-quark pair in a densely populated phase space
- Fries et al.
  - In case of thermalized partons at RHIC, fragmentation wins over recombination only above  $p_T = 5$  GeV/c
  - This explains  $p/\pi$  ratio
  - “Such a phase phase may be appropriately called a QGP”



**Talk by J. Velkovska**



# Summary

- **p+p reference well under control at 200 GeV**
  - $x_T$  scaling works
  - binary scaling works for direct photons
- **suppression of  $\pi^0$ 's and charged hadrons in central Au+Au**
  - charged hadrons less suppressed at medium  $p_T$
  - $x_T$  scaling
    - works for pions:
      - hard scattering
    - breaks for charged hadrons in central Au+Au:
      - not only hard scattering
- **non-suppression in d+Au**
  - suppression in Au+Au must be final-state effect
- **suppression also in 62.4 GeV Au+Au**
  - evidence for smooth  $\sqrt{s}$  dependence
  - better reference needed to say more
- **high  $p/\pi$  ratio in central Au+Au**
  - possibly recombination from thermalized partons






*backup*

---

# PHENIX-Setup




(pseudorapidity  $|\eta| < 0.35$ )

- Relevant for this talk:
- Detectors in the central spectrometer arms
  - $\pi^0$  via  $\pi^0 \rightarrow \gamma\gamma$ :
    - Lead scintillator calorimeter (**PbSc**)
    - Lead glass calorimeter (**PbGl**)
- Centrality , vertex
  - Beam Beam Counter (**BBC**)  
 $3.0 < |\eta| < 3.9$
  - Zero Degree Calorimeter (**ZDC**)



# PHENIX Detector at Collision Point




East Carriage  
( Moved in Place )

Ring Imaging  
Cerenkov  
Drift Chamber

Beam-Beam Counter  
Central Magnet

West Carriage


# PHENIX in Run 4




|        | 100GeV/u<br>(mb) <sup>-1</sup> | Relative<br>to Run-2 | 31.2GeV/u<br>(mb) <sup>-1</sup> |
|--------|--------------------------------|----------------------|---------------------------------|
| PHENIX | 1370                           | 15x                  | 21.8                            |
| STAR   | 1270                           | 21x                  | 20.7                            |
| BRAHMS | 560                            | 13x                  | 12.2                            |
| PHOBOS | 540                            | 7x                   | 12.3                            |

# PHENIX in Run 4

- 200 GeV Au+Au data sample:
  - $1.5 \times 10^9$  min bias events recorded,
  - $241 \text{ mb}^{-1}$  integrated luminosity
  - 60 times the  $24 \times 10^6$  minbias events of Run 2
  - 10 times the  $24 \text{ mb}^{-1}$  sampled in Run 2 by triggered events
- 62.4 GeV Au+Au data sample:
  - $58 \times 10^6$  min bias events recorded




# $\pi^0$ production in p+p



- Good agreement with NLO pQCD
  - Factorization theorem:

$$\sigma_{AB \Rightarrow hX} \propto f_{a/A}(x_a, Q^2_a) \otimes f_{b/B}(x_b, Q^2_b) \otimes \sigma_{ab \Rightarrow cd} \otimes D_{h/c}(z_c, Q^2_c)$$





- Constrains Fragmentation Function  $D(\text{Gluon-}\pi)$
- Reference for Au+Au spectra

Phys. Rev. Let 91, 241803 (2003)

200 GeV - Run2




# $\pi^0$ production in p+p



- New run3 pp data
- Reach out to  $p_T > 16$  GeV
- Good reference for run3 dAu data – reduced syst. errors
- Cross section determination under investigation

200 GeV – Run3

# $\pi^0$ Spectra in Au+Au at 200 GeV



- Use p+p as base line measurement


spectra up to 15 GeV/c



central

Nuclear  
Physics


Particle  
Physics



peripheral


Phys. Rev. Lett. 91, 072301 (2003)

# Centrality dependence




integrated  $\pi^0$  yield  
above  $p_T = 4$  GeV/c

# Direct photons centrality



# $(h^+ + h^-)/2$ to $\pi^0$ ratio in 200GeV Au-Au

- Averaged charged hadrons to  $\pi^0$  ratios.
- Lines are drawn at  $h/\pi^0 = 1.6$ , which is predicted by results from past experiments
- At high  $p_T$ , ratio reaches the asymptotic values of 1.6
- In the intermediate  $p_T$  region, excess is seen
  - Consistent with the  $p/\pi^0$  ratio data showing that more protons are produced
  - Strong centrality dependence



# Initial ↔ Final State Effects


- Initial state effects
  - lead to  $R_{AA} \neq 1$  at high  $p_T$
  - but are not related to properties of hot and dense nuclear matter
- Possible initial state effects:
  - Initial state multiple soft scatterings (Cronin effect)  
 $R_{AA} > 1$
  - Modifications of nuclear structure functions in nuclei (Shadowing)  
 $R_{AA} < 1$
  - gluon saturation (Color Glass Condensate)  
 $R_{AA} < 1 ?$
- Final state effects
  - dense partonic medium parton energy loss (and recombination)
  - dense hadronic medium hadronic energy loss

# The Control Experiment: d+Au



- Initial state nuclear effects present in both A+A and N+A collisions
- Final state medium effects only present in A+A collisions

# $R_{AA}$ for $\pi^0$ in Central Collisions Different Energies



- Cronin Effect at lower energies
- Expectation
  - $R_{AA} > 1$
- Observed
  - factor 4-5 suppression at 130 and 200 GeV


A.L.S. Angelis PLB 185, 213 (1987)  
WA98, EPJ C 23, 225 (2002)  
PHENIX, PRL 88 022301 (2002)  
PHENIX submitted to PRL,  
nucl-ex/0304022

**Explanation ??**





# h/pi ratio


AuAu 62.4GeV



AuAu 62.4GeV



# $\pi^0$ Spectra at 62.4 GeV



- $\pi^0$  data in min. bias, 0-10%, and 10-30%, and 30-60% most central collisions up to  $p_T = 7 \text{ GeV}/c$
- Small systematic uncertainty : 9~12%

# Charged hadron reference

$\pi, k, p$  data from ISR are first combined to obtain charged hadron data at low  $\sqrt{s}$

- Charged hadron data are then interpolated between ISR, UA1 and PHENIX to obtain the reference data at  $\sqrt{s}$  62.4 GeV
- A fit using modified hagedon functional form is used to obtain parameterization for charged hadrons
- Right Fig. shows the charged reference/1.6 and compared with  $\pi^0$  and Breakstone which is not used in the fit. ( $(h^+ + h^-)/2\pi^0 = 1.6 \pm 0.16$  measured in ISR and RHIC)
- Charged hadron have  $\pm 25\%$  systematic errors and the upper error increase to about 50% at 7 GeV/c

