

# PDF Issues In MSSM Higgs Searches:

$$p\bar{p} \rightarrow b\bar{b}A \rightarrow b\bar{b}b\bar{b}$$

Jared Yamaoka

Rutgers University, CDF

Dec. 14, 2004



# Outline

- Introduction
- Cross Section
- Acceptance
  - Increased statistic (3x) since last time
- x Value
- Conclusion and Observations


## Why Search $b\bar{b}h$ Channel?

In the minimal supersymmetric extension to the Standard Model (MSSM), the  $b\bar{b}h$  Yukawa coupling is proportional to  $\tan \beta$ , thus the cross section grows as  $\tan^2 \beta$  with respect to SM.

Typical lowest order Feynman diagrams for the signal channel.



## DZero Run II vs. CDF Run I




DZero Run II Limit; March 2004

Using 130 pb<sup>-1</sup>

CDF Run I Limit; October 2000

Using 91 pb<sup>-1</sup>



How can DZero Run II limit be worse?!

## What is going on?

To see what might be causing the discrepancy between the Run I and the Run II analysis, we looked at the PDF's used in each analysis. CTEQ3L was used in the Run I analysis, but CTEQ5L is used in the Run II analysis.

- Differences in cross section due to PDF.
  - Using *PYTHIA* v6.216
  - Using PPHTT v1.1 from M. Spira
- Differences in acceptance.
  - Recreated CDF Run I event cuts.
- CDF Run I got lucky.
  - Run I analysis had less than expected background so it was able to set a better limit.

## CTEQ3L vs. CTEQ5L: *PYTHIA*

Signal Cross Section (pb)

Mass A	$\tan\beta$	CTEQ3L	CTEQ5L
		PYTHIA	PYTHIA
90	30	10.0	6.7
90	50	27.0	18.3
100	30	6.7	4.4
100	50	18.3	12.0

There seems to be about a factor of 1.5 difference in the cross sections across the board.

## CTEQ3L vs. CTEQ5L: *PPHTT* (As a Cross Check)

Signal Cross Section (pb)

Mass A	$\tan\beta$	CTEQ3L	CTEQ5L
		PPHTT v1.1	PPHTT v1.1
90	30	13.9	9.3
90	50	37.7	25.9
100	30	8.7	5.2
100	50	24.2	15.9

PPHTT v1.1 is a cross section calculator from M. Spira. It uses a leading order (LO) calculation where the scale used for the running b mass in the Yukawa coupling  $Q = (M_H + 2 * M_b)/2$ .

PPHTT shows the same trend as PYTHIA.

## Back to *PYTHIA*: CTEQ3L vs. CTEQ5L

Signal Cross Section (pb) *PYTHIA*

Mass A	$\tan\beta$	Process	CTEQ3L	CTEQ5L
90	30	gg	10	6.6
		qq	5.1e-2	5.2e-2
90	50	gg	27	18
		qq	0.14	0.14
100	30	gg	6.6	4.3
		qq	3.4e-2	3.4e-2
100	50	gg	18	12
		qq	9.2e-2	9.3e-2

CTEQ5L has a softer the gluon/gluon interaction than CTEQ3L.  
However the quark/quark interactions seem to be the same.



## CDF Run I Selection Cuts

We did our best to model the Run I selection cuts using current CDF Run II software.

- L2

- 4 Jets  $E_T > 15$  GeV
- $\Sigma E_T > 125$  GeV

- Kinematics

- $M_A$  dependent cuts on jet energy  
(This case  $M_A = 90$  GeV)
  - \* Hardest Jet  $> 42$  GeV
  - \*  $2^{nd}$  Hardest Jet  $> 34$  GeV
  - \*  $3^{rd}$  Hardest Jet  $> 14$  GeV

- b-Tagging

- At least 3 of the 4 hardest jets are b-tagged.

- bJetKin

- $\Delta\phi > 109^\circ$  between the 2 hardest b-tagged jets.

## Effect of the PDF on Acceptance: qq

*PYTHIA* Monte Carlo ( $M_A = 90$ ;  $\tan\beta = 50$ )

		CTEQ3L(qq)	CTEQ5L(qq)
$\sigma$ (pb)		0.14	0.14
Num MC		51k	59k
L2	Events	10935	12777
	Accept.(%)	21	22
	$\sigma \times Accept$	0.030	0.030
Kinematics	Events	2381	2774
	Accept.(%)	4.7	4.7
	$\sigma \times Accept$	0.007	0.007
b-Tagging	Events	330	356
	Accept.(%)	0.65	0.60
	$\sigma \times Accept$	0.0009	0.0008
bJetKin	Events	232	246
	Accept.(%)	0.46 $\pm$ .03	0.42 $\pm$ .03
	$\sigma \times Accept$	0.00063 $\pm$ .00004	0.00058 $\pm$ .00004

The ratio between the PDF's in the quark/quark process:

$$0.00063/0.00058 = 1.1$$

## Effect of the PDF on Acceptance: gg

*PYTHIA* Monte Carlo ( $M_A = 90$ ;  $\tan\beta = 50$ )

		CTEQ3L(gg)	CTEQ5L(gg)
$\sigma$		26.9	18.2
Num MC	Increased Stat.	352k	358k
L2	Events	2526	2376
	Accept.(%)	0.7	0.7
	$\sigma \times Accept$	0.19	0.13
Kinematics	Events	385	336
	Accept.(%)	0.11	0.09
	$\sigma \times Accept$	0.030	0.016
b-Tagging	Events	23	18
	Accept.(%)	0.007	0.005
	$\sigma \times Accept$	0.0018	0.0009
bJetKin	Events	19	11
	Accept.(%)	$0.0054 \pm .0012$	$0.0031 \pm .0010$
	$\sigma \times Accept$	$0.0015 \pm .0003$	$0.00056 \pm .00017$

The ratio between the PDF in the glue/glue process:

$$0.0015/0.00056 = 2.7$$

## Effect of the PDF on Acceptance: Total (qq + gg)

*PYTHIA* Monte Carlo ( $M_A = 90$ ;  $\tan\beta = 50$ )


		CTEQ3L(total)	CTEQ5L(total)
$\sigma$		27.04	18.31
L2	Accept.(%)	0.81	0.86
	$\sigma \times Accept$	0.22	0.16
Kinematics	Accept.(%)	0.13	0.13
	$\sigma \times Accept$	0.035	0.023
b-Tagging	Accept.(%)	0.010	0.010
	$\sigma \times Accept$	0.0027	0.0019
bJetKin	Accept.(%)	0.0078 $\pm$ .0012	0.0063 $\pm$ .0010
	$\sigma \times Accept$	0.0021 $\pm$ .0003	0.0011 $\pm$ .00017

The total ratio between the PDF's:

$$0.0021/0.0011 = 1.9)$$


## Some Kinematic Plots

$P_T$  of the Hardest b




Out in the high  $P_T$  tails, it seems the slopes of the glue/glue process diverge.

# x Values gg



## x Values qq



## Conclusion

- CTEQ3L to CTEQ5L, the cross section dropped by a factor of 1.5.
- The acceptance also dropped by a factor of 1.2.
- The total (qq+gg) ratio ( $\sigma * acceptance$ ) is factor of 1.9.
- Qualitatively this seems to account for some of the difference in the DZero Run II result and CDF Run I result.
- Now that this archeology is done, I'll move on, unhindered, toward my analysis result.

## General Observations

- PDF's make significant difference in this analysis.
- There needs to be good ways to estimate the errors associated with PDF's. CTEQ6 does this but with lots of effort.
- Experimenters need to have guidance on how to use the new NLO calculations.