

LHC program for very rare B decays

P. Rezníček^{a*}, For the ATLAS, CMS and LHCb collaboration

^aIPNP, Faculty of Mathematics and Physics, Charles University in Prague
V Holešovičkách 2, 18000 Prague-8, Czech Republic

This paper gives an overview of ATLAS, CMS and LHCb potential in measurement of di-muonic rare B-decays $B_s^0 \rightarrow \mu^+\mu^-$ and $B_d^0 \rightarrow \mu^+\mu^-$. Branching ratios (BR) of these decays are small due to helicity suppression and Flavour Changing Neutral Currents (FCNC) forbidden at tree level. Accounting clear theoretical picture for the BR prediction and simple experimental signature, measurement of the BR provides excellent probe of New Physics effects. While present experimental upper limits from Tevatron are roughly $50\times$ above Standard Model (SM) prediction, LHC experiments will reach the SM branching already around the end of initial low-luminosity stage of LHC (30 fb^{-1}). In the paper, offline analysis and the expected number of signal and background events are presented. Because of small BR, background coming from misidentification effects and rare exclusive and exotic decays can be important. Therefore part of this paper is also devoted to study of these background sources.

1. Introduction

In the Standard Model (SM), Flavour Changing Neutral Currents (FCNC) can occur through higher order diagrams only. Di-muonic FCNC B-decays $B_s^0 \rightarrow \mu^+\mu^-$ and $B_d^0 \rightarrow \mu^+\mu^-$ are also helicity suppressed, resulting in branching ratios $(3.42 \pm 0.52) \cdot 10^{-9}$ and $(1.00 \pm 0.14) \cdot 10^{-10}$ respectively [1]. The $B_d^0 \rightarrow \mu^+\mu^-$ rate is in SM further suppressed by the ratio of $|V_{td}/V_{ts}|^2$. These small rates provide great room for New Physics (NP) effects, that can enhance/suppress the BR significantly. Observation of both B_d^0 and B_s^0 decays is important in determining the flavour structure of NP, because in some models like e.g. R-parity violating SUSY [2] the relative suppression of B_d^0 with respect to B_s^0 may not retain.

Present best limits on the BRs, provided by Tevatron: CDF measurements at 780 pb^{-1} and DØ expectation at 700 pb^{-1} [3,4] - are shown in table 1. Expected improvement by the end of Tevatron run is by factor of $5\times-8\times$ [4], therefore remaining one order above SM values. Because NP can also suppress the branching, it is important to have sensitivity below SM rates. This is what LHC experiments are able to provide, especially after LHC enters the high luminosity stage.

*e-mail: Pavel.Reznicek@cern.ch

Table 1

Present BR limits at 95% CL and SM prediction

Branching ratio:	$B_s^0 \rightarrow \mu^+\mu^-$	$B_d^0 \rightarrow \mu^+\mu^-$
CDF (780 pb^{-1})	$1.0 \cdot 10^{-7}$	$3.0 \cdot 10^{-8}$
DØ (700 pb^{-1})	$2.3 \cdot 10^{-7}$	–
SM prediction	$3.4 \cdot 10^{-9}$	$1.0 \cdot 10^{-10}$

2. Trigger Strategies

Trigger strategies for di-muonic very rare decays at ATLAS, CMS and LHCb depends on each experiment. Details can be found in [5].

LHCb trigger is divided into three steps. The first level (L0) requires two high transverse momenta ($p_T \sim 1 - 5 \text{ GeV}$) tracks found in muon stations. At the next stage (L1) topological cuts are performed on reconstructed B-vertex. Finally, high-level trigger (HLT) consists of full event reconstruction and application of specific cuts.

At ATLAS and CMS, the situation is different from LHCb due to detector acceptance (central geometry compared to forward) and higher instantaneous luminosity ($10^{33}/10^{34} \text{ cm}^{-2}\text{s}^{-1}$ vs. $2 \cdot 10^{32} \text{ cm}^{-2}\text{s}^{-1}$). The p_T cuts on muons are higher at the first stage (LVL1): $\sim 6 \text{ GeV}$ at ATLAS and $\sim 3 \text{ GeV}$ at CMS, but vertex trigger is not feasible here. Vertexing and specific cuts are applied at the last level: Event Filter (EF).

3. Offline Analysis

The offline analysis is based on similar cuts as in CDF and DØ measurements. When distinguishing combinatorial background from the signal, the simplicity of the experimental signature limits variety of cuts, that can be used, to these:

- B-hadron invariant mass window
- Secondary vertex displacement and quality
- Pointing of B-hadron momentum to primary vertex (PV)
- Isolation in tracker and/or in calorimeters

The invariant mass window is driven by detector resolution. But it should be accounted that the resolution may not be sufficient to distinguish B_d^0 from B_s^0 . In that case joint analysis is needed. Secondary vertex displacement and quality cuts reduce combinatorial background from primary-vertex tracks. Isolation requirement rejects secondary vertices with more than two tracks and pointing to primary vertex constraint prevents from existence of detector invisible particles originating in the same vertex.

3.1. ATLAS

The ATLAS offline analysis combines high- p_T muon tracks to di-muon pairs with invariant mass $M_{\mu^-\mu^+} = M_{B_s^0}^{+140\text{MeV}}_{-70\text{MeV}}$. The asymmetry is to separate B_s^0 from B_d^0 . $B_s^0 \rightarrow \mu^+\mu^-$ mass resolution at ATLAS is 80 MeV. In the present study, isolation in ATLAS tracker is used, requiring no charged tracks with $p_T > 0.8$ GeV in cone $\theta < 15^\circ$ around B-meson momentum. Vertex fit with pointing constraint is performed and vertices with transverse decay length significance $L_{xy}/\sigma(L_{xy}) < 11$ or $\chi^2 > 15$ are cut out. The expected number of SM signal and background events, corresponding upper limits on BR and single event sensitivity (SES) are shown in table 2. The upper limits are calculated according to [7]. SES is defined as $1/(\text{total no. of BG events}) \times \sigma_{\text{BG}}/(\sigma_{B_s^0} \cdot \alpha) \cdot \epsilon_\mu^2$. Where $\sigma_{B_s^0}$ is cross-section and α acceptance of $B_s^0 \rightarrow \mu^+\mu^-$ decay with $p_T(\mu) > 6$ GeV and $|\eta(\mu)| < 2.5$ ($\sigma_{B_s^0} \cdot \alpha = 0.42$ μb), $\sigma_{\text{BG}} = 600$ pb is $b\bar{b} \rightarrow \mu^+\mu^-X$ cross-section within the same $p_T(\mu)$ and $\eta(\mu)$ cuts, and ϵ_μ stands for muon identification efficiency.

Table 2

ATLAS offline analysis - $B_s^0 \rightarrow \mu^+\mu^-$

L_{int}	SES	S/BG	90% CL limit
100 pb ⁻¹	$2.7 \cdot 10^{-8}$	$\sim 0/0.2$	$6.4 \cdot 10^{-8}$
10 fb ⁻¹	$2.7 \cdot 10^{-10}$	$\sim 7/20$	$7.0 \cdot 10^{-9}$
30 fb ⁻¹	$0.9 \cdot 10^{-10}$	$\sim 21/60$	$6.6 \cdot 10^{-9}$

3.2. CMS

CMS mass resolution is 46 MeV. The analysis uses $|M_{\mu^+\mu^-} - M_{B_s^0}| < 40$ MeV invariant mass cut. Both tracker and calorimetry isolations are applied, requiring no track in $\Delta R < (0.5 \cdot \Delta R_{\mu^+\mu^-} + 0.4)$ and $E_T < 4/6\text{GeV}$ (high/low luminosity) in the same ΔR in electromagnetic and hadronic calorimeters. The muon tracks have to have minimal distance in-between $d < 50$ μm and its significance $d/\sigma(d) < 2$. B_s^0 transverse decay length cuts $L_{xy}/\sigma(L_{xy}) > 12$ (15 at L_{high}) and its uncertainty $\sigma(L_{xy}) < 80$ μm . Pointing constraint is represented by cosine of angle between momentum and vertex direction to be above 0.9. The preliminary number of signal and background events shown in table 3 correspond to 4σ SM observation at 30 fb⁻¹ and 6.3σ at 130 fb⁻¹ (1 year at L_{high}). But new study is in progress to study full selections, HLT and offline cuts.

Table 3

CMS offline analysis - $B_s^0 \rightarrow \mu^+\mu^-$

L_{int}	Signal	Background
30 fb ⁻¹	7.0	< 1.0
100 fb ⁻¹	26.0	< 6.4

3.3. LHCb

LHCb constructs di-muons from $p_T > 1.3$ GeV muon candidates. With 18 GeV mass resolution, the analysis selects ± 60 MeV mass window around $M_{B_s^0}$ [8]. Cut on muon tracks impact parameter is applied: $IP_\mu/\sigma_\mu > 3$. In the vertex, B_s^0 impact parameter is required to be $IP_{B_s^0}/\sigma_{B_s^0} < 3$, vertex $\chi^2 < 3^2$ and pointing constraints angle momentum-vertex below 5 mrad. This study results in expectation of ≥ 30 events/year. Background rejection was tested on $30 \cdot 10^6$ $b\bar{b} \rightarrow \text{inclusive}$ events, where none with $M_{\mu^+\mu^-} > 4$ GeV passed the cuts.

4. Backgrounds

The main source of background come from random combinatorics. The two muon candidates can originate either from semileptonic decays of b and \bar{b} quarks or from cascade decays of one of the $b\bar{b}$ quarks. Due to extremely low BR of the signal, relatively rare effects can become important. But for Monte Carlo (MC) studies it has to be accounted that common generators do not include rare decay channels ($BR \leq \sim 10^{-5}$). Misidentification can also not-negligibly contribute to the background. Some of the effects can be suppress or excluded by convenient mass resolution.

The number of MC background production is limited by slowness of full detector simulation chain ($100\times$ events/day/CPU). Therefore correlation between cuts have to be studied and rejection of events already at generator level applied (e.g. based on invariant mass of high- p_T muons, selection of special channels etc.). Fast approximate simulations (like e.g. Atlfast in ATLAS) are not advisable, since the background events come from rather tail resolutions effects.

4.1. Two-body hadronic decays

An example of rare-cases are two-body hadronic decays of B-mesons, when one or both mesons have short lifetime, so that they decay inside tracker (ID):

- $B_d^0 \rightarrow D_{\rightarrow\mu^+X_s}^+ D_{\rightarrow\mu^-X_s}^-$ in ID: $BR \sim 10^{-6}$
- $B_d^0 \rightarrow D_{\rightarrow\mu^+X_d}^+ D_{\rightarrow\mu^-X_d}^-$ in ID: $BR \sim 10^{-8}$

and similar decays $B_d^0 \rightarrow K^+ D^-$, $B_s^0 \rightarrow K^+ D_s^-$ or $B_s^0 \rightarrow D_s^{*+} D_s^{*-}$. None of these channels is included in common MC generators and therefore separated study have to be performed.

4.2. Very rare decays $B^{0\pm} \rightarrow (\pi^{0\pm}, \gamma)\mu^+\mu^-$

Branching ratio of these decays is $\sim 2 \cdot 10^{-8}$ [6]. The background come from soft π/γ escaping detection. In case of $B^{0\pm} \rightarrow \pi^{0\pm}\mu^+\mu^-$ decay, the resulting invariant mass is $M_{\mu^+\mu^-} \sim M_{B^{0\pm}} - M_{\pi^{0\pm}}$, which remains in the detector resolution of ATLAS only. Background from B^\pm is expected to be less significant than from B^0 , due to the lower p_T limit excluding the pion detection: ATLAS example:

- for π^\pm not to be detected by inner tracker, $p_T \leq 0.5$ GeV
- for π^\pm not to be detected by electromagnetic calorimeter, $p_T \leq \sim$ few GeV

Initial study based on particle-level simulation was performed. Di-muon invariant mass distribution from these two background channels compared to signal peaks indicates, that the $B^0 \rightarrow \gamma\mu^+\mu^-$ do not significantly contribute, while $B^{0\pm} \rightarrow \pi^{0\pm}\mu^+\mu^-$ needs more checking.

At this point, it should be also mentioned, that these two very rare decay channels may be worth to study as signal too, since some of their properties (like e.g. di-muon invariant mass spectrum) are also sensitive to NP contributions [6].

4.3. Four-leptonic B-decays

Another decay channels that were found to be able to produce $B_{s,d}^0 \rightarrow \mu^+\mu^-$ background are purely leptonic decays $B_{(c)}^+ \rightarrow \mu^+\mu^-l^+\nu_l$. If the p_T of one of the leptons is out of reconstruction capabilities (e.g. $p_T \leq 0.5$ GeV at ATLAS), then there are only two tracks observed from the B-meson vertex and the invariant mass of the di-lepton pair can be close to $B_{s,d}^0$ mass. Branching ratio of these decays are $5 \cdot 10^{-6}$ and $8 \cdot 10^{-5}$ for B^+ and B_c^+ respectively. In spite of higher BR, the contribution from B_c^+ will be less significant due to $400\times$ lower production cross-section and $4\times$ shorter lifetime, therefore more efficiently rejected by secondary vertex cut. The di-muonic spectrum from these four-leptonic decays is shown in figure 1, showing just initiating particle-level simulation biased by not-requiring B-meson momentum pointing to primary vertex.

4.4. Misidentification effects

Considering typical hadron misidentification probability of being an muon $\sim 0.5\%$, such effects are obviously not negligible with respect to the $\sim 10^{-9}$ branching. The most simpler background comes from two body hadronic decays: $B_{d,s}^0 \rightarrow K^\pm\pi^\mp$, $B_{d,s}^0 \rightarrow K^\pm K^\mp$, $B_{d,s}^0 \rightarrow \pi^\pm\pi^\mp$ etc. The fake probability can be estimated by: $BR(B_d^0 \rightarrow K^\pm\pi^\mp) \times (0.5\%)^2 = 2 \cdot 10^{-5} \times (0.005)^2 = 0.5 \cdot 10^{-9}$, which is of the same order as the BR of signal channels. The back-

ground contribution can be calculated by convoluting fake probability with K and π spectrum. Such an study at LHCb resulted in having ~ 2 events per 2 fb^{-1} (in $\pm 2 \cdot \sigma$ mass window). Decays of π and K to muons were found not producing significant background (at ATLAS).

Fake signal events can also originate from two-body hadronic decays with soft muon in final state, e.g.: $B^+ \rightarrow (J/\psi \rightarrow \mu^+\mu^-)K^+$ ($BR \sim 6 \cdot 10^{-5}$). Considering $K - \mu$ misidentification and soft μ^+ not reconstructed by tracker (e.g. $p_T \leq 0.5 \text{ GeV}$ at ATLAS with probability ~ 0.1), the fake rate can be calculated as: $6 \cdot 10^{-5} \times 0.5\% \times 0.1 \sim 10^{-8}$. Other contributing channels are $B_{(c)}^+ \rightarrow (J/\psi \rightarrow \mu^+\mu^-)\pi^+$ and $B^+ \rightarrow (\psi(2S) \rightarrow \mu^+\mu^-)K^+$.




Figure 1. Di-muonic invariant mass spectrum from four-leptonic B-decays and misidentification in $B_d^0 \rightarrow \pi^- \mu^+ \nu_\mu$. Particle level simulation only without pointing to PV constraint.

Finally the fake events can also be generated by semileptonic B-decays like $B_d^0 \rightarrow \pi^- \mu^+ \nu_\mu$ with $BR \sim 10^{-4}$. As in previous case, accounting $\pi - \mu$ misidentification and soft-neutrino phase space (probability ~ 0.1), the fake rare would be $0.5 \cdot 10^{-7}$. Similar channels to be accounted are $B_s^0 \rightarrow K^- \mu^+ \nu_\mu$ and $B^+ \rightarrow K^+ \mu^+ \mu^-$. The di-muonic spectrum coming from semileptonic B-decay $B_d^0 \rightarrow \pi^- \mu^+ \nu_\mu$ is shown in figure 1.

5. Summary

$B_{s,d}^0 \rightarrow \mu^+ \mu^-$ branching ratios measurement provides powerful tool for indirect New Physics

search. Due to $BR \sim 10^{-9}$, present experiments will detect signal only when the BR is strongly enhanced by New Physics, while LHC sensitivity below SM expectation allow for discovering both enhancement and suppression of BR.

When estimating the background at LHC, rare and exotic decays have to be taken into account (some of them would be interesting to measure as signal too). Present experiments are not so sensitive to be challenged by this kind of background, however LHC and any future projects will have to consider such effects carefully. The other non-negligible contribution can come from hadron-muon misidentification.

The offline analyses foreseen, that Tevatron limits will be overcome by 1st year measurement at low luminosity and by the end of 3rd year (end of low-luminosity LHC stage), SM branching should be reached. ATLAS, CMS and LHCb are also putting large effort to ensure continuation of very rare decay program at high luminosity stage.

REFERENCES

1. G. Buchalla and A. J. Buras, Nucl. Phys. B400, 225 (1993); A.J. Buras, Phys. Lett. B 566, 115 (2003).
2. R. Arnowitt et al., Phys. Lett. B 538, 121 (2002).
3. J. Rieger, Rare decays of B-hadron at Tevatron, BEACH2006 www.hep.lancs.ac.uk/Beach2006
4. V. Krutelyov, Rare B-decays at Tevatron, HCP2006, hcp2006.phy.duke.edu
5. The LHCb collaboration, LHCb Trigger System, CERN/LHCC 2003-031 (2003); T. Speer, ATLAS and CMS detectors and triggers for B-physics, BEACH2006 www.hep.lancs.ac.uk/Beach2006
6. D.Melikhov, N.Nikitin, Phys. Rev. D70, 114028, (2004)
7. J. Heinrich, Bayesian limit software: multi-channel with correlated backgrounds and efficiencies, CDF Internal Note 7587 (2005)
8. B. de Paula, F. Marinho, S. Amato, Analysis of the Rare $B_s^0 \rightarrow \mu^+ \mu^-$ Decay with the Reoptimized LHCb Detector, LHCb Internal Note LHCb-2003-16 (2003)