



Contribution ID: **1006**

Type: **Oral Presentation**

## **Searching for Dark Matter Using the NOvA Upward-going Muon Trigger (15' + 5')**

*Thursday 4 August 2016 17:00 (20 minutes)*

The NOvA collaboration has constructed a 14,000 ton, fine-grained, low-Z, total absorption tracking calorimeter at an off-axis angle to an upgraded NuMI neutrino beam. This detector, with its excellent granularity and energy resolution and relatively low-energy neutrino thresholds, was designed to observe electron neutrino appearance in a muon neutrino beam, but it also has unique capabilities suitable for more exotic efforts. In fact, if sufficient cosmic ray background rejection can be demonstrated, NOvA will be capable of a competitive indirect dark matter search for low-mass Weakly-Interacting Massive Particles (WIMPs). The cosmic ray muon rate at the NOvA far detector is approximately 100 kHz and provides the primary challenge for triggering and optimizing such a search analysis. We present the first dark matter search results using data collected with the upward-going muon trigger.

**Authors:** TSARIS, Aristeidis (Fermilab); OKSUZIAN, Iuri (UVa); GROUP, Robert (University of Virginia); MINA, Robert (University of Virginia)

**Presenter:** TSARIS, Aristeidis (Fermilab)

**Session Classification:** Dark Matter Detection

**Track Classification:** Dark Matter Detection