


Searches for R-Parity violating SUSY with the ATLAS detector



ICHEP2016CHICAGO

Aug. 3-10, 2016



Suen Hou
Academia Sinica

for the ATLAS collaboration



R-Parity violating search

Search for R-parity violating SUSY

$$R = (-1)^{2s+3(B-L)}$$

Super-potential with RPV of lepton or baryon number

$$\mathcal{W}_{\Delta L=1} = \lambda_{ijk} L_i L_j \bar{E}_k + \lambda'_{ijk} L_i Q_j \bar{D}_k + \kappa_i L_i H_d$$


$$\mathcal{W}_{\Delta B=1} = \lambda''_{ijk} \bar{U}_i \bar{D}_j \bar{D}_k$$

Analyses updated to **13 TeV** data

$LL\bar{E}$ 4 leptons

ATLAS-CONF-2016-075


13.3 fb⁻¹



$\bar{U}\bar{D}\bar{D}$ stop 4-jet

ATLAS-CONF-2016-084


15.4 fb⁻¹



$\bar{U}\bar{D}\bar{D}$ multi-jet

ATLAS-CONF-2016-057


14.8 fb⁻¹



$\bar{U}\bar{D}\bar{D}$ SS/3Leptons

ATLAS-CONF-2016-037

13.2 fb⁻¹



ATLAS RPV searches


SUSY RPV publications (July 2016)

Model	observables	mass limit	$\sqrt{s} = 7, 8 \text{ TeV}$	$\sqrt{s} = 13 \text{ TeV}$	reference
LFV $pp \rightarrow \tilde{\nu}_\tau + X, \tilde{\nu}_\tau \rightarrow e\mu/\epsilon\tau/\mu\tau$	$\tilde{\nu}_\tau$ $e\mu, e\tau, \mu\tau$ -			1.7 TeV	1503.04430
Bilinear RPV CMSSM	\tilde{q}, \tilde{g} $2 e, \mu$ (SS) 0-3 b			1.45 TeV	1404.2500
$\tilde{\chi}_1^+ \tilde{\chi}_1^-, \tilde{\chi}_1^+ \rightarrow W \tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow ee\tilde{\nu}_\mu, e\mu\tilde{\nu}_e$	$\tilde{\chi}_1^\pm$ $4 e, \mu$ -		760 GeV		1405.5086
$\tilde{\chi}_1^+ \tilde{\chi}_1^-, \tilde{\chi}_1^+ \rightarrow W \tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \tau\tau\tilde{\nu}_e, e\tau\tilde{\nu}_\tau$	$\tilde{\chi}_1^\pm$ $3 e, \mu + \tau$ -		450 GeV		1405.5086
$\tilde{g}\tilde{g}, \tilde{g} \rightarrow qq\tilde{q}$	\tilde{g} 0 6-7 jets		917 GeV		1502.05686
$\tilde{g}\tilde{g}, \tilde{g} \rightarrow qq\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow qq\tilde{q}$	\tilde{g} 0 6-7 jets		980 GeV		1502.05686
$\tilde{g}\tilde{g}, \tilde{g} \rightarrow t_1 t_1, t_1 \rightarrow bs$	\tilde{g} $2 e, \mu$ (SS) 0-3 b		880 GeV		1404.2500
$\tilde{t}_1 \tilde{t}_1, \tilde{t}_1 \rightarrow bs$	\tilde{t}_1 0 2 jets + 2 b	345 GeV			ATLAS-CONF-2016-022
$\tilde{t}_1 \tilde{t}_1, \tilde{t}_1 \rightarrow b\bar{b}$	\tilde{t}_1 $2 e, \mu$ $2 b$			0.4-1.0 TeV	ATLAS-CONF-2015-015

8 TeV analyses


$LL\bar{E}+LQ\bar{D}$ di-leptons

1503.04430




$LQ\bar{D}$ $\tilde{t} \rightarrow \ell b$

ATLAS-CONF-2015-015




$LQ\bar{D}$ 1-lepton

ATLAS-CONF-2015-018



bilinear

ATLAS-CONF-2015-018




λ_{12k} : 4-lepton RPV search

Chargino pair production to neutralino (LSP)

Lepton number RPV by $\lambda_{121}, \lambda_{122}$ of

$$\tilde{\chi}_1^0 \rightarrow e^+e^-\nu, \tilde{\chi}_1^0 \rightarrow e^\pm\mu^\mp\nu, \tilde{\chi}_1^0 \rightarrow \mu^+\mu^-\nu$$

- Data of 13 TeV, 13.3 fb⁻¹
 - ≥ 4 leptons (e, μ),
 - Z veto, $m_{e^+e^-} \notin [81.2, 101.2]$
 - effective mass: $m_{\text{eff}} = p_T(\ell) + p_T(\text{jet}) + E_t^{\text{miss}}$
- Signal region: $\geq 4\ell, m_{\text{eff}} > 600, 800 \text{ GeV}$
- Validation conducted using events in $m_{\text{eff}} < 600 \text{ GeV}$ dominated by ZZ, t \bar{t} , Z+jets



see also, talk by D. Zhang:
“EW production, Gauginos and sleptons”

λ_{12k} : 4 leptons RPV exclusion

Chargino pair production assuming RPV



with $\lambda_{ijk} L_i L_j \bar{E}_k$ coupling of

$$\tilde{\chi}_1^0 \rightarrow \ell_k^\pm \ell_{i/j}^\mp \nu_{j/i}$$


Exclusion of $\tilde{\chi}_1^\pm$

with mass up to 1.1 TeV

for $m(\tilde{\chi}_1^0) > 400$ GeV



Resonance pair: 4-jet search



4-jet final states in search of

- top squark pair, RPV λ''_{3jk} coupling to quarks
- coloron pair

Event selection: 13 TeV data, 15.4 fb^{-1}

- anti- k_t jet of $R=0.4$, $p_T > 120 \text{ GeV}$
- 4 jets, paired by
$$\Delta R_{min} = \sum_{i=1,2} |\Delta R_i - 1.0|$$
- background suppression
$$\Delta R_{min} < 0.03 m_{avg}/\text{GeV}$$

 for two resonances of equal masses
$$m_{avg} < (m_1 + m_2)/2$$
- b-tag not required



Resonance pair, 4-jet final state

Pair production of heavy resonances
each decay into a jet-pair

- jet-pair $|\cos\vartheta^*|$ are more central
- masses of the two jet pairs about equal

$$A = \frac{|m_1 - m_2|}{m_1 + m_2}$$

Distributions compared to stop $m(\tilde{t}) = 400$ GeV, coloron $m_C = 1250$ GeV



4-jets discriminant

- Signal region (**SR**):

$$\mathcal{A} < 0.05, \quad |\cos\vartheta^*| < 0.5$$

- Discriminant variable:

$$m_{\text{avg}} = (m_1 + m_2) / 2$$



- Background in **SR**:

SM multi-jets, by distributions in quadrants of \mathcal{A} , $|\cos\vartheta^*|$ (*ABCD method*)

- Validation :

m_{avg} distributions in

$$0.05 < \mathcal{A} < 0.15, \quad |\cos\vartheta^*| < 0.5$$






Resonance pair: 4-jet exclusion

- m_{avg} in signal region:
events counted for stop, coloron
in mass windows of hypotheses

- Exclusion 95% CL
(13 TeV, 15.4 fb⁻¹)

stop λ''_{3jk} RPV $250 < m(\tilde{t}) < 405$ GeV
 $445 < m(\tilde{t}) < 510$ GeV

coloron $250 < m_C < 1500$ GeV






λ''_{ijk} : multi-jet RPV gluino search

Gluino pair production RPV, λ''_{ijk} coupling to jets

Event selection: 13 TeV, 14.8 fb⁻¹

- multiple jets and b-tag
- anti-k_t of **large $R=1.0$** , leading jet of $p_T > 440$ GeV
- topological variable: $M_J^\Sigma = \sum^4 m^{\text{jet}}$
jet mass of 4 leading jets
- discriminant on pseudorapidity: $|\Delta\eta_{12}|$
difference of the 2 leading jets



λ''_{ijk} : multi-jet validation

Glauino Signal (**SR**), Validation (**VR**) and Control (**CR**) regions

divided on $N_{\text{jet}}, |\Delta\eta_{12}|, \text{b-tag/inclusive}$

b-tag of small $R=0.4$ jets

N_{jet}	$ \Delta\eta_{12} > 1.4$	$ \Delta\eta_{12} < 1.4$
= 3	3j CR	—
≥ 4	4j VR	4j SR
≥ 5	5j VR	5j SR



Data-driven background

single-jet mass template from **CR** binned in p_T, η

jets of sample \rightarrow mass randomly picked from the template, for each jet p_T, η

M_J^Σ background calculated

Topological M_J^Σ : examined for samples in **VR**, **SR**



λ''_{ijk} : multi-jet exclusion


Observables in SR's
w/w.o. b-tag, M_J^Σ cuts

Region	M_J^Σ cut	observed	predicted
4jSRb1	> 0.8 TeV	24	$36.5 \pm 7.7 \pm 3.9 \pm 7.5$
4jSR	> 0.8 TeV	66	$88.2 \pm 10.7 \pm 8.6 \pm 11.8$
5jSRb1	> 0.6 TeV	15	$9.0 \pm 2.6 \pm 0.9 \pm 1.5$
5jSR	> 0.6 TeV	35	$27.5 \pm 5.5 \pm 3.3 \pm 3.5$




RPV gluino pair excluded

for

$m(\tilde{g})$ up to 1080 GeV



$m(\tilde{g})$ to 1050 ~1550 GeV
vs $m(\chi_1^0)$



Sample of 5jSRb1

λ''_{ijk} : Same-Sign di-lepton, 3 leptons


Same-Sign dilepton (e, μ) or 3 leptons + jets

- Analysis conducted for **RPC** and **RPV** scenarios
- Results of **RPV** λ''_{ijk} coupling are presented here

Event selection: 13 TeV, 13.2 fb⁻¹

Final states are clean to SM

- lepton (e or μ), $p_T > 10$ GeV
- SS or 3L
- jet of anti- k_t $R=0.4$, $p_T > 50$ GeV
- b-jets of $p_T > 20$ GeV



λ''_{ijk} : SS, 3-Lepton signals

Discriminant: effective mass

$$m_{\text{eff}} = \sum p_T^{\text{lep}} + \sum p_T^{\text{jet}} + E_T^{\text{miss}}$$

Background:

- SM processes: $t\bar{t}V$, diboson, .. by MC
- Detector charge mis-ID :
 electrons, estimated using $Z \rightarrow ee$ events
 fake, non-prompt leptons :
 estimated with data, $Z \rightarrow \ell\ell$ in p_T, η



Validation:

- Regions close to signal selection
- Regions with data dominated by $t\bar{t}V$ or diboson compared to MC

Signal region:

Signal region	$N_{\text{lept}}^{\text{signal}}$	$N_{b\text{-jets}}^{20}$	N_{jets}	$p_{T,\text{jets}}$	m_{eff} [GeV]
SR1b-DD	≥ 2	≥ 1	≥ 4	50	> 1200
SR3b-DD	≥ 2	≥ 3	≥ 4	50	> 1000
SR1b-GG	≥ 2	≥ 1	≥ 6	50	> 1800

} $d\bar{d}$
 $\tilde{g}\tilde{g}$




λ''_{ijk} : SS/3L distributions


Number of events selected

	SR1b-GG	SR1b-DD	SR3b-DD
Observed	1	10	2
Total SM	1.24 ± 0.34	8.4 ± 2.0	1.14 ± 0.35
$t\bar{t}Z$	0.18 ± 0.06	2.0 ± 0.7	0.21 ± 0.07
$t\bar{t}W$	0.25 ± 0.09	1.3 ± 0.4	0.13 ± 0.05
Diboson	0.05 ± 0.06	0.5 ± 0.4	< 0.1
Rare	0.46 ± 0.25	1.9 ± 1.0	0.53 ± 0.28
Fake/non-prompt le $_\tau$	0.15 ± 0.15	1.5 ± 1.3	0.15 ± 0.15
Charge-flip	0.15 ± 0.07	1.21 ± 0.18	0.11 ± 0.04


SR1b-GG



SR1b-DD



SR3b-DD






λ''_{ijk} : SS/3L RPV exclusion




RPV $\lambda''_{321} \lambda''_{313}$
exclusion, 95% CL

$m(\tilde{g}) \approx 1.3 \text{ TeV}$

for $m(\tilde{t})=600 \text{ GeV}$




$m(\tilde{d}_R)$ to 700 GeV



Summary

- Predictions of R-Parity Violating SUSY are searched
- Analyses updated with 13 TeV data reveal no evidence
- Higher mass limits excluded for SUSY particles

Model	mass limit	reference
LFV $pp \rightarrow \tilde{\nu}_\tau + X, \tilde{\nu}_\tau \rightarrow e\mu / e\tau / \mu\tau$	$\tilde{\nu}_\tau$ 1.9 TeV	1607.08079
Bilinear RPV CMSSM	\tilde{q}, \tilde{g} 1.45 TeV	1404.2500
$\tilde{\chi}_1^+ \tilde{\chi}_1^-, \tilde{\chi}_1^+ \rightarrow W \tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow e e \nu, e \mu \nu, \mu \mu \nu$	$\tilde{\chi}_1^\pm$ 1.14 TeV	ATLAS-CONF-2016-075
$\tilde{\chi}_1^+ \tilde{\chi}_1^-, \tilde{\chi}_1^+ \rightarrow W \tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \tau \tau \nu_e, e \tau \nu_\tau$	$\tilde{\chi}_1^\pm$ 450 GeV	1405.5086
$\tilde{g}\tilde{g}, \tilde{g} \rightarrow qqq$	\tilde{g} 1.08 TeV	ATLAS-CONF-2016-057
$\tilde{g}\tilde{g}, \tilde{g} \rightarrow qq\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow qqq$	\tilde{g} 1.55 TeV	ATLAS-CONF-2016-057
$\tilde{g}\tilde{g}, \tilde{g} \rightarrow \tilde{t}_1 t, \tilde{t}_1 \rightarrow bs$	\tilde{g} 1.3 TeV	ATLAS-CONF-2016-037
$\tilde{t}_1 \tilde{t}_1, \tilde{t}_1 \rightarrow bs$	\tilde{t}_1 410 GeV 450-510 GeV	NF-2016-022, ATLAS-CONF-2016-084
$\tilde{t}_1 \tilde{t}_1, \tilde{t}_1 \rightarrow b\ell$	\tilde{t}_1 0.4-1.0 TeV	ATLAS-CONF-2015-015



Backup