ATLAS measurements of the ridge in *pp* and *p*+Pb collisions

ATLAS measurements of correlations between particle pairs in relative azimuthal angle ($\Delta\phi$) and pseudorapidity separation ($\Delta\eta$) in pp collisions at \sqrt{s} =2.76, 5.02 and 13 TeV, and in p+Pb collisions at $\sqrt{s_{NN}}$ =5.02 TeV are presented. Prior measurements have shown that in pp collisions with a large multiplicity of produced particles, a long-range structure (the "ridge") develops along $\Delta\eta$ at $\Delta\phi \sim 0$. However, due to the presence of the large away-side jet, the full $\Delta\phi$ dependence of the long-range correlation could not be studied. In this analysis, a template fitting procedure is implemented to determine the contributions from dijets to the correlations, using low-multiplicity

events, and to extract the genuine long-range correlation. The long-range correlations are shown to be present even in events with a small multiplicity of produced particles, implying that the long-range correlations are not unique to rare high multiplicity events. The properties of the correlation in pp and p+Pb collisions shown to

be remarkably similar.

Preferred Track

QCD in small systems

Collaboration

ATLAS

Primary author: TU, Xiao (Columbia University (US))
Co-author: COLLABORATION, ATLAS (CERN)
Presenter: TU, Xiao (Columbia University (US))
Session Classification: Poster Session