

Supersymmetric explanation of CP violation in $K \rightarrow \pi\pi$ decays


Teppei Kitahara

Karlsruhe Institute of Technology (KIT), TTP
with **Ulrich Nierste** and **Paul Tremper**

Phys. Rev. Lett. 117, 091802 (2016)
[arXiv:1604.07400], arXiv:1607.06727

KAON2016

September 15, 2016,
Birmingham University



Kaon & CP violation:1

- Precise measurement for Kaon decay discovered the two type of CP violations: Indirect (mixing) (ϵ_K) & Direct CP violation (ϵ'_K)

[Christenson, Cronin, Fitch, Turlay, 64' with Nobel prize]


$$\mathcal{A}(K_L \rightarrow \pi^+ \pi^-) \propto \epsilon_K + \epsilon'_K$$

$$\text{with } \epsilon_K = \mathcal{O}(10^{-3}) \neq 0$$

$$\mathcal{A}(K_L \rightarrow \pi^0 \pi^0) \propto \epsilon_K - 2\epsilon'_K$$


$$\epsilon'_K = \mathcal{O}(10^{-6}) \neq 0$$

[NA48/CERN and KTeV/FNAL 99']



$$\epsilon_K \propto \text{Im}[(\text{CKM})^2]$$

$$K^0 \longleftrightarrow \bar{K}^0$$



$$\epsilon'_K \propto \text{Im}[\text{CKM}]$$

Kaon & CP violation:2

$$\epsilon_K \equiv \frac{2\eta_{+-} + \eta_{00}}{3} \in \mathbb{C}$$

$$\epsilon'_K \equiv \frac{\eta_{+-} - \eta_{00}}{3} \in \mathbb{C}$$

$$\eta_{00} \equiv \frac{\mathcal{A}(K_L \rightarrow \pi^0 \pi^0)}{\mathcal{A}(K_S \rightarrow \pi^0 \pi^0)}$$

$$\eta_{+-} \equiv \frac{\mathcal{A}(K_L \rightarrow \pi^+ \pi^-)}{\mathcal{A}(K_S \rightarrow \pi^+ \pi^-)}$$

$$\begin{aligned} \frac{\epsilon'_K}{\epsilon_K} &\simeq \frac{1}{\sqrt{2}|\epsilon_K|} \frac{\text{Re}A_2}{\text{Re}A_0} \left(\frac{\text{Im}A_2}{\text{Re}A_2} - \frac{\text{Im}A_0}{\text{Re}A_0} \right) \\ &= \frac{1}{\sqrt{2}|\epsilon_K|} \frac{\text{Re}A_2}{(\text{Re}A_0)^2} \left(-\text{Im}A_0 + \frac{\text{Re}A_0}{\text{Re}A_2} \text{Im}A_2 \right) \end{aligned}$$

with $\mathcal{A}(K^0 \rightarrow (\pi\pi)_I) \equiv \mathcal{A}_I e^{i\delta_I}$
 isospin amplitude $\mathcal{A}(\bar{K}^0 \rightarrow (\pi\pi)_I) \equiv \bar{\mathcal{A}}_I e^{i\delta_I} = \mathcal{A}_I^* e^{i\delta_I}$

General remarks

- This formula is modified by $m_u \neq m_d$ [Cirigliano,Pich,Ecker,Neufeld,PRL 03']
- Theoretical value of ϵ'_K/ϵ_K is real number
- $|\epsilon_K|$, $\text{Re}A_0$, and $\text{Re}A_2$ have been measured by experiments very precisely
- Theorist calculates $\text{Im}A_0$, and $\text{Im}A_2$ for ϵ'_K/ϵ_K
- Experiments can precisely probe ϵ'_K/ϵ_K by the following combination

$$\text{Re} \left[\frac{\epsilon'_K}{\epsilon_K} \right] \simeq \frac{1}{6} \frac{|\eta_{+-}|^2 - |\eta_{00}|^2}{|\eta_{+-}|^2} = \frac{1}{6} \left(1 - \frac{\frac{\text{Br}(K_L \rightarrow \pi^0 \pi^0)}{\text{Br}(K_S \rightarrow \pi^0 \pi^0)}}{\frac{\text{Br}(K_L \rightarrow \pi^+ \pi^-)}{\text{Br}(K_S \rightarrow \pi^+ \pi^-)}} \right)$$

Kaon & CP violation:3

$$\frac{\epsilon'_K}{\epsilon_K} = \frac{1}{\sqrt{2}|\epsilon_K|_{\text{exp}}} \frac{\omega_{\text{exp}}}{(\text{Re}A_0)_{\text{exp}}} \left(-\text{Im}A_0 + \frac{1}{\omega_{\text{exp}}} \text{Im}A_2 \right) \quad \text{where} \quad \frac{1}{\omega} \equiv \frac{\text{Re}A_0}{\text{Re}A_2} = 22.46 \text{ (exp.)}$$

■ Numerical Remarks

- $\text{Im}A_0$ ($l=0, \Delta l=1/2$) term is dominated by gluon-penguin, while $\text{Im}A_2$ ($l=2, \Delta l=3/2$) term is dominated by EW-penguins ($\propto m_t^2$), and **they have opposite sign contributions**

- Since $\text{Im}A_2$ is proportional to α but enhanced by $1/\omega$, its contribution is comparable to $\text{Im}A_0$

$$\mathcal{O}(\alpha_s) \stackrel{!}{\sim} \frac{1}{\omega} \mathcal{O}(\alpha)$$

- Two terms contribute destructively each other. Actually, ϵ'_K/ϵ_K is canceled out at $m_t \sim O(220)$ GeV [Buchalla, Buras, Harlander, 90': LO result]
- The LO QCD contribution does not contribute to $\text{Im}A_2$. Thus NLO QED corrections are *leading order* to $\text{Im}A_2$ term

Kaon & CP violation:4

- The Isospin amplitude can be decomposed into Wilson coefficients (C_i) and hadronic matrix elements ($\langle Q_i \rangle$)


$$A_{I=0,2} = \langle (\pi\pi)_{I=0,2} | \mathcal{H}_{\text{eff}}^{|\Delta S|=1} | K^0 \rangle$$

$$= \sum_i C_i \langle (\pi\pi)_{I=0,2} | Q_i | K^0 \rangle \equiv \sum_i C_i \langle Q_i \rangle_{I=0,2} \quad Q_i \text{ are four-fermi operators}$$

Composition of ϵ'_K / ϵ_K with respect to the operator basis

Negative contribution

Positive contribution




ϵ'_K / ϵ_K

[TK, Nierste, Tremper 16']

The first lattice result for $\langle Q_i \rangle$

- The calculation of the hadronic matrix elements ($\langle Q_i \rangle$), being non-perturbative quantities, is a major challenge, and have been estimated by the effective theories (e.g. chiPT, dual QCD model, NJL model, ...)
- But their results have a tension among each other (next slide)
- Recently, a determination of all hadronic matrix elements by lattice QCD is obtained **with controlled errors (first lattice result)**

[RBC-UKQCD, PRL115 (2015)]



[Figure in RBC-UKQCD, PRL115 (2015)]

- Now, one can estimate ϵ'_K/ϵ_K without using the effective theories

Current situation of $\epsilon'_K \propto \frac{\text{Im}A_0}{\text{Re}A_0} - \frac{\text{Im}A_2}{\text{Re}A_2} \propto \text{Im}A_0 - \left(\frac{\text{Re}A_0}{\text{Re}A_2}\right) \text{Im}A_2$

Bertolini, Eeg,
Fabbrichesi, Lashin 97'

Pallante, Pich 00'
Hambye,
Peris, Rafael 03'

Buras, Gerard 15'

RBC-UKQCD 15'

BGJJ 15'

Our work KNT 16'


E371(FNAL) 93'

NA31(CERN) 93'

NA48(CERN) 02'

KTeV(FNAL) 11'

PDG average



$$B_6^{(1/2)} \sim 1.6, B_8^{(3/2)} \sim 0.9$$

$$B_6^{(1/2)} \sim 1.6, B_8^{(3/2)} \sim 0.9$$

$$B_6^{(1/2)} \approx 3, B_8^{(3/2)} \approx 3.5$$

$$B_6^{(1/2)} \leq B_8^{(3/2)} \leq 1$$

$$B_6^{(1/2)} \leq B_8^{(3/2)} \leq 0.76$$

$$B_6^{(1/2)} = 0.57, B_8^{(3/2)} = 0.76$$

} Observed values

large N limit (convention)

$$B_6^{(1/2)} = B_8^{(3/2)} = 1$$

dual QCD prediction

$$B_6^{(1/2)} \leq B_8^{(3/2)} < 1, B_8^{(3/2)} = 0.8$$

$$\text{Re } \epsilon'_K / \epsilon_K \times 10^4$$

$$\left(\frac{\text{Re}A_0}{\text{Re}A_2}\right)$$

Exp.	χ PT	dual QCD	Lattice (l=0,2)
22.45 ± 0.05	~ 14	16.0 ± 1.5	31.0 ± 11.1

Singularity

J: NLO-RG evolution matrix
$$\hat{J}_s - \left[\hat{J}_s, \frac{\hat{\gamma}_s^{(0)T}}{2\beta_0} \right] = \frac{\beta_1}{\beta_0} \frac{\hat{\gamma}_s^{(0)T}}{2\beta_0} - \frac{\hat{\gamma}_s^{(1)T}}{2\beta_0},$$

- Go on the diagonalized basis of $\gamma_s^{(0)T}$, the equation becomes

$$\left(\hat{V}^{-1} \hat{J}_{s,e} \hat{V} \right)_{ij} = \frac{\dots}{2\beta_0 \mp \left((\hat{\gamma}_{s,D}^{(0)T})_{jj} - (\hat{\gamma}_{s,D}^{(0)T})_{ii} \right)}.$$

- Unfortunately, when $f=3$, $2\beta_0 = 18$, $\hat{\gamma}_{s,D}^{(0)T} \supset +2, -16$, then the denominator vanishes with a generally non-zero numerator \rightarrow Singularity
- The other J matrices also have similar singularity when $f= 3,4,5,6$

$$\hat{U}_f(\mu_1, \mu_2) = \hat{K}(\mu_1) \hat{U}_0(\mu_1, \mu_2) \hat{K}'(\mu_2),$$

with

$$\hat{K}(\mu_1) = \left(\hat{1} + \frac{\alpha_{EM}}{4\pi} \hat{J}_{se} \right) \left(\hat{1} + \frac{\alpha_s(\mu_1)}{4\pi} \hat{J}_s \right) \left(\hat{1} + \frac{\alpha_{EM}}{\alpha_s(\mu_1)} \hat{J}_e \right),$$

$$\hat{K}'(\mu_2) = \left(\hat{1} - \frac{\alpha_{EM}}{\alpha_s(\mu_2)} \hat{J}_e \right) \left(\hat{1} - \frac{\alpha_s(\mu_2)}{4\pi} \hat{J}_s \right) \left(\hat{1} - \frac{\alpha_{EM}}{4\pi} \hat{J}_{se} \right),$$

← Singularities

Removing the Singularities:1

- In order to eliminate the singularities, we generalize the Roma group's ansatz by adding a logarithmic scale dependence to the J matrices

Our singularity-free analytic solution

$$\hat{U}_f(\mu_1, \mu_2) = \hat{K}(\mu_1) \hat{U}_0(\mu_1, \mu_2) \hat{K}'(\mu_2),$$

with

$$\begin{aligned} \hat{K}(\mu_1) &= \left(\hat{1} + \frac{\alpha_{EM}}{4\pi} \hat{J}_{se}(\alpha_s(\mu_1)) \right) \left(\hat{1} + \frac{\alpha_s(\mu_1)}{4\pi} \hat{J}_s(\alpha_s(\mu_1)) \right) \\ &\quad \times \left(\hat{1} + \frac{\alpha_{EM}}{\alpha_s(\mu_1)} \hat{J}_e(\alpha_s(\mu_1)) + \left(\frac{\alpha_{EM}}{\alpha_s(\mu_1)} \right)^2 \hat{J}_{ee}(\alpha_s(\mu_1)) \right), \\ \hat{K}'(\mu_2) &= \left(\hat{1} - \frac{\alpha_{EM}}{\alpha_s(\mu_2)} \hat{J}_e(\alpha_s(\mu_2)) - \left(\frac{\alpha_{EM}}{\alpha_s(\mu_2)} \right)^2 \left(\hat{J}_{ee}(\alpha_s(\mu_2)) - \left(\hat{J}_e(\alpha_s(\mu_2)) \right)^2 \right) \right) \\ &\quad \times \left(\hat{1} - \frac{\alpha_s(\mu_2)}{4\pi} \hat{J}_s(\alpha_s(\mu_2)) \right) \left(\hat{1} - \frac{\alpha_{EM}}{4\pi} \hat{J}_{se}(\alpha_s(\mu_2)) \right), \end{aligned}$$

where

$$\begin{aligned} \hat{J}_s &\rightarrow \hat{J}_s(\alpha_s(\mu)) = \hat{J}_{s,0} + \hat{J}_{s,1} \ln \alpha_s(\mu), \\ \hat{J}_e &\rightarrow \hat{J}_e(\alpha_s(\mu)) = \hat{J}_{e,0} + \hat{J}_{e,1} \ln \alpha_s(\mu), \\ \hat{J}_{se} &\rightarrow \hat{J}_{se}(\alpha_s(\mu)) = \hat{J}_{se,0} + \hat{J}_{se,1} \ln \alpha_s(\mu) + \hat{J}_{se,2} \ln^2 \alpha_s(\mu). \\ \hat{J}_{ee} &\rightarrow \hat{J}_{ee}(\alpha_s(\mu)) = \hat{J}_{ee,0} + \hat{J}_{ee,1} \ln \alpha_s(\mu). \end{aligned}$$

[TK, Nierste, Tremper 16']

Removing the Singularities:2

- Then, J_s matrices are the solution of the following equations

$$\hat{J}_{s,0} - \left[\hat{J}_{s,0}, \frac{\hat{\gamma}_s^{(0)T}}{2\beta_0} \right] = \frac{\beta_1}{\beta_0} \frac{\hat{\gamma}_s^{(0)T}}{2\beta_0} - \frac{\hat{\gamma}_s^{(1)T}}{2\beta_0} - \hat{J}_{s,1}$$

$$\hat{J}_{s,1} - \left[\hat{J}_{s,1}, \frac{\hat{\gamma}_s^{(0)T}}{2\beta_0} \right] = 0,$$

- Overview of our solution
 - All singularity terms are regulated into **logarithmic terms**
 - Some logarithmic terms are consistent with literature
 - Our solution does not rely on a specific basis and permits a much faster, easier and, in particular, more stable computational algorithm
 - Our next-to-leading order RG evolution matrix has an additional **new** correction of $O(\alpha^2/\alpha_s^2)$, which appears only at this order

numerically $\alpha^2/\alpha_s^2 \sim \alpha$, but we find this contribution is small

ϵ'_K Discrepancy

- In the SM, ϵ'_K/ϵ_K is significantly suppressed by the GIM suppression AND by the accidental cancellation between QCD and EW penguin contributions


- SM expectation value at NLO (without effective theory) [TK, Nierste, Tremper 16']

$$\left(\frac{\epsilon'_K}{\epsilon_K}\right)_{\text{SM-NLO}} = (0.96 \pm 4.68 \pm 1.52 \pm 0.60 \pm 0.24) \times 10^{-4}$$

Lattice NNLO isospin violating mt

- We have calculated ϵ'_K/ϵ_K in the Standard Model at the next-to-leading order. The result is **2.9 sigma** below the experimental measured value. It highlights a tension between the Standard-Model prediction and experiment.

$$\text{Re}\left(\frac{\epsilon'_K}{\epsilon_K}\right)_{\text{exp}} = (16.6 \pm 2.3) \times 10^{-4}.$$



Preliminary for NP part

- The SM prediction of ϵ'_K/ϵ_K is 2.9 sigma below the experimental values, which give strong motivation for searching for NP contributions
- ϵ'_K/ϵ_K is highly sensitive to CP violation of NP

SM loop suppression *GIM suppression* accidental cancelation

VS.

NP (loop suppression) *(large coupling) * NP scale suppression

- One should also consider the other flavour constraints
- Actually, some models can explain this discrepancy, e.g. Littlest Higgs model, 331 model, generic Z' models, 750GeV model (dead?), and SUSY

[Buras,Fazio,Girrbach 14', Buras,Buttazzo,Knegjens 15, Buras 15', Buras,Fazio 15', 16', Goertz,Kamenik,Katz,Nardecchia 15', Blanke,Buras,Recksiegel 16',TK,Nierste,Tremper 16', Tanimoto, Yamamoto 16',Endo,Mishima,Ueda,Yamamoto 16']

Our calculation strategy for MSSM

- Our work

- CP violating phase in the MSSM

- CKM matrix
 - squark mass matrix
 - μ (Higgsino mass)
 - gaugino mass
 - A term



Included




take to be Real
in light of severe constraint
from EDM experiments

- We calculate SUSY QCD (gluino) corrections and chargino/neutralino-Z penguin contribution in light of strong coupling and Isospin symmetry breaking
 - TeV scale SUSY & SUSY scale matching, mass eigenbasis calc., NLO-QCD and QED RGE corrections


Gluino box (“Trojan penguin”)

[Kagan, Neubert, PRL83(1999),
Grossman, Kagan, Neubert, JHEP10(1999)]

- In spite of QCD correction, gluino box diagram **can** break isospin symmetry through mass difference between right-handed squark masses
- *“It is neither (pure) penguins nor of electroweak origin. Nevertheless, at low energies their effects are parameterized by an extension of the usual basis of electroweak penguin operators.”*




Movie: Penguins of Madagascar




Sub leading contributions

- Gluino chromomagnetic penguin operator can give subleading contribution, but there is no reliable results for hadronic matrix element
[Buras,Colangelo,Ishidori,Romanino,Silvestrini,00']
- Gluino photon-penguin breaks isospin sym. explicitly, but is suppressed by α/α_s
[Langacker,Sathiapalan,84',Grossman,Worah,97',Abel,Cottingham,Whittingham,98']
- Z-penguin contribution needs to break the EW sym. like $\mathcal{L}_{\text{eff}} = \frac{\lambda_{ij}}{M^2} |H|^2 \bar{d}_i \not{D} d_j$,
Hence, chargino Z-penguin contribution is always larger than gluino Z-penguin
[Colangelo,Isidori,98'@K→πνν]



Overview for calculation of SUSY ϵ'_K

- We calculated the following six-type one-loop SUSY contributions
- SUSY matching scale is given as the input parameter



Main Constraint: ϵ_K ($\Delta S=2$, ID-CPV)

- Although ϵ'_K ($\Delta S=1$, D-CPV) is sensitive to NP, once ϵ_K ($\Delta S=2$, ID-CPV) constraint is taken into account, NP effects in $\Delta S=1$ is highly suppressed
- NP hierarchy in $|\Delta S| = 1$ vs. $|\Delta S| = 2$ transitions;

$$\epsilon_K^{\text{SM}} \propto \frac{\text{Im}(\tau^2)}{M_W^2} \quad \epsilon_K^{\prime\text{SM}} \propto \frac{\text{Im}\tau}{M_W^2} \quad \tau = -\frac{V_{td}V_{ts}^*}{V_{ud}V_{us}^*} \sim (1.5 - i0.6) \cdot 10^{-3}$$

If the NP contribution comes with the $\Delta S = 1$ parameter δ and is mediated by heavy particles of mass M , one finds

$$\epsilon_K^{\text{NP}} \propto \frac{\text{Im}(\delta^2)}{M^2} \quad \epsilon_K^{\prime\text{NP}} \propto \frac{\text{Im}\delta}{M^2}$$


$$\frac{\epsilon_K^{\prime\text{NP}}}{\epsilon_K^{\prime\text{SM}}} \leq \frac{\frac{\epsilon_K^{\text{NP}}}{\epsilon_K^{\text{SM}}}}{\frac{\epsilon_K^{\prime\text{SM}}}{\epsilon_K^{\text{SM}}}} = \mathcal{O}\left(\frac{\text{Re}\tau}{\text{Re}\delta}\right)$$

\uparrow
 $\epsilon_K^{\text{NP}} \leq \epsilon_K^{\text{SM}}$

With $M > 1$ TeV, NP effects can only be relevant for $|\delta| \gg |\tau|$ and this equation seemingly forbids detectable NP contributions to ϵ'_K

Loophole of constraint from ϵ_K



- The leading contribution is given by $\overline{d}_L s_L \overline{d}_R s_R$



$$\propto \left(\frac{m_K}{m_s + m_d} \right)^2$$

this contribution is suppressed when $\Delta_{\bar{D},12} \simeq 0$

- The next contribution is given by $\overline{d}_L s_L \overline{d}_L s_L$




Crossed diagram gives relatively negative contributions

- $m_{\tilde{g}} \gtrsim 1.5 m_{\tilde{q}}$, these contributions almost cancel out [Crivellin, Davidkov, PRD81(2010)]

Constraint from ϵ_K

$M_S = 10 \text{ TeV}$



[TK, Nierste, Tremper, PRL(2016)]



- Actually, there are several expected values of ϵ_K depending on the input CKM parameters
 - $|V_{cb}|$; measured in inclusive $b \rightarrow cl\nu$ decays..... ϵ_K is consistent with exp. value
 - $|V_{cb}|$; measured in exclusive $B \rightarrow D(*)l\nu$ decays..... ϵ_K is 3σ below the exp. value

SUSY contributions to ϵ'_K

- We take universal SUSY mass spectrum without gauginos and right-handed up-type squark mass




[TK, Nierste, Tremper, PRL(2016)]

- ✓ nEDM, ΔM_K , DDbar mixing are weaker constraints than ϵ_K

Conclusions

- ϵ'_K/ϵ_K is a good measure of the CP violation from new physics
- The lattice group and the SM calculations have revealed that the SM expected value deviates significantly from exp. data ($\sim 3\sigma$)
- In the MSSM, gluino box diagram with mass different of the right-handed squark contributes ϵ'_K/ϵ_K significantly
- Heavy gluino can relax the constraint from ϵ_K
- Prospects
 - Correlation with other hadronic channels
 - Higher order corrections: e.g. 2-loop gluino box
 - UV model, GUT?
 - Large A scenario, vacuum stability

TK, Nierste, Tremper, Endo, Mishima, Yamamoto(K) STAY TUNED



made by
Philipp Frings

Backup



Numerical results

- Wilson coefficients @ $\mu = 1.3$ GeV $C_i(\mu) \equiv z_i(\mu) - \frac{V_{ts}^* V_{td}}{V_{us}^* V_{ud}} y_i(\mu)$ new results

i	$z_i(\mu)$	$y_i(\mu)$	$\mathcal{O}(1)$	$\mathcal{O}(\alpha_{EM}/\alpha_s)$	$\mathcal{O}(\alpha_s)$	$\mathcal{O}(\alpha_{EM})$	$\mathcal{O}(\alpha_{EM}^2/\alpha_s^2)$
1	-0.3903	0	0	0	0	0	0
2	1.200	0	0	0	0	0	0
3	0.0044	0.0274	0.0254	0.0001	0.0007	0.0012	0
4	-0.0131	-0.0566	-0.0485	-0.0003	-0.0069	-0.0009	0
5	0.0039	0.0068	0.0124	0.0001	-0.0059	0.0001	0
6	-0.0128	-0.0847	-0.0736	-0.0003	-0.0099	-0.0008	0
$7/\alpha_{EM}$	0.0042	-0.0344	0	-0.1120	0	0.0757	0.0019
$8/\alpha_{EM}$	0.0020	0.1158	0	-0.0222	0	0.1373	0.0007
$9/\alpha_{EM}$	0.0053	-1.3834	0	-0.1269	0	-1.2582	0.0017
$10/\alpha_{EM}$	-0.0013	0.4877	0	0.0214	0	0.4668	-0.0004

- Hadronic matrix elements @ $\mu = 1.3$ GeV

i	$\langle Q_i(\mu) \rangle_0^{\text{MS-NDR}} (\text{GeV})^3$	i	$\langle Q_i(\mu) \rangle_2^{\text{MS-NDR}} (\text{GeV})^3$
1	-0.145 ± 0.046	1	0.01006 ± 0.00002
2	0.105 ± 0.015	2	0.01006 ± 0.00002
3	-0.041 ± 0.066	3	—
4	0.209 ± 0.066	4	—
5	-0.180 ± 0.068	5	—
6	-0.342 ± 0.122	6	—
7	0.160 ± 0.065	7	0.135 ± 0.012
8	1.556 ± 0.376	8	0.874 ± 0.054
9	-0.197 ± 0.069	9	0.01509 ± 0.00003
10	0.053 ± 0.037	10	0.01509 ± 0.00003

Lattice simulation is calculated at $\mu=1.5$ GeV ($l=0$) and $\mu=3.0$ GeV ($l=2$) with 2+1 flavour

We exploit CP-conserving data (with z_i) to reduce hadronic uncertainties

[TK, Nierste, Tremper 16']

Overview of effective models

- Chiral perturbation theory

- Effective theory of the QCD Goldstone bosons: $\Phi = \begin{pmatrix} \sqrt{\frac{1}{2}}\pi^0 + \sqrt{\frac{1}{6}}\eta & \pi^+ & K^+ \\ \pi^- & -\sqrt{\frac{1}{2}}\pi^0 + \sqrt{\frac{1}{6}}\eta & K^0 \\ K^- & \bar{K}^0 & -\sqrt{\frac{2}{3}}\eta \end{pmatrix}$

$$\mathcal{L} = -\frac{G_F}{\sqrt{2}} V_{ud} V_{us}^* \left(g_8 f^4 \text{tr} (\lambda L_\mu L^\mu) + g_{27} f^4 \left(L_{\mu 23} L_{11}^\mu + \frac{2}{3} L_{\mu 21} L_{13}^\mu \right) + \mathcal{O}(g_E W) \right)$$

with $L_\mu = -iU^\dagger D_\mu U$ $U = \exp \left(i \frac{\sqrt{2}\Phi}{f} \right)$

- dual QCD method [Bardeen, Buras, Gerard 87', 14']

- Effective theory of the truncated pseudo-scalar and vector mesons:

$$\mathcal{L} = \frac{f^2}{4} \text{tr} (\partial_\mu U \partial^\mu U^\dagger) - \frac{1}{4} \text{tr} (V_{\mu\nu} V^{\mu\nu}) - \frac{f^2}{2} \text{tr} (\partial_\mu \xi^\dagger \xi + \partial_\mu \xi \xi^\dagger - 2igV_\mu)^2 \quad \text{with} \quad U = \xi \xi$$

- Chiral quark model

- Mean-field approximation of the full extended NJL model

$$\mathcal{L} = \mathcal{L}_{QCD} - M (\bar{q}_R U q_L + \bar{q}_L U^\dagger q_R)$$