

The 3rd NPKI Workshop @ Seoul

*"The lesson from the first results
of Run 2 of the LHC"*

Highlights and Prospects from LHCb

Tim Gershon
University of Warwick



NPKI 2016

15th June 2016

The LHCb detector

- In high energy collisions, $b\bar{b}$ pairs produced predominantly in forward or backward directions
- LHCb designed as a forward spectrometer

The LHCb Detector
JINST 3 (2008) S08005



Precision primary and secondary vertex measurements

Excellent K/π separation capability

The LHCb Run 1 trigger

JINST 8 (2013) P04022

Challenge is


- to efficiently select most interesting events
- while maintaining manageable data rates

Main backgrounds

- “minimum bias” inelastic pp scattering
- other charm and beauty decays



Handles

- high p_T signals (muons)
- displaced vertices



Run II data taking

- At 13 TeV, LHCb's flavour physics programme gains from higher \sqrt{s} (increased production) and 25 ns bunch spacing (lower pile up)
- During LS1: some subdetector consolidation; new HERSCHEL forward shower counters; change of data flow in trigger




First results from Run II


Open charm production

JHEP 03 (2016) 159


$D \rightarrow K\pi$




$D_s \rightarrow KK\pi$



$D^* \rightarrow D\pi; D \rightarrow K\pi$





Increases of production cross-section from $\sqrt{s} = 7 \rightarrow 13$ TeV at upper end of range of expectation



First results from Run II



J/ψ production

JHEP 10 (2015) 172




high- p_T
b-hadrons tend to have lower background & better tagging

Increases from $\sqrt{s} = 7,8 \rightarrow 13 \text{ TeV}$ at upper end of range of expectation



Data taken so far in Run II

LHCb Integrated Luminosity at p-p in 2016




Not only flavour physics ...

- Most of the recent results from LHCb are on its “core” flavour physics programme
 - CP violation, the Unitarity Triangle and rare B decays
- LHCb also has unique non-flavour capability
 - Top production in the forward region (PRL 115 (2015) 112001)
 - Determination of $\sin^2\theta_W$ (JHEP 11 (2015) 190)
 - Search for hidden sector bosons (PRL 115 (2015) 161802)
 - Ideas to search for dark photons ...

Proposals for dark photon searches at LHCb

arXiv:1509.06765, arXiv:1603.08926



CP violation & the Unitarity Triangle

The Unitarity Triangle


- The CKM matrix must be unitary

$$V_{CKM}^+ V_{CKM} = V_{CKM} V_{CKM}^+ = 1$$


- Provides numerous tests of constraints between independent observables, such as

$$|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1$$

$$V_{ud} V_{ub}^* + V_{cd} V_{cb}^* + V_{td} V_{tb}^* = 0$$



<http://ckmfitter.in2p3.fr>
see also <http://www.utfit.org>



Consistency of measurements tests the Standard Model and provides model-independent constraints on New Physics

$|V_{ub}/V_{cb}|$ from $\Lambda_b \rightarrow p\mu\nu/\Lambda_b \rightarrow \Lambda_c\mu\nu$

Nature Phys. 11 (2015) 743

- Long standing discrepancy between exclusive and inclusive determinations of both V_{ub} and V_{cb}


PDG 2014

$$|V_{cb}| = (42.4 \pm 0.9) \times 10^{-3} \text{ (inclusive)} \quad |V_{ub}| = (4.41 \pm 0.15 \text{ } ^{+0.15}_{-0.17}) \times 10^{-3} \text{ (inclusive),}$$

$$|V_{cb}| = (39.5 \pm 0.8) \times 10^{-3} \text{ (exclusive)} \quad |V_{ub}| = (3.23 \pm 0.31) \times 10^{-3} \text{ (exclusive).}$$

- Use of b baryon decays provides complementary alternative to B mesons
- At LHCb, exploit displaced vertex to reconstruct corrected mass

$$M_{corr} = \sqrt{p_{\perp}^2 + M_{p\mu}^2} + p_{\perp}$$






$$|V_{ub}/V_{cb}| \text{ from } \Lambda_b \rightarrow p\mu\nu/\Lambda_b \rightarrow \Lambda_c\mu\nu$$

- Can then reconstruct $q^2 = m(\mu\nu)^2$
 - Select events with $q^2 > 15 \text{ GeV}^2$
 - Highest rate, best resolution & most reliable theory (lattice) predictions
- Use isolation MVA to suppress background
- Fit M_{corr} to obtain signal yields

- Rules out models with RH currents
- Compatible with UT fit (β, γ)

Nature Phys. 11 (2015) 743



$$\frac{\mathcal{B}(\Lambda_b \rightarrow p\mu^-\bar{\nu}_\mu)_{q^2 > 15 \text{ GeV}^2/c^4}}{\mathcal{B}(\Lambda_b \rightarrow \Lambda_c\mu\nu)_{q^2 > 7 \text{ GeV}^2/c^4}} = (1.00 \pm 0.04(\text{stat}) \pm 0.08(\text{syst})) \times 10^{-2}$$

$$\frac{|V_{ub}|}{|V_{cb}|} = 0.083 \pm 0.004(\text{expt}) \pm 0.004(\text{lattice})$$



$|V_{td}/V_{ts}|$ from $\Delta m_d/\Delta m_s$

arXiv:1604.03475

- Δm_s now precisely known
- limitation on knowledge of UT side from lattice (improving fast) and Δm_d
- new measurement uses $B^0 \rightarrow D^{(*)-} \mu \nu$ decays

$\Delta m_s = 17.768 \pm 0.023 \pm 0.006 \text{ ps}^{-1}$ (LHCb NJP 15 (2013) 053021)

latest lattice calculations: arXiv:1603.04306, arXiv:1602.03560



$$\Delta m_d = (505.0 \pm 2.1 \text{ (stat)} \pm 1.0 \text{ (syst)}) \text{ ns}^{-1}$$

single most precise determination
precision of previous world average


Importance of γ from $B \rightarrow DK$

- γ plays a unique role in flavour physics

the only CP violating parameter that can be measured through tree decays (*)

(*) more-or-less

- A benchmark Standard Model reference point
 - doubly important after New Physics is observed




Variants use different B or D decays
require a final state common to both D^0 and \bar{D}^0

γ from $B^+ \rightarrow DK^+$, $D \rightarrow KK, \pi\pi, K\pi$

arXiv:1603.08993


$D \rightarrow K\pi$ (favoured)



small asymmetries due to production and detection effects

$B \rightarrow D\pi$ control mode helps to separate effects

$D \rightarrow \pi K$ (“ADS” suppressed)




large CP violating asymmetries – first 5σ observation in a single $B \rightarrow DK$ channel

effects also possible in $B \rightarrow D\pi$


γ from $B^+ \rightarrow DK^+$, $D \rightarrow KK, \pi\pi, K\pi$

arXiv:1603.08993

$D \rightarrow \pi\pi$ ("GLW" CP+ state)




$D \rightarrow KK$ ("GLW" CP+ state)



CP violating asymmetries visible
but not 5σ significant

γ from $B^+ \rightarrow DK^+$, $D \rightarrow KK, \pi\pi, K\pi$

arXiv:1603.08993



Measurements reaching percent level precision

Some tension in the A_{CP+} average ($\chi^2 = 16/4$ dof) but no other sign of experimental disagreements

γ from $B^0 \rightarrow DK^{*0}$, $D \rightarrow K_S \pi\pi$, $K_S KK$


arXiv:1604.01525, arXiv:1605.01082

$B^0 \rightarrow DK^{*0}$ rarer, but with larger interference effects, than $B^+ \rightarrow DK^+$


$D \rightarrow KK, \pi\pi, K\pi$ previously studied in PR D90 (2014) 112002

Now consider “GGSZ” modes with both model-independent (arXiv:1604.01525) and -dependent (arXiv:1605.01082) analyses

$D \rightarrow K_S \pi\pi$ (both MI & MD)



$D \rightarrow K_S KK$ (MI only)




B_S^0 decays to same final states provide control channels

γ from $B^0 \rightarrow DK^{*0}$

arXiv:1604.01525, arXiv:1605.01082

For $B^0 \rightarrow DK^{*0}$, width of the K^{*0} resonance introduces a dilution factor that depends on the $B^0 \rightarrow DK^+\pi^-$ Dalitz plot

This has been studied with $D \rightarrow K\pi$ (PRD 92 (2015) 012012), KK and $\pi\pi$ (arXiv:1602.03455) decays






Interference effects in the $D_2^*-K^*$ overlap region enhance sensitivity to γ

γ from $B^0 \rightarrow DK^{*0}$

arXiv:1604.01525

arXiv:1605.01082

arXiv:1602.03455



$D \rightarrow K_S \pi \pi$

$D \rightarrow KK, \pi \pi$

Comparison of results in terms of $x_{\pm} = r_B \cos(\delta_{B^{\pm}} \pm \gamma)$, $y_{\pm} = r_B \sin(\delta_{B^{\pm}} \pm \gamma)$

RED: (x_+, y_+) , BLUE (x_-, y_-)

γ combination

LHCb-CONF-2016-001

Many observables with sensitivity to γ

- $B^+ \rightarrow DK^+$, $D \rightarrow h^+h^-$, GLW/ADS, 3 fb^{-1} [4]
- $B^+ \rightarrow DK^+$, $D \rightarrow h^+\pi^-\pi^+\pi^-$, quasi-GLW/ADS, 3 fb^{-1} [4]
- $B^+ \rightarrow DK^+$, $D \rightarrow h^+h^-\pi^0$, quasi-GLW/ADS, 3 fb^{-1} [5]
- $B^+ \rightarrow DK^+$, $D \rightarrow K_s^0 h^+h^-$, model-independent GGSZ, 3 fb^{-1} [6]
- $B^+ \rightarrow DK^+$, $D \rightarrow K_s^0 K^+\pi^-$, GLS, 3 fb^{-1} [7]
- $B^0 \rightarrow DK^+\pi^-$, $D \rightarrow h^+h^-$, GLW-Dalitzz, 3 fb^{-1} [8]
- $B^0 \rightarrow DK^{*0}$, $D \rightarrow K^+\pi^-$, ADS, 3 fb^{-1} [9]
- $B^0 \rightarrow DK^{*0}$, $D \rightarrow K_s^0 \pi^+\pi^-$, model-dependent GGSZ, 3 fb^{-1} [10]
- $B^+ \rightarrow DK^+\pi^+\pi^-$, $D \rightarrow h^+h^-$, GLW/ADS, 3 fb^{-1} [11]
- $B_s^0 \rightarrow D_s^\mp K^\pm$, time-dependent, 1 fb^{-1} [12],

[4] LHCb collaboration, R. Aaij *et al.*, *Measurement of CP observables in $B^\pm \rightarrow DK^\pm$ and $B^\pm \rightarrow D\pi^\pm$ with two- and four-body D meson decays*, LHCb-PAPER-2016-003, in preparation.

[5] LHCb collaboration, R. Aaij *et al.*, *A study of CP violation in $B^\mp \rightarrow Dh^\mp$ ($h = K, \pi$) with the modes $D \rightarrow K^\mp \pi^\pm \pi^0$, $D \rightarrow \pi^+\pi^-\pi^0$ and $D \rightarrow K^+K^-\pi^0$* , *Phys. Rev. D* **91** (2015) 112014, [arXiv:1504.05442](#)

[6] LHCb collaboration, R. Aaij *et al.*, *Measurement of the CKM angle γ using $B^\pm \rightarrow DK^\pm$ with $D \rightarrow K_s^0 \pi^+\pi^-$, $K_s^0 K^+K^-$ decays*, *JHEP* **10** (2014) 097, [arXiv:1408.2748](#)

[7] LHCb collaboration, R. Aaij *et al.*, *A study of CP violation in $B^\pm \rightarrow DK^\pm$ and $B^\pm \rightarrow D\pi^\pm$ decays with $D \rightarrow K_s^0 K^\pm \pi^\mp$ final states*, *Phys. Lett. B* **733** (2014) 36, [arXiv:1402.2982](#)

[8] LHCb collaboration, R. Aaij *et al.*, *Constraints on the unitarity triangle angle γ from Dalitz plot analysis of $B^0 \rightarrow DK^+\pi^-$ decays*, LHCb-PAPER-2015-059, submitted to *Phys. Rev. Lett.*

[9] LHCb collaboration, R. Aaij *et al.*, *Measurement of CP violation parameters in $B^0 \rightarrow DK^{*0}$ decays*, *Phys. Rev. D* **90** (2014) 112002, [arXiv:1407.8136](#)

[10] LHCb collaboration, R. Aaij *et al.*, *Measurement of the CKM angle γ using $B^0 \rightarrow DK^{*0}$ with $D \rightarrow K_s^0 \pi^+\pi^-$ decays*, LHCb-PAPER-2016-007, in preparation.

[11] LHCb collaboration, R. Aaij *et al.*, *Study of $B^- \rightarrow DK^-\pi^+\pi^-$ and $B^- \rightarrow D\pi^-\pi^+\pi^-$ decays and determination of the CKM angle γ* , *Phys. Rev. D* **92** (2015) 112005, [arXiv:1505.07044](#)


[12] LHCb collaboration, R. Aaij *et al.*, *Measurement of CP asymmetry in $B_s^0 \rightarrow D_s^\mp K^\pm$ decays*, *JHEP* **11** (2014) 060, [arXiv:1407.6127](#)

New results discussed on previous slides

γ combination


LHCb-CONF-2016-001

Many observables with sensitivity to γ



$$\gamma = (70.9^{+7.1}_{-8.5})^\circ$$


World average including BaBar, Belle, CDF results will give marginally better precision



CP violation in $B^0_{(s)}$ mixing

arXiv:1605.09768



- Evidence of non-SM CP violation in inclusive dimuon asymmetry from the D0 collaboration
 - PRD 89 (2014) 012002
- Semileptonic asymmetries $a_{sl}(B^0_{(s)})$ and $a_{sl}(B^0_s)$ however consistent with SM $\sim (0,0)$
 - $a_{sl}(B^0)$ by BaBar, Belle, LHCb, D0
 - $a_{sl}(B^0_s)$ by LHCb (new), D0
- Possibility of additional contributions to inclusive dimuon asymmetry under investigation
 - PR D87 (2013) 074020



$$a_{sl}(B^0_s) = (0.39 \pm 0.26 \pm 0.20)\%$$

Limits on BSM contributions to $\Delta B=2$

Define $M_{12}^q = M_{12}^{\text{SM},q} \Delta_q$ and obtain constraints on $(\text{Re } \Delta_q, \text{Im } \Delta_q)$
 (here not including anomalous D0 dimuon asymmetry result, and other recent results)




Rare (and some not so rare) decays

$$B_s \rightarrow \mu^+ \mu^-$$

Killer app. for new physics discovery

Very rare in Standard Model due to


- absence of tree-level FCNC
- helicity suppression
- CKM suppression
- ... all features which are not necessarily reproduced in extended models




$$B(B_s \rightarrow \mu^+ \mu^-)^{SM} = (3.66 \pm 0.23) \times 10^{-9}$$

$$B(B_s \rightarrow \mu^+ \mu^-)^{MSSM} \sim \tan^6 \beta / M_{A0}^4$$

Intensively searched for over 30 years!




$$B_s \rightarrow \mu^+ \mu^-$$



Combination of CMS and LHCb data results in first observation of $B_s \rightarrow \mu^+\mu^-$ and first evidence for $B^0 \rightarrow \mu^+\mu^-$

Results consistent with SM at 2 σ level



Recent results from ATLAS (not included here) have almost similar sensitivity



Full angular analysis of $B^0 \rightarrow K^{*0} \mu^+ \mu^-$

JHEP 02 (2016) 104


- $B^0 \rightarrow K^{*0} \mu^+ \mu^-$ provides superb laboratory to search for new physics in $b \rightarrow s l^+ l^-$ FCNC processes
 - rates, angular distributions and asymmetries sensitive to NP
 - **experimentally clean signature**
 - many kinematic variables ... **with clean theoretical predictions**
- Full set of observables measured – only a subset shown



Tension with SM in the P_5' observable

JHEP 02 (2016) 104

- Dimuon pair is predominantly spin-1
 - either vector (V) or axial-vector (A)
- There are 6 non-negligible amplitudes
 - 3 for VV and 3 for VA ($K^{*0}\mu^+\mu^-$)
 - expressed as $A_{0,\perp,\parallel}^{L,R}$ (transversity basis)



- P_5' related to difference between relative phase of longitudinal (0) and perpendicularly (\perp) polarised amplitudes for VV and VA
 - constructed so as to minimise form-factor uncertainties



$$P_5' = \sqrt{2} \frac{\text{Re} (A_0^L A_{\perp}^{L*} - A_0^R A_{\perp}^{R*})}{\sqrt{(|A_0^L|^2 + |A_0^R|^2) (|A_{\parallel}^L|^2 + |A_{\parallel}^R|^2 + |A_{\perp}^L|^2 + |A_{\perp}^R|^2)}}$$

Sensitive to NP in V or A couplings (Wilson coefficients $C_9^{(i)}$ & $C_{10}^{(i)}$)

$$B_s \rightarrow \phi \mu^+ \mu^-$$

JHEP 09 (2015) 179


- Full angular analysis performed
- Not self-tagging → complementarity to $K^{*0} \mu^+ \mu^-$
 - only a subset of many observables shown



Tension in branching fraction, but angular observables consistent with SM

Global fit to Wilson coefficients

(slide from Sebastian Descotes-Genon @ FPCP 2016)





- p-value=71% (goodness of fit), $\text{pull}_{SM} = 4.5\sigma$ (metrology)
- BRs and angular obs both favour $C_9^{NP} \simeq -1$ in all “good” scenarios

Lepton universality – R_K

PRL 113 (2014) 151601

Deficit of $B \rightarrow K\mu^+\mu^-$ compared to expectation
also seen in $K\mu^+\mu^-/Ke^+e^-$ ratio (R_K)

Example mass fit for Ke^+e^-
Note huge tail due to energy loss





$$R_K(1 < q^2 < 6 \text{ GeV}^2) = 0.745^{+0.090}_{-0.074} \pm 0.036$$

$B \rightarrow D^{(*)} \tau \nu$

- Powerful channel to test lepton universality
 - ratios $R(D^{(*)}) = B(B \rightarrow D^{(*)} \tau \nu) / B(B \rightarrow D^{(*)} \mu \nu)$ could deviate from SM values, e.g. in models with charged Higgs
- Heightened interest in this area
 - anomalous results from BaBar
 - other hints of lepton universality violation, e.g. R_K , $H \rightarrow \tau \mu$



PRL 109 (2012) 101802
& PRD 88 (2013) 072012



B → D*τν at LHCb

PRL 115 (2015) 112001


- Identify $B \rightarrow D^*\tau\nu$, $D^* \rightarrow D\pi$, $D \rightarrow K\pi$, $\tau \rightarrow \mu\nu\bar{\nu}$
 - Similar kinematic reconstruction to $\Lambda_b \rightarrow p\mu\nu$
 - Assume $p_{B,z} = (p_{D^*} + p_{\mu})_z$ to calculate $M_{\text{miss}}^2 = (p_B - p_{D^*} - p_{\mu})^2$
 - Require significant B, D, τ flight distances & use isolation MVA
- Separate signal from background by fitting in M_{miss}^2 , q^2 and E_{μ}
 - Shown below high q^2 region only (best signal sensitivity)



$$R(D^*) = 0.336 \pm 0.027 \pm 0.030$$

$B \rightarrow D^{(*)}TV$

Tension with SM at 4.0σ



Careful averaging needed to account for statistical and systematic correlations

Prospects

- **Data-taking progressing well**
 - Expect to collect ~5/fb of 13 TeV data during Run II
 - Improve current precision by at least a factor of 2
- **During LS2 (2019-20) will install upgraded detector**
 - Will allow higher luminosity and improved trigger efficiency
 - Designed to accumulate 50/fb in ~5 years of operation
- **Possibilities for subsequent upgrade under discussion**
 - During LS3 (concomitant with HL-LHC upgrades) to extend capability (e.g. additional tracking coverage, calorimeter replacement)
 - During LS4 to allow significantly higher luminosity and/or alternative physics programme (e.g. $H \rightarrow c\bar{c}$)
 - Your ideas welcome!


The 3rd NPKI Workshop @ Seoul

"The lesson from the first results of Run 2 of the LHC"

- LHCb surpassed Run I performance expectations
 - huge physics output, in “core” flavour observables but also much more
 - **modes with neutrinos, previously thought to be impossible**
 - ... and don't forget pentaquarks (and other topics not covered today)
 - **several potential hints of BSM effects to be explored further**
- Important improvements in the trigger for Run II
- Data taking going well so far
 - first physics papers on Run II data already submitted
 - much to look forward to!
- **Beyond Run II will install LHCb upgrade to enable even high luminosity**
 - **also starting to think of even longer term possibilities**



Quark flavour mixing a.k.a. CKM phenomenology




$$V_{CKM} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} = \begin{pmatrix} 1 - \frac{1}{2}\lambda^2 & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \frac{1}{2}\lambda^2 & A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{pmatrix} + O(\lambda^4)$$

- CKM theory is highly predictive
 - huge range of phenomena over a massive energy scale predicted by only 4 independent parameters (+ G_F + m_q + QCD)
- CKM matrix is hierarchical
 - distinctive flavour sector of Standard Model not necessarily replicated in extended theories → strong constraints on NP models
- CKM mechanism introduces CP violation
 - only source of CP violation in the Standard Model ($m_\nu = \theta_{QCD} = 0$)

Two routes to heaven


for quark flavour physics



CP violation
(extra sources must exist)

But


- No guarantee of the scale
- No guarantee of effects in the quark sector
- Realistic prospects for CPV measurement in vs due to large θ



Rare decays
(strong theoretical arguments)

But

- How high is the NP scale?
- Why have FCNC effects not been seen?






Absence of clear NP signals at ATLAS/CMS → argument for searches via rare decays stronger

Search for hidden sector bosons

Search for narrow $\mu\mu$ peak in $B \rightarrow K^{*0}\mu\mu$ decays corresponding to χ with either negligible or non-negligible lifetime



PRL 115 (2015) 161802



No significant peak away from known resonances

$|V_{ub}/V_{cb}|$ from $\Lambda_b \rightarrow p\mu\nu/\Lambda_b \rightarrow \Lambda_c\mu\nu$

Nature Phys. 11 (2015) 743



$$\frac{\mathcal{B}(\Lambda_b \rightarrow p\mu^-\bar{\nu}_\mu)_{q^2 > 15 \text{ GeV}^2/c^4}}{\mathcal{B}(\Lambda_b \rightarrow \Lambda_c\mu\nu)_{q^2 > 7 \text{ GeV}^2/c^4}} = (1.00 \pm 0.04(\text{stat}) \pm 0.08(\text{syst})) \times 10^{-2}$$


$$\frac{|V_{ub}|}{|V_{cb}|} = 0.083 \pm 0.004(\text{expt}) \pm 0.004(\text{lattice})$$

- Rules out models with RH currents
- Compatible with UT fit (β, γ)





γ combination

LHCb-CONF-2016-001

Many observables with sensitivity to γ



$$\gamma = (70.9^{+7.1}_{-8.5})^\circ$$





World average including BaBar, Belle, CDF results will give marginally better precision

Charm mixing with $D \rightarrow K\pi\pi\pi$

LHCb-PAPER-2015-057

Multibody charm decays also of interest to study charm oscillations (also to constrain hadronic parameters needed in the γ fit)



Charm mixing parameters $< 1\%$



Still not established whether
 $x \equiv \Delta m_D / \Gamma_D \neq 0$

Charm CP violation

LHCb-PAPER-2015-055

No evidence for CP violation in the charm system, whether in mixing, decay or mixing-decay interference

Latest: $\Delta A_{CP} \equiv A_{CP}(D \rightarrow KK) - A_{CP}(D \rightarrow \pi\pi) = (-0.10 \pm 0.08 \pm 0.03) \%$





Much stronger constraints obtained with minimal assumption on CPV in decays

Charm CP violation

LHCb-PAPER-2015-055

No evidence for CP violation in the charm system, whether in mixing, decay or mixing-decay interference

Latest: $\Delta A_{CP} \equiv A_{CP}(D \rightarrow KK) - A_{CP}(D \rightarrow \pi\pi) = (-0.10 \pm 0.08 \pm 0.03) \%$





Much stronger constraints obtained with minimal assumption on CPV in decays

B^0 and B_s^0 mixing phases: $\sin(2\beta)$ & ϕ_s

PRL 115 (2015) 031601

$\sin(2\beta) \equiv \sin(2\phi_1)$ **HFAG**
Moriond 2015
PRELIMINARY




LHCb: PRL 114 (2015) 041801;
PL B736 (2014) 186;
ATLAS: arXiv:1601.03297;
CMS: PL B757 (2016) 97

Possible penguin pollution controlled by SU(3) partners
LHCb: PL B742 (2015) 38, JHEP 11 (2015) 082

$$B_s \rightarrow \mu^+ \mu^-$$


ATLAS preliminary
Moriond 2016



Cleanest of 3 BDT bins

Able to distinguish B^0 and B_s^0 peaks

$B(B^0 \rightarrow \mu^+ \mu^-) < 4.2 \times 10^{-10}$ at 95% CL



compatibility
of the simultaneous fit
with the SM:
p-value = 0.048 (2.0σ)

$$B(B_s^0 \rightarrow \mu^+ \mu^-) = 0.9_{-0.8}^{+1.1} \times 10^{-9}$$

$$B(B_s^0 \rightarrow \mu^+ \mu^-) [10^{-9}]$$


Sensitivity comparable
to CMS and LHCb
One to watch in Run 2

Full angular analysis of $B^0 \rightarrow K^{*0} \mu^+ \mu^-$

JHEP 02 (2016) 104

Comparison to other experiments
(until now, only LHCb does a full angular analysis)


CMS (PLB 753 (2016) 424) quite competitive,
especially at high q^2




Beyond Run II – the LHCb Upgrade

- Beyond LHC Run II, the data-doubling time for LHCb becomes too long
 - Due to 1 MHz readout limitation and associated hardware (L0) trigger
- However, there is an excellent physics case to push for improved precision and an ever-broader range of observables
- **Will upgrade the LHCb detector in the LHC LS2 (2018-20)**
 - Upgrade subdetector electronics to 40 MHz readout
 - Make all trigger decisions in software
 - Operation at much higher luminosity with improved efficiency
 - order of magnitude improvement in precision (compared to today)
- Upgrade will be performed during LSII (now expected to be 2019-20)
 - Restart data taking in 2021 at instantaneous luminosity up to $2 \cdot 10^{33}/\text{cm}^2/\text{s}$
 - Upgrade detector qualified to accumulate 50/fb

LHC upgrade and the all important trigger





higher luminosity
 → need to cut harder at L0 to keep rate at 1 MHz
 → lower efficiency




- readout detector at 40 MHz
- implement trigger fully in software → efficiency gains
- run at L_{inst} up to $2 \cdot 10^{33} / \text{cm}^2 / \text{s}$

LHC upgrade and the all important trigger



- readout detector at 40 MHz
- implement trigger fully in software → efficiency gains
- run at L_{inst} up to $2 \cdot 10^{33}/\text{cm}^2/\text{s}$

LHCb detector upgrade



LHCb & upgrade sensitivities

Table 28: Statistical sensitivities of the LHCb upgrade to key observables. For each observable the expected sensitivity is given for the integrated luminosity accumulated by the end of LHC Run 1, by 2018 (assuming 5 fb^{-1} recorded during Run 2) and for the LHCb Upgrade (50 fb^{-1}). An estimate of the theoretical uncertainty is also given – this and the potential sources of systematic uncertainty are discussed in the text.

Type	Observable	LHC Run 1	LHCb 2018	LHCb upgrade	Theory
B_s^0 mixing	$\phi_s(B_s^0 \rightarrow J/\psi \phi)$ (rad)	0.050	0.025	0.009	~ 0.003
	$\phi_s(B_s^0 \rightarrow J/\psi f_0(980))$ (rad)	0.068	0.035	0.012	~ 0.01
	$A_{\text{sl}}(B_s^0)$ (10^{-3})	2.8	1.4	0.5	0.03
Gluonic penguin	$\phi_s^{\text{eff}}(B_s^0 \rightarrow \phi \phi)$ (rad)	0.15	0.10	0.023	0.02
	$\phi_s^{\text{eff}}(B_s^0 \rightarrow K^{*0} \bar{K}^{*0})$ (rad)	0.19	0.13	0.029	< 0.02
	$2\beta^{\text{eff}}(B^0 \rightarrow \phi K_S^0)$ (rad)	0.30	0.20	0.04	0.02
Right-handed currents	$\phi_s^{\text{eff}}(B_s^0 \rightarrow \phi \gamma)$	0.20	0.13	0.030	< 0.01
	$\tau^{\text{eff}}(B_s^0 \rightarrow \phi \gamma)/\tau_{B_s^0}$	5%	3.2%	0.8%	0.2%
Electroweak penguin	$S_3(B^0 \rightarrow K^{*0} \mu^+ \mu^-; 1 < q^2 < 6 \text{ GeV}^2/c^4)$	0.04	0.020	0.007	0.02
	$q_0^2 A_{\text{FB}}(B^0 \rightarrow K^{*0} \mu^+ \mu^-)$	10%	5%	1.9%	$\sim 7\%$
	$A_1(K \mu^+ \mu^-; 1 < q^2 < 6 \text{ GeV}^2/c^4)$	0.09	0.05	0.017	~ 0.02
	$\mathcal{B}(B^+ \rightarrow \pi^+ \mu^+ \mu^-)/\mathcal{B}(B^+ \rightarrow K^+ \mu^+ \mu^-)$	14%	7%	2.4%	$\sim 10\%$
Higgs penguin	$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-)$ (10^{-9})	1.0	0.5	0.19	0.3
	$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-)/\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-)$	220%	110%	40%	$\sim 5\%$
Unitarity triangle angles	$\gamma(B \rightarrow D^{(*)} K^{(*)})$	7°	4°	1.1°	negligible
	$\gamma(B_s^0 \rightarrow D_s^\mp K^\pm)$	17°	11°	2.4°	negligible
	$\beta(B^0 \rightarrow J/\psi K_S^0)$	1.7°	0.8°	0.31°	negligible
Charm	$A_\Gamma(D^0 \rightarrow K^+ K^-)$ (10^{-4})	3.4	2.2	0.5	–
CP violation	ΔA_{CP} (10^{-3})	0.8	0.5	0.12	–