

Bottomia physics at RHIC and LHC

Georg Wolschin

Heidelberg University
Institut für Theoretische Physik
Philosophenweg 16
D-69120 Heidelberg




ICNFP_Kolymbari_2016

Topics

1. Introduction: Υ suppression at RHIC and LHC
2. Model for bottomium suppression
 - 2.1 Complex potential: Screening and damping
 - 2.2 Gluon-induced dissociation
 - 2.3 Hydrodynamic expansion
 - 2.4 Feed-down cascade
3. Comparison with STAR and CMS data
4. Prediction for 5.02 TeV PbPb
5. Conclusion

1. Introduction: Υ Suppression in PbPb @ LHC



Υ suppression as a sensitive probe for the QGP

- No significant effect of regeneration
- $m_b \approx 3m_c \rightarrow$ cleaner theoretical treatment
- More stable than J/ψ


$$E_B(\Upsilon_{1S}) \approx 1.10 \text{ GeV}$$
$$E_B(J/\psi) \approx 0.64 \text{ GeV}$$

CMS Collab., CMS-PAS-HIN-10-006 (2011)

ICNFP_Kolymbari_2016

$\Upsilon(nS)$ states are suppressed in PbPb @ LHC:

CMS



A clear QGP indicator

1. $\Upsilon(1S)$ ground state is suppressed in PbPb:

$$R_{AA}(\Upsilon(1S)) = 0.56 \pm 0.08 \pm 0.07 \text{ in min. bias}$$

2. $\Upsilon(2S, 3S)$ states are > 4 times stronger suppressed in PbPb than $\Upsilon(1S)$

$$R_{AA}(\Upsilon(2S)) = 0.12 \pm 0.04 \text{ (stat.)} \pm 0.02 \text{ (syst.)}$$


$$R_{AA}(\Upsilon(3S)) = 0.03 \pm 0.04 \text{ (stat.)} \pm 0.01 \text{ (syst.)}$$

$$R_{AA} = \frac{N_{PbPb}(Q\bar{Q})}{N_{coll} N_{pp}(Q\bar{Q})}$$


CMS Collab., PRL 109, 222301 (2012)
[Plot from CMS database]

ICNFP_Kolymbari_2016

Sequential suppression of $\Upsilon(nS)$ and J/ψ states



$\Upsilon(nS)$ states are suppressed in AuAu @ RHIC



2. The model: Screening, Gluodissociation and Collisional broadening of the $\Upsilon(nS)$ states

- Debye screening of all states involved: **Static suppression**
- The **imaginary part** of the potential (effect of collisions) contributes to the broadening of the $\Upsilon(nS)$ states: **damping**
- Gluon-induced dissociation: **dynamic suppression**, in particular of the $\Upsilon(1S)$ ground state due to the large thermal gluon density
- Reduced feed-down from the excited Υ/χ_b states to $\Upsilon(1S)$ substantially modifies the populations: **indirect suppression**

F. Vaccaro, F. Nendzig and GW, *Europhys.Lett.* 102, 42001 (2013); J. Hoelck and GW, unpublished
F. Nendzig and GW, *Phys. Rev. C* 87, 024911 (2013); *J. Phys. G* 41, 095003 (2014)
F. Brezinski and GW, *Phys. Lett.B* 70, 534 (2012)

2.1 Screening and damping treated in a nonrelativistic potential model

$$V_{nl}(r, T) = -\frac{\sigma}{m_D(T)} e^{-m_D(T)r} - C_F \alpha_{nl}(T) \left(\frac{e^{-m_D(T)r}}{r} + iT \phi(m_D(T)r) \right)$$

$$\phi(x) = \int_0^\infty \frac{dz}{(1+z^2)^2} \left(1 - \frac{\sin xz}{xz} \right), m_D(T) = T \sqrt{4\pi \alpha_s(2\pi T) \frac{2N_c + N_f}{6}}$$

From the literature

Screened potential: m_D = Debye mass,

$\alpha_{nl}(T)$ the strong coupling constant;


$C_F = (N_c^2 - 1) / (2N_c)$

$\sigma \approx 0.192$ the string tension (Jacobs et al.; Karsch et al.)

Imaginary part: Collisional damping (Laine et al. 2007, Beraudo et al. 2008, Brambilla et al. 2008) for $2\pi T \gg \langle 1/r \rangle$; different form for $2\pi T \ll \langle 1/r \rangle$.


Screened real part of potential, T-dep. imag. part

Constrain $\text{Re}V_s(r)$ by lattice QCD data on the singlet free energy



Screening

Take $\text{Im}V_s(r)$ from pQCD calculations




From: A. Mocsy et al.

Damping

Radial wave function of Y(1S) at temperatures T

Solutions of the Schoedinger equation with complex potential $V(r, T, \alpha_s)$ for the radial wave functions $g_{nl}(r, T)$,

$$[H(r, T, \alpha_s) - E + i\Gamma/2]g(r) = 0$$



From: J. Hoelck and
GW, unpublished

ICNFP_Kolymbari_2016

2.2 Cross section for gluodissociation

Born amplitude for the interaction of gluon clusters according to Bhanot&Peskin in dipole approximation / Operator product expansion, extended to include the screened coulombic + string eigenfunctions as outlined in Brezinski and Wolschin, PLB 70, 534 (2012)

$$\sigma_{diss}^{nS}(E) = \frac{2\pi^2 \alpha_s E}{9} \int_0^\infty dk \delta \left(\frac{k^2}{m_b} + \epsilon_n - E \right) |w^{nS}(k)|^2$$
$$w^{nS}(k) = \int_0^\infty dr r g_{n0}^s(r) g_{k1}^a(r)$$

for the Gluodissociation cross section of the $Y(nS)$ states, and correspondingly for the $\chi_b(nP)$ states.

Gluodissociation cross section




Figure 3. Gluodissociation cross section σ_{diss} (left scale) of the $\Upsilon(1S)$ and $\Upsilon(2S)$ and the thermal gluon distribution (right scale) plotted for temperature $T = 170$ (solid curves) and 250 MeV (dotted curves) as functions of the gluon energy E_g .

F. Nendzig and GW, J. Phys. G41, 095003 (2014)

ICNFP_Kolymbari_2016

Thermal gluodissociation cross section

Average the gluodissociation cross section over the Bose-Einstein distribution of the thermal gluons in the QGP to obtain the dissociation width at temperature T for each of the six bottomia states involved


$$\Gamma_{\text{diss, } nl}(T) \equiv \frac{g_d}{2\pi^2} \int_0^\infty \frac{dE_g E_g^2 \sigma_{\text{diss, } nl}(E_g)}{e^{E_g/T} - 1}$$

(g_d = 16)

With rising temperature, the peak of the gluon distribution moves to larger gluon energies E_g, whereas the dissociation cross sections move to smaller E_g, giving rise to a maximum in the gluodissociation width for fixed coupling α_s.
(Larger cross sections at higher temperatures due to **running coupling** counteract.)

Damping and gluodissociation widths for six bottomia states

$$\Gamma_{\text{tot}}(T) = \Gamma_{\text{damp}}(T) + \Gamma_{\text{diss}}(T)$$




F. Nendzig and GW, J. Phys. G41, 095003 (2014) ; arXiv:1406.5103

ICNFP_Kolymbari_2016

2.3 Hydrodynamic expansion (ideal)

Temperature profile for central collisions at different times τ

Use total decay widths $\Gamma_{\text{tot}}(b, x, y)$ of the bottomia states for each impact parameter b and time step t in the transverse (x^1, x^2) plane



Dynamical fireball evolution

Dependence of the local temperature T on impact parameter b , time t , and transverse coordinates x, y evaluated in ideal hydrodynamic calculation with transverse expansion

$$T(b, \tau_{init}, x^1, x^2) = T_0 \left(\frac{N_{mix}(b, x^1, x^2)}{N_{mix}(0, 0, 0)} \right)^{1/3}$$

$$N_{mix} = \frac{1-f}{2} N_{part} + f N_{coll}, \quad f = 0.145$$


The number of produced $b\bar{b}$ -pairs is proportional to the number of binary collision, and the nuclear overlap

$$N_{b\bar{b}}(b, x, y) \propto N_{coll}(b, x, y) \propto T_{AA}(b, x, y)$$

QGP suppression factor (without feed-down and CNM effects):

$$R_{AA}^{QGP} = \frac{\int d^2b \int dx dy T_{AA}(b, x, y) e^{-\int_{t_F}^{\infty} dt \Gamma_{tot}(b, t, x, y)}}{\int d^2b \int dx dy T_{AA}(b, x, y)}$$

Model ingredients




$$\alpha_s(Q) = \frac{\alpha(\mu)}{1 + \alpha(\mu)b_0 \ln \frac{Q}{\mu}}, \quad b_0 = \frac{11N_c - 2N_f}{6\pi}$$

F. Nendzig and GW, J. Phys. G41, 095003 (2014)

$\alpha_{nl}(T)$ depends on the solution $g_{nl}(r, T)$ of the Schrödinger eq.: Iterative solution

2.4 Feed-down cascade including χ_{nP} states


Relative initial populations in pp computed using an inverted cascade from the final populations measured by CMS and CDF(χ_b)
 $[N_{\text{final}}(1S)=1]$




3. Theoretical vs. exp. (STAR, CMS) suppression

- Screening (potential model)
- Gluodissociation (OPE with string tension included)
- Collisional damping (imaginary part of potential)
- Reduced feed-down from excited states

t_F : γ formation time
 T_0 @ t_F : initial central temperature



Theoretical vs. exp. (CMS) suppression factors



$t_F = 0.4 \text{ fm/c}$: Y formation time
 $T_0 = 550 \text{ MeV}$: central temp.
at $b = 0$ and $t = t_F$


2.76/ 5.02 TeV PbPb

Room for additional suppression mechanisms for the excited states:
Hadronic dissociation, mostly by pions, is one possibility. Thermal pions are insufficient; direct pions may contribute, and magnetic dissociation.

J. Hoelck, F. Nendzig and GW,
arXiv:1602.00019 (2016)

Transverse momentum dependence of $\Upsilon(1S)$ suppression in PbPb at 2.76/5.02 TeV

Γ -averaging, min. bias 0-100%



5. Conclusion Υ at RHIC and LHC

- ❖ The suppression of the $\Upsilon(1S)$ ground state in UU collisions at RHIC and PbPb at LHC energies through gluodissociation, damping, screening, and reduced feed-down has been calculated for min. bias, and as function of centrality, and is found to be in good agreement with the CMS result. Screening is not decisive for the $1S$ state except for central collisions. The flat p_T dependence is understood based on the relativistic Doppler effect.
- ❖ The enhanced suppression of the $\Upsilon(2S, 3S)$ relative to the $1S$ state in PbPb as compared to pp collisions at LHC energies (CMS) leaves room for additional suppression mechanisms, in particular for peripheral collisions where discrepancies to the CMS data persist. Hadronic and/or magnetic dissociation of the excited states may be relevant.

❖ Thank you for your attention,
and for organizing ICNFP2016 !

