Forward-backward multiplicity correlations in pp collisions at LHC within the QGSM

UiO : Universitetet i Oslo

Alexander von Humboldt Stiftung/Foundation

L. Bravina (UiO)

in collaboration with

J. Bleibel and E. Zabrodin

MEPhI

НИИЯФ МГУ

FIAS Frankfurt Institute for Advanced Studies

EBERHARD KARLS UNIVERSITÄT TÜBINGEN

5th International Conference on New Frontiers in Physics (ICNFP´2016) July 9, 2016, Kolymbari, Crete, Greece

Outline

- Forward-backward multiplicity correlations in hadronic interactions
- Quark-gluon string model (QGSM)
- FB correlations at intermediate energies
- Origin of FB correlations at LHC energies; comparison with data
- Conclusions

I. Forward-backward multiplicity correlations in hadronic interactions

Observation of strong correlations (ISR)

S. Uhlig et al., NPB 132 (1978) 15

central region



fragmentation region



Main results

see also talk by V.Vechernin

$$\langle n_F(n_B)\rangle = a + b_{corr} n_B$$

$$b_{corr} = \frac{\langle n_F n_B\rangle - \langle n_F\rangle^2}{\langle n_F^2\rangle - \langle n_F\rangle^2}$$

- The FB multiplicity correlations are positive
- Correlations are stronger for anti-pp collisions compared to pp ones (at energies below 100 GeV)
- No sizeable FB correlations were found in e^+e^- annihilation up to 93 GeV
- Correlations are due to central region $|x_F| < 0.1$

(some) references

theory/models

A.~Capella, A.~Krzywicki, PRD 18 (1978) 4120.
A.~Capella, J.Tran Thanh Van, ZPhC 18 (1983) 85.
J.~Dias~de~Deus, PLB 100 (1981) 177.
T.T.~Chou, C.N.~Yang, PLB 135 (1984) 175.
B.~Carazza, Nuovo Cim. 99 (1988) 731.
S.~Barshay, PLB 199 (1987) 121.
L.K.~Chen, D.~Kiang, C.K.~Chew, PLB 408 (1997) 422.
N.~Armesto, M.A.~Braun, E.G.~Ferreiro, C.~Pajares, NPA 661 (1999) 325c.
M.A.~Braun, C.~Pajares, V.V.~Vechernin, PLB 493 (2000) 54.
V.V.~Vechernin, NPA 939 (2015) 21.
A.~Capella et al.,Phys. Rep. 236 (1994) 225.
N.S.~Amelin et al., PRL 73 (1994) 2813.
Y.-L.~Yan et al., NPA 834 (2010) 320c.
K.~Wraight, P.~Skands, EPJC 71 (2011) 1628.
V.~Kovalenko, V.~Vechernin, arXiv:1410.3884 [hep-ph].

Experiment

K.~Alpgaard et al., UA5 Collab., Phys. Lett. B 123 (1983) 361.
G.G.~Alner et al., UA5 Collab., Phys. Rep. 154 (1987) 247.
R.E.~Ansorge et al., UA5 Collab., Z. Phys. C 37 (1988) 191.
S.~Uhlig, et al., Nucl. Phys. B 132 (1978) 15.
L.V.~Bravina et al., Mirabelle Collab., Sov. J. Nucl. Phys. 50 (1989) 245.
V.V.~Aivazyan et al., NA22 Collab., Z. Phys. C 42 (1989) 533.
T.~Alexopoulos et al., E735 Collab., Phys. Lett. B 353 (1995) 155.
J.~Adam et al., ALICE Collab., J. High Energy Phys. JHEP05 (2015) 097
W.~Braunschweig et al., TASSO Collab., Z. Phys. C 45 (1989) 193.
R.~Akers, et al., OPAL Collaboration, Phys. Lett. B 320(1994) 417.

II. Quark-Gluon String Model (QGSM)

Diagrams at intermediate energies



(a) -- planar
(b) -- cylindrical
(c) – undeveloped cylinder
(d) – quark rearrangement
(e) – single diffraction of small mass
(f) – single diffraction of large mass
(g)-(i) – annihilation diagrams
+ double diffraction diagrams similar to (e), (f)

Because of the different sets of diagrams for pp and anti-pp collisions (particularly, annihilation) there should be a difference in FB multiplicity correlations for these two reactions.

Diagrams at ultrarelativistic energies



(a) -- multi-cylinder
(b) – semihard
(+ soft) Pomeron
(c) – single diffraction of large mass
(d) – single diffraction of small mass
(e) – double diffraction of small mass
(f) – double diffraction of large mass

At ultrarelativistic energies only few diagrams are left, whereas cross sections of annihilation and pre-asymptotic processes quickly drop
=> No difference between pp and anti-pp collisions

III. FB multiplicity correlations at intermediate energies

(anti)pp @ 32 GeV/c (data)

L.B. et al., Sov. J. Nucl. Phys. 50 (1989) 245

TABLE I. Parameters of the approximation by the function $\langle n_F \rangle = a + bn_B$ for $\bar{p}p$ and pp interactions at 32 GeV/c.

Reaction	$\bar{p}p$		
	a	b	χ^2/deg. fr.
Inelastic interactions	2,03±0,02	0,192±0,007	22/(10−2−1)
Nondiffraction interactions	2,18±0,04	0,177±0,010	24/(10−2−1)
Particles with $\lvert x \rvert < 0.1$ in inelastic interactions	1,09±0,01	0,237±0,006	23/(12−2−1)
Particles with $\lvert x \rvert > 0.1$ in inelastic interactions	1,144±0,008	0,013±0,006	0,29/(6−2−1)

Reaction	pp		
	a	b	χ^2/deg. fr.
Inelastic interactions	2,07±0,04	0,098±0,016	92/(11−2−1)
Nondiffraction interactions	2,25±0,05	0,07±0,02	36/(11−2−1)
Particles with $\lvert x \rvert < 0.1$ in inelastic interactions	1,11±0,04	0,175±0,015	19/(9−2−1)
Particles with $\lvert x \rvert > 0.1$ in inelastic interactions	1,262±0,009	−0,075±0,008	1,5/(6−2−1)

No long-range correlations between the fragmentation regions

(anti)pp @ 32 GeV/c vs. QGSM

antiproton-proton **proton-proton**



L.B. et al., Sov. J. Nucl. Phys. 50 (1989) 245

Closed points: inelastic collisions
Open points: non-diffractive collisions
Solid curves: QGSM
Dashed curve: Lund

(anti)pp @ 32 GeV/c vs. QGSM



Open points: inelastic collisions
Solid curves: QGSM
Dashed curves: Lund

(a) |x| > 0.1

(b) |x| < 0.1

No long-range correlations

Positive FB correlations come from the region |x| < 0.1 corresponding to |y| < 1, i.e. midrapidity region

L.B. et al., Sov. J. Nucl. Phys. 50 (1989) 245

FB correlations within the subprocesses



(a) – cylindrical diagrams

(b) – undeveloped cylinder

(c) – diffraction diagrams

(d) – annihilation diagrams

(e) – planar diagrams

No correlations in individual processes, however, addition of all these processes with different $\langle n_F \rangle$ leads to appearance of positive correlations $\langle n_F \rangle (n_B)$

L.B. et al., Sov. J. Nucl. Phys. 50 (1989) 245

IV. Origin of FB multiplicity correlations in QGSM at LHC energies

pp @ 20 GeV to 14 TeV in QGSM



(1) The slope is almost linear; (2) no difference between pp and anti-pp at high energies; (3) slope parameter b increases with rising energy; (4) saturation of the increase at high multiplicities. Why?

Rel. weight of soft and hard Pomerons



The ratio hard/soft increases from 0.29 : 1.14 (900 GeV)
to 1.10 : 1.54 (13 TeV)

Contribution of soft and hard Pomerons

Soft Pomerons



Hard Pomerons



Change of the slope with energy



The slope increases. Also, the distributions are more horizontal at very small and very high multiplicities. Why?

Contributions of n-Pomeron processes



If we select processes going via 1,2,…N Pomerons, then the distributions are remarkably flat (no FB correlations). However, average multiplicities are different.
This leads to positive correlations when one adds all processes together. The same is true for the diffraction, but the slope is not so steep, therefore

$b_{NSD} > b_{inel}$

more details ...



Comparison with LHC data



… fine tuning might be needed



Agreement is good, however

CONCLUSIONS

- *In pp and $\bar{p}p$ collisions positive forward-backward correlations of the multiplicity of particles are observed in QGSM in a broad energy range from 8 GeV to 14 TeV*
- *At energies below 100 GeV the correlations are higher for $\bar{p}p$ interactions*
- *The observed dependences are due to hadrons coming from the central area $|x_F| < 0.1$*
- *No correlations are observed for events corresponding to the same mechanism of particle production*
- *At lowest and highest multiplicities the distributions are more flat*
- ***The analysis shows that these correlations are not "true" correlations due to the mechanism of particle production but arise as a consequence of the addition of events corresponding to different production mechanisms with different average multiplicities of secondary hadrons. This is in line with the observations of other models.***

Back-up Slides