

# High-Precision Higgs Masses in the Complex MSSM

Sebastian Paßehr<sup>†</sup>

in collaboration with

Wolfgang Hollik<sup>‡</sup>, arXiv:1401.8275 [hep-ph], arXiv:1409.1687 [hep-ph],

Thomas Hahn<sup>‡</sup>, arXiv:1508.00562 [hep-ph],

Sophia Borowka<sup>§</sup>, Georg Weiglein<sup>†</sup>


<sup>†</sup>DESY, Hamburg, <sup>‡</sup>Max Planck Institute for Physics, Munich, <sup>§</sup>Institute for Physics, Universität Zürich

SUSY 2016,  
Melbourne, Australia,  
5th of July 2016



# Experimental discovery and mass measurement

Higgs-like particle discovered, [ATLAS, arXiv:1207.7214 [hep-ex]],  
 [CMS, arXiv:1207.7235 [hep-ex]],  
 e. g. signal in  $H \rightarrow \gamma\gamma$ , [ATLAS, arXiv:1406.3827 [hep-ex]],



- very good agreement with SM Higgs boson,
- but: SM has many deficiencies,
- test models beyond the Standard Model,  
 e. g. Supersymmetry, here: MSSM,
- experimental value:  
 $125.09 \pm 0.21(\text{stat}) \pm 0.11(\text{syst}) \text{ GeV}$   
 [ATLAS, CMS, arXiv:1503.07589].

two complex  $SU(2)$ -Higgs doublets (eight degrees of freedom):

$$\mathcal{H}_1 = \begin{pmatrix} v_1 + \frac{1}{\sqrt{2}} (\phi_1^0 - i \chi_1^0) \\ -\phi_1^- \end{pmatrix}, \quad \mathcal{H}_2 = e^{i\zeta} \begin{pmatrix} \phi_2^+ \\ v_2 + \frac{1}{\sqrt{2}} (\phi_2^0 + i \chi_2^0) \end{pmatrix},$$

positive real vacuum expectation values  $v_1, v_2$ ,  $\tan \beta \equiv v_2/v_1$ ,  
relative phase  $\zeta$ ,

with superpotential

$$\mathcal{W}_{\text{MSSM}} = \mu H_1 \cdot H_2 - h_{u,ij} Q_i \cdot H_2 U_j^C - h_{e,ij} H_1 \cdot L_i E_j^C - h_{d,ij} H_1 \cdot Q_i D_j^C.$$

tree-level mass eigenstates:

$CP$  even  $h, H$ ;  $CP$  odd  $A$ ; charged  $H^\pm$

MSSM  $CP$  conserving at tree level

Higgs masses at  $k$  loop order given by poles of propagator matrix

$$\Delta_{hHAG}^{(k)}(p^2) = i \left[ p^2 \mathbf{1} - \mathbf{M}_{hHAG}^{(k)}(p^2) \right]^{-1},$$

matrix of renormalized two-point vertex functions:

$$\hat{\Gamma}_{hHAG}^{(k)}(p^2) = - \left[ \Delta_{hHAG}^{(k)}(p^2) \right]^{-1},$$

masses determined by

$$\det \left[ \hat{\Gamma}_{hHAG}^{(k)}(p^2) \right]_{p^2 = x_i^2} = 0, \quad m_{h_i}^2 = \Re[x_i^2], \quad i \in \{1, 2, 3\},$$

(fourth solution belongs to Goldstone boson, equal to zero).

- lowest order:  $\mathbf{M}_{hHAG}^{(k)}(p^2) \Big|_{k=0} = \mathbf{M}_{hHAG}^{(0)}$ , diagonal,
- higher order:  $\mathbf{M}_{hHAG}^{(k)}(p^2) \Big|_{k \geq 1} = \mathbf{M}_{hHAG}^{(0)} - \sum_{j=1}^k \hat{\Sigma}_{hHAG}^{(j)}(p^2)$ ,


shift by renormalized self-energies

$$\hat{\Sigma}_{hHAG}^{(j)}(p^2) = \begin{pmatrix} \hat{\Sigma}_h^{(j)}(p^2) & \hat{\Sigma}_{hH}^{(j)}(p^2) & \hat{\Sigma}_{hA}^{(j)}(p^2) & \hat{\Sigma}_{hG}^{(j)}(p^2) \\ \hat{\Sigma}_{hH}^{(j)}(p^2) & \hat{\Sigma}_H^{(j)}(p^2) & \hat{\Sigma}_{HA}^{(j)}(p^2) & \hat{\Sigma}_{HG}^{(j)}(p^2) \\ \hat{\Sigma}_{hA}^{(j)}(p^2) & \hat{\Sigma}_{HA}^{(j)}(p^2) & \hat{\Sigma}_A^{(j)}(p^2) & \hat{\Sigma}_{AG}^{(j)}(p^2) \\ \hat{\Sigma}_{hG}^{(j)}(p^2) & \hat{\Sigma}_{HG}^{(j)}(p^2) & \hat{\Sigma}_{AG}^{(j)}(p^2) & \hat{\Sigma}_G^{(j)}(p^2) \end{pmatrix},$$


$\hat{\Sigma}_{hA}^{(j)}(p^2)$ ,  $\hat{\Sigma}_{HA}^{(j)}(p^2) \neq 0 \Rightarrow$  complex parameters (e.g.  $\mu$ ,  $A_t$ ,  $A_b$ ,  $M_3$ , ...).

# Mixed particles at higher orders


- lowest order mass eigenstates:



- higher orders: off-diagonal entries in  $\hat{\Sigma}_{hHAG}^{(j)}(p^2)$  induce mixing,




# lightest Higgs mass with FeynHiggs



- upper bound at tree level:  $M_Z$
- huge one-loop corrections: +50%
- two-loop in complex MSSM:  
 leading  $\mathcal{O}(\alpha_t \alpha_s)$   
[Heinemeyer, Hollik, Rzehak, Weiglein, arXiv:0705.0746]  
 leading  $\mathcal{O}(\alpha_t^2)$   
[Hollik, SP, arXiv:1401.8275, 1409.1687]
- orange band:  $\pm 1\text{GeV}$

$m_h^{\text{mod}}$  scenario:  $M_{H^\pm} = 0.6\text{TeV}$ ,  $\mu = 0.2\text{TeV}$ ,  
 $M_{\text{SUSY}} = 1.0\text{TeV}$ ,  $A_f = 1.5\text{TeV}$ ,  $M_{\tilde{g}} = 1.5\text{TeV}$

# focus at two-loop order




- huge effect by  $\mathcal{O}(\alpha_t \alpha_s)$ :  
 $\approx -17\text{GeV}$
- also big effect by  $\mathcal{O}(\alpha_t^2)$ :  
 $\approx +5\text{GeV}$

$m_h^{\text{mod}}$  scenario:  $M_{H^\pm} = 0.6\text{TeV}$ ,  $\mu = 0.2\text{TeV}$ ,  
 $M_{\text{SUSY}} = 1.0\text{TeV}$ ,  $A_f = 1.5\text{TeV}$ ,  $M_{\tilde{g}} = 1.5\text{TeV}$


- extension for leading  $\mathcal{O}(\alpha_t \alpha_s)$ :  
full lowest order QCD, i. e.  $\mathcal{O}(\alpha_{\text{any}} \alpha_s)$ ,  $\alpha_{\text{any}} = \alpha, \alpha_t, \alpha_b, \dots$   
no approximations applied,  
all subleading terms included,  
momentum dependence taken into account
- extension for leading  $\mathcal{O}(\alpha_t^2)$ :  
Yukawa<sup>4</sup> terms, i. e.  $\mathcal{O}(\alpha_t^2 + \alpha_t \alpha_b + \alpha_b^2)$   
gauge-less approximation applied,  
subleading Yukawa terms included,  
no momentum dependence

results generated with help of FeynArts, FormCalc, TwoCalc  
following previously developed scripts [Hahn, SP, arXiv:1508.00562]

# $\mathcal{O}(\alpha_{\text{any}}\alpha_s)$ , Feynman diagrams



# $\mathcal{O}(\alpha_t^2 + \alpha_t \alpha_b + \alpha_b^2)$ , Feynman diagrams



- $\mathcal{O}(\alpha_{\text{any}}\alpha_s)$

subrenormalization, only  $\mathcal{O}(\alpha_s)$ :

$\delta m_{\tilde{t}_{1,2}}, \delta m_{\tilde{t}_{12}}, \delta m_t, \delta m_{\tilde{b}_2}$  on-shell

$\delta m_b, \delta A_b$   $\overline{\text{DR}}$

genuine two loop:

$\delta m_{H^\pm}, \delta T_{h,H,A}, \delta m_W, \delta m_Z$  on-shell

$\delta Z_{\mathcal{H}_1}, \delta Z_{\mathcal{H}_2}, \delta t_\beta$   $\overline{\text{DR}}$

- $\mathcal{O}(\alpha_t^2 + \alpha_t\alpha_b + \alpha_b^2)$

subrenormalization,  $\mathcal{O}(\alpha_t + \alpha_b)$ :

$\delta m_{\tilde{t}_{1,2}}, \delta m_{\tilde{t}_{12}}, \delta m_t, \delta m_{\tilde{b}_2}, \delta \mu$  on-shell

$\delta m_b, \delta A_b$   $\overline{\text{DR}}$


$\frac{\delta M_Z}{M_Z}, \frac{\delta M_W}{M_W}$  on-shell

$\delta m_{H^\pm}, \delta T_{h,H,A}$  on-shell

$\delta Z_{\mathcal{H}_1}, \delta Z_{\mathcal{H}_2}, \delta t_\beta$   $\overline{\text{DR}}$

genuine two loop:

$\delta m_{H^\pm}, \delta T_{h,H,A}$  on-shell




- opposite sign of new contributions
- momentum-dependent integrals evaluated with help of SecDec

[Borowka, Carter, Heinrich, arXiv:1011.5493, 1204.4152, 1303.1157]

$m_h^{\text{mod}}$  scenario:  $M_{H^\pm} = 0.6\text{TeV}$ ,  $\mu = 0.2\text{TeV}$ ,  
 $M_{\text{SUSY}} = 1.0\text{TeV}$ ,  $A_f = 1.5\text{TeV}$ ,  $M_{\tilde{g}} = 1.5\text{TeV}$

# phase dependence: $\phi_{M_3}$

— two loop (old,  $\phi_{M_3} = 0$ )      — two loop (new)  
- - - - two loop (old,  $\phi_{M_3} \neq 0$ )




- in general: large shift by phase
- new terms of  $\mathcal{O}(\alpha_{\text{any}}\alpha_s)$  depend only slightly on  $\phi_{M_3}$ : shift by  $\approx -20\text{MeV}$

$m_h^{\text{mod}}$  scenario:  $M_{H^\pm} = 0.6\text{TeV}$ ,  $\mu = 0.2\text{TeV}$ ,  $t_\beta = 15$ ,  
 $M_{\text{SUSY}} = 1.0\text{TeV}$ ,  $A_f = 1.5\text{TeV}$ ,  $M_{\tilde{g}} = 1.5\text{TeV}$

# phase dependence: $\phi_{A_t}$

- two loop (old,  $\phi_{A_t} = 0$ )
- two loop (new)
- ⋯ two loop (old,  $\phi_{A_t} \neq 0$ )
- one loop



- in general: large shift by phase
- new terms of  $\mathcal{O}(\alpha_t^2 + \alpha_t \alpha_b + \alpha_b^2)$  depend only slightly on  $\phi_{A_t}$ : shift by  $\approx -30\text{MeV}$

$m_h^{\text{mod}}$  scenario:  $M_{H^\pm} = 0.6\text{TeV}$ ,  $\mu = 0.2\text{TeV}$ ,  $t_\beta = 15$ ,  
 $M_{\text{SUSY}} = 1.0\text{TeV}$ ,  $A_f = 1.5\text{TeV}$ ,  $M_{\tilde{g}} = 1.5\text{TeV}$

- analyze separately the contributions to  $\mathcal{O}(\alpha_{\text{any}}\alpha_s)$  by momentum-dependent terms, gauge terms,  $m_b$  terms
- compare  $\mathcal{O}(\alpha_{\text{any}}\alpha_s)$  in limit  $m_b \rightarrow 0$  and real parameters with previous result
- compare  $\mathcal{O}(\alpha_q\alpha_{q'})$  in limit of real parameters with previous result
- implement in FeynHiggs
- implement running  $m_t$  consistently
- make results available in on-shell scheme and  $\overline{\text{DR}}$  scheme
- investigate effect on charged Higgs mass in real MSSM