# Progress in GEM and PreCompound models

J. M. Quesada Geant4 Collaboration Workshop Catania, 19.10.2009


### IAEA benchmark of spallation data

- The benchmark includes nucleon-induced reactions on nuclei from carbon to uranium
- Energy range: 20 MeV to 3 GeV
- Exp. data from thin target experiments for model testing
- After intensive model improvement, Geant4 decided to participate
- This benchmark has triggered a series of critical model improvements in pre-compound & deexcitation models in Geant4

# Bug fixing in PreCompound Model

- Fixed bug in probability distribution of cluster (<sup>2</sup>H, <sup>3</sup>H, <sup>3</sup>He, <sup>4</sup>He) emission
  - level density parameter (a) of the emitted cluster (a physical nonsense!) instead of the one of the residual nucleus)
  - a~A/10
  - It enters into the emission probability through (proportional to) g<sup>Aj</sup>, where
    - $g=(6/\pi^2)$  a
    - Aj is the mass number of the cluster


#### Light cluster emission: improvement



# **Bug fixing GEM**

- Fixed bug in probability distribution of particle emission :
  - hbar\_Plack instead of hbarc at the denominator
    - A factor 9 10<sup>4</sup> (!!) was spureously introduced
- Several fixes in the implementation of Furihata's formulation
- Fix in kinetic energy samplig which caused an infinite loop in some cases

#### Isotope production at 1000 MeV in inverse kinematics



# **Bug fixing in Fission Model**


- Fixed bug in total probability of fission :
  - Missing parenthesis at the denominator:
    a<sub>fis</sub> must enter mutiplying instead of dividing
- Tuning of the width of the gaussian distribution for simetric fission

#### Isotope production at 1000 MeV in inverse kinematics


**BEFORE** 

9.2p01

Without GEM (buggy at that time)



#### Isotope production at 1000 MeV in inverse kinematics



## Fixes in Femi BreakUp


- Excitation energies of available fragments in Fermi BreakUp
  - where in keV instead of MeV
- Bug in implementation of Gamma function in the statistical weight of the channels
  - Even partitions were spureously biased by a factor 0.5

# Geant4 simulations of thermal neutrons irradiation on pad detectors with converter layers

Part 2 (with corrections: Quesada's files \*)

- Detector Efficiency
- Spectrums of energy deposited on planar detector convered by converter layer (boron clusters)

#### Detector Efficiency (different clusters)




$$^{10}B + n \rightarrow \alpha (1.47 MeV) + ^{7}Li(0.84 MeV) + \gamma (0.48 MeV)$$
 93.7%

$$^{10}B + n \rightarrow \alpha (1.78 MeV) + ^{7}Li(1.01 MeV)$$
 6.3%


#### Quesada's files


("G4FermiFragmentsPool.cc" and "G4ExcitationHandler.cc")

#### 106 thermal neutrons




range 3-5 µm detection efficiency maximized






5  $\mu$ m converter layer (  $[B_{12}H_{12}]^{2-}$  )


300 µm sensitive detector (Si)



# Improvement in Statistical Multifragmentation Model

- Statistical Multifragmentation Model should act only once in de-excitation.
  - G4ExcitationHandler has been corrected and code intercomparison by Igor Pshenichnov finally is OK.






### The handler

- It handles the different models which are included in the de-excitation process
- A reshaping of it has been accomplished (M.A. Cortés):
  - More efficient & clearer algorithms
  - A bug was fixed:
    - A (likely) still excited fragment was skipped at each iteration of the de-excitation loop.
- Now we have full control of what is going on inside it

## Flux diagram of G4ExcitationHandler



### Conclusions

- Many improvements have been made during the last year
- But...
  - There are still pending issues

To be continued ....

Thanks for your attention!

## **Backup slides**

Results of Geant4 in IAEA bechmark as they were in May 2009 (previous to GEM fixing and fission parameters tuning)

# Results obtained with nuclear models of Geant4 in IAEA Benchmark of Spallation

J. M. Quesada on behalf of the Geant4 Hadronic Group IAEA, Vienna, 05.05.2009

## General Introduction


#### What is Geant4?

- Geant4 is the C++, object-oriented successor to GEANT3
- Designed primarily with high energy physics in mind
  - but now used in medical and space applications as well
- It is a toolkit:
  - large degree of functionality and flexibility are provided
  - many different codes provided, including alternates covering the same regions of applicability
  - choice of which to use is up to user, but guidance provided by Geant4 developers
- All major physics processes covered:
  - electromagnetic, hadronic, decay, photo- and electro-nuclear

# Geant4 Hadronic Processes and Models

- Hadronic processes include
  - Elastic
  - Inelastic
  - Capture at rest
  - Neutron capture
  - Neutron-induced fission
  - Lepton-nuclear
  - Gamma-nuclear
- Each of the above processes is implemented by one or more:
  - models (which contain the physics algorithm)
  - cross sections (which determine mean free path, etc.)

### Geant4 hadronic models



#### Geant4 cascade models

- The large energy region considered in this benchmarking includes different interaction regimes.
- In order to predict the production cross sections, different reaction mechanisms must be considered
  - Cascade
  - Pre-equilibrium
  - Equilibrium de-excitation
- Cascade models all have nuclear de-excitation models embedded in them

## Why several cascade models?

#### Binary:

- a time-dependent model which depends as little as possible on parameterization and therefore can be expected to be more predictive
- is an *in house* development, including its own precompound and evaporation models.

#### Bertini:

- came from the INUCL code which was intended as an allinclusive model.
- It came with its own precompound and evaporation models. Neither of these are very different in origin from those in Binary, but the implementations are different.

# Geant4 ongoing developments not included in this benchmark

- CHIPS, Chiral invariant phase space, :
  - Quark-level event generator for the fragmentation of hadronic systems into hadrons.
  - Includes nonrelativistic phase space of nucleons to explain evaporation

#### INCL/ABLA

- C++ translation of INCL intranuclear cascade code
- C++ translation of ABLA evaporation/fission code

## Geant4 Bertini Cascade: Origin

- A re-engineered version of the INUCL code of N. Stepanov (ITEP)
- Employs many of the standard INC methods developed by Bertini (1968)
  - using free particle-particle collisions within cascade
  - step-like nuclear density
- Similar methods used in many different intra-nuclear transport codes

### Applicability of the Bertini Cascade

- inelastic scattering of p, n,  $\pi$  , K,  $\Lambda$  ,  $\Sigma$  ,  $\Xi$
- incident energies: 0 < E < 10 GeV</p>
  - upper limit determined by lack of partial final state cross sections and the end of the cascade validity region
  - lower limit due to inclusion of internal nuclear de-excitation models
- in principle, can be extended to:
  - anti-baryons
  - ion-ion collisions

# Origin and Applicability of the Binary Cascade

- H.P. Wellisch and G. Folger (CERN)
- Henning Weber (Frankfurt group)
- Based in part on Amelin's kinetic model
- Incident p, n

- light ions
  - ■0 < E < ~3 GeV/A
- $\pi$ 
  - ■0 < E < ~1.5 GeV

### Binary Cascade Model

- Hybrid between classical cascade and full QMD model
- Detailed model of Nucleus
  - nucleons placed in space according to nuclear density
  - nucleon momentum according to Fermi gas model
- Nucleon momentum is taken into account when evaluating cross sections, i.e. collision probability
- Collective effect of nucleus on participant nucleons described by optical potential
  - numerically integrated equation of motion

#### The Nuclear model

- Nucleon momenta are sampled assuming Fermi gas model
- Nuclear density
  - harmonic oscillator for A < 17
  - Woods-Saxon for others
- Sampling is done in a correlated manner:
  - local phase-space densities are constrained by Pauli principle
  - sum of all nucleon momenta must equal zero

# Inverse reaction cross sections (preequilibrium & equilibrium)

Inverse reaction cross sections play a mayor role in the calculation of (competing) emission probabilities.

Theory driven (old) parameterization (Dostrovski et al, 1959)

#### New parameterization:

• More realistic parameterization of reaction cross sections (Kalbach), calculated with global optical model potentials, in turn fitted to reproduce available experimental data (angular distributions, elastic scattering, total cross sections, etc..).

## Remarks


- No *ad hoc* tuning of level density parameter ratio  $a_{fis}/a_{evap}$ . (preliminary trials show that it is critical, as reported in previous works).
- No soft transition from pre-equilibrium (i.e. increment of equilibium at the expenses of pre-equilibrium).
- Very important: parameters tuned in a "model suite" shuldn't be assumed to work in a different environment, i.e. with different coupled models.

Quite likely, ad hoc tuning of parameters will be necessary in order to reproduce fission data.


#### **RESULTS**

(Geant4 official release 9.2 patch p01)


## Neutron production at 63 MeV




## Neutron production at 1200 MeV




## Proton production at 62 MeV




## Proton production at 175 MeV




## Proton production at 1200 MeV




## Pion production at 730 MeV




## Pion production at 2205 MeV




## Deuteron production at 63 MeV




## Deuteron production at 1200 MeV




# Tritium production at 63 MeV




# Tritium production at 1200 MeV




# <sup>3</sup>He production at 175 MeV




# <sup>3</sup>He production at 1200 MeV




# Alpha production at 63 MeV




# Alpha production at 175 MeV



# Alpha production at 1200 MeV



## Fission at 1000 MeV



# Conclusions (1)

- Bertini agrees better with data for:
  - protons (high energy)
  - pions (high and low energy)
  - fission
- Binary agrees better with data for:
  - low and medium energy protons
  - almost all light ion production, although agreement is not good in either case
- Many cases where neither model is better overall
  - one model may be better for forward angles, the other for backward angles

# Conclusions (2)

- The fact that we cannot say that one model is clearly better than the other emphasizes the need for alternate models in same energy range
- This benchmark study demonstrated areas where improvement is needed. As a result:
  - recently made improvements to precompound
  - plan to add coalescence models for cascade stage
  - improvements to fission are possible

Thanks for your attention

# Backup slides

#### **Geant4 Collaboration**












**Jefferson** 



















Collaborators also from nonmember institutions, including Budker Inst. of Physics IHEP Protvino MEPHI Moscow Pittsburg University University of Sevilla CIEMAT


## **Brief History**

- Dec'94 : Project started
- Dec'98: First public release
- Geant4 was used by BaBar experiment at SLAC since 2000
- Geant4 is used for Monte Carlo simulation of particle transport for ATLAS, CMS, LHCb since 2004
- Hadronic physics packages are an important part of Geant4 for LHC
  - Signal acceptance
  - Background estimation





## Cascade Modeling Concept



- The Bertini model is a classical cascade:
  - it is a solution to the Boltzmann equation on average
    Bertini Cascade Model
    no scattering matrix calculated
- Core code:
  - elementary particle collider: uses free cross sections
    - Up to and including 6-body final state partial cross sections for pi+p, pi-p, pp, pn from the CERN compilations (V. Flaminio et al., 1983 and 1984). K+, K- partial cross sections also from Flaminio (1983).
    - pi+n, pi-n, nn cross sections are obtained through isospin arguments
  - Generated secondaries:
    - pions, nucleons, kaons, hyperons.
    - No resonances
    - Deuterons, tritons, 3He, alphas (from avaparation phace only)

#### Bertini Cascade Modeling Sequence (1)

- Nuclear entry point sampled over projected area of nucleus
- Incident particle is transported in density dependent nuclear medium
  - mean free path from total particle-particle cross sections
  - Nucleus modeled as 3 concentric, constant-density shells plus reflection/transmission shell boundaries.
  - nucleons have Fermi gas momentum distribution
  - Pauli exclusion invoked
- Projectile interacts with a single nucleon
  - hadron-nucleon interactions based on free cross sections and angular distributions
  - pions can be absorbed on quasi-deuterons

#### Bertini Cascade Modeling Sequence (2)

- Each secondary from initial interaction is propagated in nuclear potential until it interacts or leaves nucleus
  - can have reflection from density shell boundaries
  - Coulomb barrier added recently
- As cascade collisions occur, exciton states are built up, leading to equilibrated nucleus
  - selection rules for p-h state formation:  $\Delta$  p = 0, +/1,  $\Delta$  h = 0, +/-1,  $\Delta$  n = 0, +/-2
- Model uses its own exciton routine based on that of Griffin
  - Kalbach matrix elements used
  - level densities parametrized vs. Z and A

#### Bertini Cascade Modeling Sequence (3)

- Cascade ends and exciton model takes over when secondary KE drops below 20% of its original value or 7 X nuclear binding energy
- Nuclear evaporation follows for most nuclei
  - emission continues as long as excitation is large enough to remove a particle.
- For light, highly excited nuclei, Fermi breakup
- Fission included in fully phenomenological way

#### Binary Cascade Modeling (1)

- Nucleon-nucleon scattering (t-channel) resonance excitation cross-sections are derived from p-p scattering using isospin invariance, and the corresponding Clebsch-Gordan coefficients
  - elastic N-N scattering included
- Meson-nucleon inelastic (except true absorption) scattering modelled as s-channel resonance excitation. Breit-Wigner form used for cross section.
- Resonances may interact or decay
  - nominal PDG branching ratios used for resonance decay
  - masses sampled from Breit-Wigner form

#### Binary Cascade Modeling (2)

- Calculate imaginary part of the R-matrix using free 2-body cross-sections from experimental data and parameterizations
- For resonance re-scattering, the solution of an in-medium BUU equation is used.
  - The Binary Cascade at present takes the following strong resonances into account:
    - The delta resonances with masses 1232, 1600, 1620, 1700, 1900, 1905, 1910, 1920, 1930, and 1950 MeV
    - Excited nucleons with masses 1440, 1520, 1535, 1650, 1675, 1680, 1700, 1710, 1720, 1900, 1990, 2090, 2190, 2220, and 2250 MeV

#### Binary Cascade Modeling (3)

- Nucleon-nucleon elastic scattering angular distributions taken from Arndt phase shift analysis of experimental data
- Pauli blocking implemented in its classical form
  - final state nucleons occupy only states above Fermi momentum
- True pion absorption is modeled as s-wave absorption on quasi-deuterons
- Coulomb barrier taken into account for charged hadrons

#### Binary Cascade Modeling (4)

- If primary below 45 MeV, no cascade, just precompound
- Cascade stops when mean energy of all scattered particles is below A-dependent cut
  - varies from 18 to 9 MeV
- When cascade stops, the properties of the residual exciton system and nucleus are evaluated, and passed to pre-equilibrium de-excitation class (G4PreCompoundModel)

#### Binary Cascade Modeling (5): pre-equilibrium

- Geant4 precompound model is an extension of the binary cascade for lower energies
- It is a variant of the exciton model used in CEM (Gudima et al, 1983)
- This stage lasts until the nuclear system reaches equilibrium.
- Transition to equilibrium is considered consistently, i.e. the physical condition is applied.
- No need of the rough estimation  $\int_{\lambda_n-1}^{t} = \lambda_{\lambda_n-1}^{t}$
- No need for enhancement of equilibrium by means a soft transtion to equilibrium

#### Binary Cascade Modelling (6): equilibrium

After pre-equilibrium the properties of the residual nucleus are evaluated, and passed to the equilibrium de-excitation handler (G4ExcitationHandler)

Three processes are considered:

1. Statistical multifragmentation (Botvina *et al*) (for E\*/A > 3 MeV).


#### Competitors:

- 1. Fission (Bohr-Wheeler model + Amelin prescript.)
- 2. Particle evaporation (Weisskopf-Erwin).


# additional **RESULTS**

(Geant4 official release 9.2 patch p01)


### Proton production at 542 MeV




### Deuteron production at 175 MeV




### Deuteron production at 542 MeV




# Tritium production at 175 MeV



## Tritium production at 542 MeV





# <sup>3</sup>He production at 63 MeV





# Ongoing development effort: CHIPS

 $^{27}$ Al(p,n) reaction at E<sub>p</sub> = 90 MeV












<sup>27</sup>Al(p. <sup>3</sup>He) reaction at E<sub>p</sub> = 90 MeV





<sup>27</sup>Al(p. <sup>4</sup>He) reaction at E<sub>p</sub> = 90 MeV





 $^{209}$ Bi(p,n) reaction at E<sub>p</sub> = 90 MeV













 $^{209}$ Bi(p,t) reaction at E<sub>p</sub> = 90 MeV



<sup>209</sup>Bi(p.<sup>3</sup>He) reaction at E<sub>p</sub> = 90 MeV



<sup>209</sup>Bi(p, <sup>4</sup>He) reaction at E<sub>p</sub> = 90 MeV

