



Contribution ID: 49

Type: **Poster Presentation of 1h45m**

## Stealth superconducting magnet technology for collider IR and injector requirements

*Wednesday, August 30, 2017 1:15 PM (1h 45m)*

The intersection regions and the beam injection channel of a high-energy collider require magnets that must act strongly upon one beam yet not at all on a closely neighboring beam. Designs are presented for three examples: a septum dipole that serves as a forward spectrometer centered on an ion beam after collision, which must clear an electron beam leaving the IP; a final-focus quadrupole that must provide high-gradient focusing of electrons with large aperture but pass a close-lying ion beam; and a high-gradient quadrupole that must operate in the background field of a spectrometer solenoid. All designs use to advantage a new superconducting cable-in-conduit that provides for compact winding, robust end geometry, and in-cable flow of liquid helium.

### Submitters Country

US

**Primary authors:** MCINTYRE, Peter (Texas A&M University); BREITSCHOPF, Jeff (Texas A&M University); Mr CHAVEZ, Daniel (Universidad Guanajuato); GERITY, James (Texas A&M University); Dr SATTAROV, Akhdiyov (Texas A&M University); Mr KELLAMS, Joshua (Accelerator Technology Corp.)

**Presenter:** MCINTYRE, Peter (Texas A&M University)

**Session Classification:** Wed-Af-Po3.01

**Track Classification:** A1 - Superconducting Accelerator Magnets