

Extra Higgs bosons

Filippo Sala

LPTHE Univ. Paris 6



mainly based on Buttazzo S Tesi 1505.05488

FCC-hh Higgs and EWSB group informal meeting, CERN, 28 Oct 2015

- ◊ Can new Higgses be the lightest new particles around?
- ◊ How to look for them?

- ◊ Can new Higgses be the lightest new particles around?

Another light scalar would be hard to justify with anthropic arguments

Extra **Singlet** scalars are ubiquitous, for example in

- Twin Higgs
- Supersymmetry
- Electroweak Baryogenesis (independent of naturalness)

Bottom-up motivation for a Singlet: Higgs couplings fit


Add singlet S and doublet H :

$$h_{\text{LHC}} = h = c_\gamma(c_\delta h_v - s_\delta H) + s_\gamma S$$


$$\left[\frac{g_{htt}}{g_{SM}^{htt}} = c_\gamma(c_\delta + \frac{s_\delta}{\tan \beta}), \quad \frac{g_{hbb}}{g_{SM}^{hbb}} = c_\gamma(c_\delta - s_\delta \tan \beta), \quad \frac{g_{hVV}}{g_{SM}^{hVV}} = c_\gamma c_\delta \right]$$

cont: LHC8 status

dashed: LHC14 projections (300 fb^{-1})



$$s_\gamma^2 = 0, 0.15$$



- ✓ Can new Higgses be the lightest new particles around?
 - ◊ How to look for them?

Generic singlet

[ϕ = singlet-like mass eigenstate]

$$\sin^2 \gamma = \frac{M_{hh}^2 - m_\phi}{m_\phi^2 - m_h^2}$$

Master formula, valid for **any** model

2 free parameters control all pheno! + $\text{BR}_{\phi \rightarrow hh}$ ($= \text{BR}_{\phi \rightarrow ZZ}$ at $m_\phi \gg m_W$)

h : signal strengths $\mu = c_\gamma^2 \times \mu_{\text{SM}}$

ϕ : $\mu(m_\phi) = s_\gamma^2 \times \mu_{\text{SM}}(m_\phi)$ [barring $\phi \rightarrow hh$]

Generic singlet

[ϕ = singlet-like mass eigenstate]

$$\sin^2 \gamma = \frac{M_{hh}^2 - m_\phi^2}{m_\phi^2 - m_h^2}$$

Master formula, valid for **any** model

2 free parameters control all pheno! + $\text{BR}_{\phi \rightarrow hh}$ ($= \text{BR}_{\phi \rightarrow ZZ}$ at $m_\phi \gg m_W$)

h : signal strengths $\mu = c_\gamma^2 \times \mu_{\text{SM}}$ ϕ : $\mu(m_\phi) = s_\gamma^2 \times \mu_{\text{SM}}(m_\phi)$ [barring $\phi \rightarrow hh$]

What does one learn from the potential $f(S)$?

$$\text{BR}_{\phi \rightarrow hh} = \frac{1}{4} - \frac{3}{4} \frac{\nu}{v_s} \frac{\sqrt{M_{hh}^2 - m_h^2}}{m_\phi} + O\left(\frac{\nu^2}{m_\phi^2}\right)$$

$$\frac{g_{h^3}}{g_{h^3}^{\text{SM}}} = 1 + \frac{2}{3} \frac{\nu}{v_s} \frac{\sqrt{M_{hh}^2 - m_h^2}}{m_\phi} \left(\frac{M_{hh}^2}{m_h^2} - 1 \right) + O\left(\frac{\nu^2}{m_\phi^2}\right)$$

Valid for **any** potential!! v_s leading new parameter



Generic singlet: Higgs couplings

$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$

Master formula, valid for **any** model

2 free parameters control all pheno! + $\text{BR}_{\phi \rightarrow hh}$ ($= \text{BR}_{\phi \rightarrow ZZ}$ at $m_\phi \gg m_W$)

h: signal strengths $\mu = c_\gamma^2 \times \mu_{\text{SM}}$



Generic singlet: Higgs couplings



$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$

Master formula, valid for **any** model

2 free parameters control all pheno! + $\text{BR}_{\phi \rightarrow hh}$ ($= \text{BR}_{\phi \rightarrow ZZ}$ at $m_\phi \gg m_W$)

h : signal strengths $\mu = c_\gamma^2 \times \mu_{\text{SM}}$

Add g_{hhh} : could be first deviation seen!



Generic singlet: Higgs couplings



$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$

Master formula, valid for **any** model

2 free parameters control all pheno! + $\text{BR}_{\phi \rightarrow hh}$ ($= \text{BR}_{\phi \rightarrow ZZ}$ at $m_\phi \gg m_W$)

h : signal strengths $\mu = c_\gamma^2 \times \mu_{\text{SM}}$

Add g_{hhh} : could be first deviation seen!



Generic singlet: Higgs couplings vs direct searches



$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$

Master formula, valid for **any** model

2 free parameters control all pheno! + $\text{BR}_{\phi \rightarrow hh}$ ($= \text{BR}_{\phi \rightarrow ZZ}$ at $m_\phi \gg m_W$)

h : signal strengths $\mu = c_\gamma^2 \times \mu_{\text{SM}}$

ϕ : $\mu(m_\phi) = s_\gamma^2 \times \mu_{\text{SM}}(m_\phi)$ [barring $\phi \rightarrow hh$]



Twin Higgs and the NMSSM

$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$

Twin Higgs: $M_{hh}^2 = (m_h^2 + m_\phi^2) v^2 / f^2$

NMSSM: $M_{hh}^2 = m_Z^2 c_{2\beta}^2 + \lambda^2 v^2 s_{2\beta}^2 + \Delta^2$


Take-home messages

Twin Higgs:

- Signal strengths μ_h more effective than direct ϕ searches, unless $m_\phi \sim f$
- no significant deviations in g_{hhh}

NMSSM:

- For μ_h to do better than direct ϕ searches, per-mille precision needed
- g_{hhh} could show significant deviations



...more in back up slides

Twin Higgs and the NMSSM

$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$

Twin Higgs: $M_{hh}^2 = (m_h^2 + m_\phi^2) v^2 / f^2$

NMSSM: $M_{hh}^2 = m_Z^2 c_{2\beta}^2 + \lambda^2 v^2 s_{2\beta}^2 + \Delta^2$


Take-home messages

Twin Higgs:

- Signal strengths μ_h more effective than direct ϕ searches, unless $m_\phi \sim f$
- no significant deviations in g_{hhh}

NMSSM:

- For μ_h to do better than direct ϕ searches, per-mille precision needed
- g_{hhh} could show significant deviations



...more in back up slides

A (necessary) input to plan future HEP strategy



Relevance to Higgs & EWSB working group

- Extra Higgses crucial to understand the EW scale
- h coupling deviations could tell a lot on the type of extra Higgs

h couplings, $\phi \rightarrow VV, hh, ghhh, \dots$

Which interplay with BSM working group?

[for extra Higgses see also F.S. 1509.08655]



Back up

Could the radial mode be the first new particle seen?

Could the radial mode be the first new particle seen?

Why TH interesting? Solves little hierarchy, without coloured top partners

If nothing new at the LHC14, TH models still quite natural!

- Add a Z_2 -symmetric copy of the SM
 - [only copy of top strictly necessary see e.g. J Serra @ MIAPP 2015]
- 8 "Higgs" degrees of freedom - vs 4 in the SM
 - 7 are massless Goldstone bosons
 - one, σ = radial mode of $\mathcal{G} \rightarrow \mathcal{H}$
 - $\langle \sigma \rangle = f$, $m_\sigma \sim f$ conceivable if UV completion is weakly coupled

Could the radial mode be the first new particle seen?

Why TH interesting? Solves little hierarchy, without coloured top partners

If nothing new at the LHC14, TH models still quite natural!

- Add a Z_2 -symmetric copy of the SM
 - [only copy of top strictly necessary see e.g. J Serra @ MIAPP 2015]
- 8 “Higgs” degrees of freedom - vs 4 in the SM
 - 7 are massless Goldstone bosons
 - one, $\sigma =$ radial mode of $\mathcal{G} \rightarrow \mathcal{H}$
 - $\langle \sigma \rangle = f$, $m_\sigma \sim f$ conceivable if UV completion is weakly coupled

Other particles? Either $M \gtrsim 4\pi f$ or very weakly coupled

Extra Higgses in Supersymmetry

Could the singlet-like scalar be the first new particle seen?

Extra Higgses in Supersymmetry

Could the singlet-like scalar be the first new particle seen?


MSSM Fine tuning worse than 1%! [v sensitivity to $m_{\tilde{t}}$ fixed by g]

NMSSM Given a fixed tuning, \tilde{t} and \tilde{g} heavier by $\sim \lambda/g$ than in MSSM

$$\text{NMSSM} = \text{MSSM} + \text{singlet } S$$

$$W = W_{\text{MSSM}} + \lambda S H_u H_d + f(S)$$

Fine tuning better than 5% \rightarrow
[green points, $\tan \beta \lesssim 5$, $\Lambda = 20$ TeV]



Gherghetta et al. 1212.5243
see also Gherghetta et al. 1401.8291
Cao et al. 1409.8431

Higgs as a PNG boson: Twin Higgs

$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\sigma^2 - m_h^2}$$

$$M_{hh}^2 = (m_h^2 + m_\sigma^2) v^2 / f^2$$

Only two free parameters f and m_σ \Rightarrow $\text{BR}_{\sigma \rightarrow hh}$ fixed everywhere

Twin SM \Rightarrow $\text{BR}_{\sigma \rightarrow \text{inv.}} \neq 0$ [equivalence theorem: $\text{BR}_{\sigma \rightarrow \text{inv.}} \rightarrow 3/7$ for $m_\sigma > m_Z \times f/v$]



Higgs as a PNG boson: Twin Higgs

$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\sigma^2 - m_h^2}$$

$$M_{hh}^2 = (m_h^2 + m_\sigma^2) v^2/f^2$$

Only two free parameters f and m_σ \Rightarrow $\text{BR}_{\sigma \rightarrow hh}$ fixed everywhere

Twin SM \Rightarrow $\text{BR}_{\sigma \rightarrow \text{inv.}} \neq 0$ [equivalence theorem: $\text{BR}_{\sigma \rightarrow \text{inv.}} \rightarrow 3/7$ for $m_\sigma > m_Z \times f/v$]



The NMSSM



$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$

$$M_{hh}^2 = m_Z^2 c_{2\beta}^2 + \lambda^2 v^2 s_{2\beta}^2 + \Delta^2$$

Δ = all loop effects, e.g. top-stop

Here $\lambda = 1.2$ $\Delta = \text{GeV}$

$\tan \beta$ “small” otherwise EWPT



The NMSSM



$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$

$$M_{hh}^2 = m_Z^2 c_{2\beta}^2 + \lambda^2 v^2 s_{2\beta}^2 + \Delta^2$$

Δ = all loop effects, e.g. top-stop

Here $\lambda = 1.2$ $\Delta = 70$ GeV

$\tan \beta$ “small” otherwise EWPT



The NMSSM



$$\sin^2 \gamma = \frac{M_{hh}^2 - m_h^2}{m_\phi^2 - m_h^2}$$

$$M_{hh}^2 = m_Z^2 c_{2\beta}^2 + \lambda^2 v^2 s_{2\beta}^2 + \Delta^2$$

Δ = all loop effects, e.g. top-stop

Here $\lambda = 0.7$ $\Delta = 80$ GeV

$\tan \beta$ “small” otherwise EWPT



A paradigmatic example: the NMSSM

NMSSM with $\lambda \sim 1$ and heavy stops & gluinos


[$\lambda \gtrsim 0.7$ needs a completion before
GUT scale]

The scalars are:

CP-even h, h_3, ϕ (from h_ν, H, S)

CP-odd A, A_s

H^\pm



A paradigmatic example: the NMSSM


NMSSM with $\lambda \sim 1$ and heavy stops & gluinos

$[\lambda \gtrsim 0.7 \text{ needs a completion before GUT scale}]$


The scalars are:

CP-even h, h_3, ϕ (from h_ν, H, S)

$$\mathcal{H}_{\text{ph}} \equiv \begin{pmatrix} h_3 \\ h \\ \phi \end{pmatrix} = R^T \begin{pmatrix} H \\ h_\nu \\ S \end{pmatrix}, \quad R = \boxed{R_\delta^{12} R_\gamma^{23} R_\sigma^{13}}$$



$\gamma = h_\nu - S$ mixing



A motivated limiting case


$m_{h_3} \gg m_{h,\phi}$ and $\sigma, \delta \rightarrow 0$

Fully mixed case and a $\gamma\gamma$ signal

Singlet-like state lighter than 125 GeV

Hard to see, could it explain this hint?

SUSY 2015, Tristan du Pree



Fully mixed case and a $\gamma\gamma$ signal

Barbieri et al 1304.3670, 1307.4937


Singlet-like state lighter than 125 GeV

Hard to see, could it explain this hint?


SUSY 2015, Tristan du Pree

$$[m_{h_3} = 500 \text{ GeV}, s_\sigma^2 = 10^{-3}, v_s = v]$$


$$\lambda = 0.1, \Delta_t = 85 \text{ GeV}$$



see also Badziak et al. 1304.5437, ...



$$\lambda = 0.8, \Delta_t = 75 \text{ GeV}$$



Extrapolation of direct searches I

We started from

- i) Collider Reach (β) Salam Weiler 2014
- ii) Thamm Torre Wulzer 1502.01701

m_0 excluded at LHC8, obtain m_1 at future collider via

$$B(s_1, L_1, m_1) = B(s_0, L_0, m_0)$$

$$B(s, L, m) \propto L \times \int d\hat{s} \frac{1}{\hat{s}} \hat{s} \hat{\sigma}(\hat{s}) \frac{d\mathcal{L}}{d\hat{s}}(s)$$

Extrapolation of direct searches I

We started from

- i) Collider Reach (β) Salam Weiler 2014
- ii) Thamm Torre Wulzer 1502.01701

m_0 excluded at LHC8, obtain m_1 at future collider via

$$B(s_1, L_1, m_1) = B(s_0, L_0, m_0)$$

$$B(s, L, m) \propto L \times \frac{\Delta \hat{s}}{m^2} \hat{s} \hat{\sigma}(\hat{s}) \left. \frac{d\mathcal{L}}{d\hat{s}}(s) \right|_{\hat{s}=m^2}$$

Extrapolation of direct searches I

We started from

- i) Collider Reach (β) Salam Weiler 2014
- ii) Thamm Torre Wulzer 1502.01701

m_0 excluded at LHC8, obtain m_1 at future collider via

$$B(s_1, L_1, m_1) = B(s_0, L_0, m_0)$$

$$B(s, L, m) \propto L \times \frac{\Delta \hat{s}}{m^2} \textcolor{red}{c} \left. \frac{d\mathcal{L}}{d\hat{s}}(s) \right|_{\hat{s}=m^2} \quad \hat{s}\hat{\sigma}(\hat{s}) = \textcolor{red}{c} \Rightarrow \frac{d\mathcal{L}}{d\hat{s}} \text{ drives the reach}$$

Extrapolation of direct searches I



We started from

- i) Collider Reach (β) Salam Weiler 2014 ii) Thamm Torre Wulzer 1502.01701

m_0 excluded at LHC8, obtain m_1 at future collider via $B(s_1, L_1, m_1) = B(s_0, L_0, m_0)$

$$B(s, L, m) \propto L \times \frac{\Delta \hat{s}}{m^2} c \frac{d\mathcal{L}}{d\hat{s}}(s) \Big|_{\hat{s}=m^2} \quad \hat{s}\hat{\sigma}(\hat{s}) = c \Rightarrow \frac{d\mathcal{L}}{d\hat{s}} \text{ drives the reach}$$

$$L_1 c_{ij} \frac{d\mathcal{L}_{ij}}{d\hat{s}}(s_1) \Big|_{\hat{s}=m_1^2} = L_0 c_{ij} \frac{d\mathcal{L}_{ij}}{d\hat{s}}(s_0) \Big|_{\hat{s}=m_0^2}$$



Extrapolation of direct searches I



We started from

- i) Collider Reach (β) Salam Weiler 2014 ii) Thamm Torre Wulzer 1502.01701

m_0 excluded at LHC8, obtain m_1 at future collider via $B(s_1, L_1, m_1) = B(s_0, L_0, m_0)$

$$B(s, L, m) \propto L \times \frac{\Delta \hat{s}}{m^2} c \frac{d\mathcal{L}}{d\hat{s}}(s) \Big|_{\hat{s}=m^2} \quad \hat{s}\hat{\sigma}(\hat{s}) = c \Rightarrow \frac{d\mathcal{L}}{d\hat{s}} \text{ drives the reach}$$

$$L_1 c_{ij} \frac{d\mathcal{L}_{ij}}{d\hat{s}}(s_1) \Big|_{\hat{s}=m_1^2} = L_0 c_{ij} \frac{d\mathcal{L}_{ij}}{d\hat{s}}(s_0) \Big|_{\hat{s}=m_0^2}$$



Extrapolation of direct searches I



We started from

- i) Collider Reach (β) Salam Weiler 2014 ii) Thamm Torre Wulzer 1502.01701



m_0 excluded at LHC8, obtain m_1 at future collider via $B(s_1, L_1, m_1) = B(s_0, L_0, m_0)$

$$B(s, L, m) \propto L \times \frac{\Delta \hat{s}}{m^2} c \frac{d\mathcal{L}}{d\hat{s}}(s) \Big|_{\hat{s}=m^2} \quad \hat{s}\hat{\sigma}(\hat{s}) = c \Rightarrow \frac{d\mathcal{L}}{d\hat{s}} \text{ drives the reach}$$

$$L_1 c_{ij} \frac{d\mathcal{L}_{ij}}{d\hat{s}}(s_1) \Big|_{\hat{s}=m_1^2} = L_0 c_{ij} \frac{d\mathcal{L}_{ij}}{d\hat{s}}(s_0) \Big|_{\hat{s}=m_0^2}$$





Extrapolation of direct searches II



Assumptions/limitations

- Not valid if systematics dominate and change significantly from s_0 to s_1

Extrapolation of direct searches II



Assumptions/limitations

- Not valid if systematics dominate and change significantly from s_0 to s_1
- $\hat{s} \gg m_{\text{bkg}}$ [i.e. not valid at $\hat{s} \sim 2m_t$ for $\phi \rightarrow hh(4b)$]
- $\frac{\Delta\hat{s}}{m^2} \ll 1$ i.e. not valid if analysis depends a lot on shape far from peak


An extra doublet-like state H

$$[\gamma, \sigma = 0, \quad m_{h_2} \gg m_{h_1}, m_{h_3}]$$


Barbieri Buttazzo Kannike Sala Tesi 1304.3670, 1307.4937

$$\frac{g_{h_3 tt}}{g_{htt}^{\text{SM}}} = s_\delta - \frac{c_\delta}{t_\beta} \quad \frac{g_{h_3 bb}}{g_{hbb}^{\text{SM}}} = s_\delta + t_\beta c_\delta \quad \frac{g_{h_3 VV}}{g_{hVV}^{\text{SM}}} = s_\delta \quad [\Delta_t = 75 \text{ GeV}]$$

Status fit LHC8:



$m_{H^\pm} > 480 \text{ GeV}$ from $B \rightarrow X_s \gamma$!



$$[\widetilde{\mathcal{M}}_{12}^2(t_\beta, \dots) = 0 \rightarrow \delta = 0]$$

h_3 phenomenology: more similar to MSSM

see e.g. Craig et al. 1504.04630

An extra doublet-like state H


$$[\gamma, \sigma = 0, \quad m_{h_2} \gg m_{h_1}, m_{h_3}]$$

Barbieri Buttazzo Kannike Sala Tesi 1304.3670, 1307.4937


$$\frac{g_{h_3 tt}}{g_{htt}^{\text{SM}}} = s_\delta - \frac{c_\delta}{t_\beta} \quad \frac{g_{h_3 bb}}{g_{hbb}^{\text{SM}}} = s_\delta + t_\beta c_\delta \quad \frac{g_{h_3 VV}}{g_{hVV}^{\text{SM}}} = s_\delta \quad [\Delta_t = 75 \text{ GeV}]$$

Projections fit LHC14 (300 fb⁻¹):

dashed: m_{H^\pm} cont: λ





$m_{H^\pm} > 480 \text{ GeV}$ from $B \rightarrow X_s \gamma$!



$$[\widetilde{\mathcal{M}}_{12}^2(t_\beta, \dots) = 0 \rightarrow \delta = 0]$$

h_3 phenomenology: more similar to MSSM

see e.g. Craig et al. 1504.04630

Status fit LHC8:[dashed: m_{H^\pm} cont: Δ_t]

Red regions excluded by direct searches at LEP and CMS

Projections fit LHC14: above regions completely excluded

[if $\frac{\mu A_t}{m_t^2}$ very large, conclusions could change...]