

Non-Minimally Coupled Inflation with a Pre-Inflation Anamorphic Contracting Era

John McDonald

Lancaster University



Two conceptual problems for inflation and the expanding Universe

1. Inflation requires smooth initial state of superhorizon size
2. Universe approaches a singularity as scale factor $a \rightarrow 0$

Both problems can be addressed via an initial contracting era

Exploit the “Anamorphic Universe” concept:

Ijjas, Steinhardt
JCAP 1510 (2015) 001

=> Early contracting era with consistent non-singular bounce

- Requires a non-minimally coupled scalar

Planck => excellent fit with Salopek-Bardeen-Bond inflation
 (“Higgs Inflation”)

- Also requires a non-minimally coupled scalar

Construct a non-minimally coupled model which is a fusion of an early anamorphic contracting era and late SBB inflation

JMcD, 0511.07835

SBB Inflation

Non-minimally coupled inflation with a (nearly) symmetric vacuum and ϕ^4 potential

Introduced by Salopek, Bardeen and Bond (SBB Inflation)

PR D40 (1989) 1753

(Earlier variants with symmetry-breaking vacuum: Spokoiny model, Induced gravity inflation)

The big advantage is that conventional TeV-scale particle theories with dimensionally natural self-couplings can play the role of the inflaton:

ϕ is the Higgs boson \leftrightarrow Higgs Inflation

Bezrukov, Shaposhnikov
PL B659 (2008) 703

ϕ is a singlet scalar e.g. a dark matter boson \leftrightarrow S-inflation

Lerner, JMcD
PR D80 (2009) 123507

or the 750 GeV resonance if it is a real singlet scalar

JMcD arXiv: 1604.01711

Marzola et al
arXiv: 1512.90136

SBB inflation is in excellent agreement with Planck

Jordan frame:

$$(8\pi G)^{1/2} = 1$$

$$(-, +, +, +)$$

$$S_J = \int d^4x \sqrt{-g} \left[\frac{\Omega^2 R}{2} - \frac{k(\phi)}{2} \partial_\mu \phi \partial^\mu \phi - V_J(\phi) \right]$$

Non-minimally
coupled scalar

$$\text{SBB Inflation} \Leftrightarrow \Omega^2 = 1 + \xi \phi^2 \quad ; \quad k(\phi) = 1 \quad ; \quad V_J(\phi) = \frac{\lambda \phi^4}{4}$$

Einstein frame:

Conformal transformation: $\tilde{g}_{\mu\nu} = \Omega^2 g_{\mu\nu}$

$$S_E = \int d^4x \sqrt{-\tilde{g}} \left[\frac{\tilde{R}}{2} + \left(-\frac{3}{4\Omega^4} \partial_\mu \Omega^2 \partial^\mu \Omega^2 - \frac{k(\phi)}{2\Omega^2} \right) \partial_\mu \phi \partial^\mu \phi - \frac{V_J(\phi)}{\Omega^4} \right]$$

$$\Omega^2 = 1 + \xi\phi^2$$

SBB Inflation =>

$$S_E = \int d^4x \sqrt{-\tilde{g}} \left[\frac{\tilde{R}}{2} + \left(-\frac{3\xi^2\phi^2}{\Omega^4} - \frac{1}{2\Omega^2} \right) \partial_\mu\phi\partial^\mu\phi - \frac{V_J(\phi)}{\Omega^4} \right]$$

Redefine in terms of a canonically normalized scalar field

$$\frac{d\chi}{d\phi} = \sqrt{\frac{\Omega^2 + 6\xi^2\phi^2}{\Omega^4}}$$

=>

$$S_E = \int d^4x \sqrt{-\tilde{g}} \left(\frac{\tilde{R}}{2} - \frac{1}{2} \tilde{g}^{\mu\nu} \partial_\mu\chi\partial_\nu\chi - V_E(\chi) \right)$$


$$V_E(\chi) = \frac{\lambda\phi^4(\chi)}{4\Omega^4}$$

$$\phi \gg 1/\sqrt{\xi}$$

=>

$$V_E(\chi) = \frac{\lambda}{4\xi^2} \left(1 + \exp\left(-\frac{2\chi}{\sqrt{6}}\right) \right)^{-2}$$

Plateau-type potential



Flat enough to support slow-roll inflation

Inflation is studied by introducing an FRW-like metric and quantizing in the Einstein frame :

$$\tilde{g}_{\mu\nu} = \text{diag}(1, \tilde{a}^2(t))$$

Slow-roll solution: $\phi_{\tilde{N}} = \sqrt{4\tilde{N}/3\xi}$

\tilde{N} = Number of
e-folds in E frame
 $\tilde{N} \approx N + \ln(1/\sqrt{N})$

The SBB inflation model is in excellent agreement with Planck:

$$n_s \approx 1 - \frac{2}{\tilde{N}} - \frac{3}{\tilde{N}^2} + O\left(\frac{1}{\tilde{N}^3}\right) = 0.965 \quad \tilde{N} = 58$$

$$r \approx \frac{12}{\tilde{N}^2} + O\left(\frac{1}{\xi\tilde{N}^2}\right) = 3.6 \times 10^{-3} \quad \text{Observable by PIXIE}$$

$$n_s = 0.9677 \pm 0.0060 \quad (68\% \text{ confidence level (CL), Planck TT + lowP + lensing})$$

$$r_{0.002} < 0.11 \quad (95\% \text{ CL, Planck TT + lowP + lensing})$$

We require a smooth initial state for SBB inflation and no initial singularity

This can be achieved via the “Anamorphic Universe” set-up

Anamorphic => Jordan frame contraction = Einstein frame expansion

Original Anamorphic Universe model:

Jordan frame contracting era = Einstein frame inflating era

• An alternative to inflation

⇒ Consistent contracting Universe model for homogeneous, flat Universe with scale-invariant density perturbations

Ijjas, Steinhardt
JCAP 1510 (2015) 001

Alternative use of the Anamorphic era: To create initial conditions for expansion and inflation

We therefore consider an initial anamorphic contracting era which evolves into an expanding era with conventional SBB inflation

JMcD arXiv:1511.07835

Anamorphic Contraction

$$\frac{a_c}{a} = \frac{\Omega(a)}{\Omega(a_c)} \times \frac{\tilde{a}_c}{\tilde{a}}$$

Expansion in the Einstein frame can appear as contraction in the physical Jordan frame if the conformal factor decreases more rapidly than the Einstein scale factor increases

Model tends to the SBB inflation model at small ϕ and anamorphic contraction at large ϕ

$$S_J = \int d^4x \sqrt{-g} \left[\frac{\Omega^2 R}{2} - \frac{k(\phi)}{2} \partial_\mu \phi \partial^\mu \phi - V_J(\phi) \right]$$

$$\Omega^2 = 1 + \xi \phi^2 \quad ; \quad k(\phi) = 1 \quad ; \quad V_J(\phi) = \frac{\lambda \phi^4}{4}$$

SBB
Inflation

$$\Omega^2 = \alpha e^{-2A\phi} \quad ; \quad k(\phi) = -\eta e^{-2A\phi} \quad ; \quad V_J(\phi) = \beta e^{-B\phi}$$

Anamorphic
contraction
[Ijjas, Steinhardt]

$$\Omega^2 = 1 + \frac{\xi \phi^2}{(1 + \gamma \phi^2 e^{2A\phi})} \quad ; \quad k(\phi) = \frac{1 - \gamma \phi^2}{1 + \gamma \phi^2 e^{2A\phi}} \quad ; \quad V_J(\phi) = \frac{\lambda \phi^4}{4 (1 + \gamma \phi^2 e^{B\phi/2})^2}$$

Interpolating
Model
[JMcD]

$$\Rightarrow \text{Anamorphic model parameters:} \quad \eta = 1 \quad \alpha = \frac{\xi}{\gamma} \quad \beta = \frac{\lambda}{4\gamma^2}$$

$$S_E = \int d^4x \sqrt{-\tilde{g}} \left[\frac{\tilde{R}}{2} - \frac{1}{2} \left(6A^2 - \frac{\eta}{\alpha} \right) \partial_\mu \phi \partial^\mu \phi - \frac{\beta e^{(4A-B)\phi}}{\alpha^2} \right]$$

$$\chi = (6A^2 - \eta/\alpha)^{1/2} \phi \quad \text{Canonically-normalized field}$$

$$V_E = \frac{\beta}{\alpha^2} \exp \left(\frac{(4A-B)\chi}{(6A^2 - \eta/\alpha)^{1/2}} \right)$$

$4A > B \Rightarrow \chi$ rolls towards zero \Rightarrow model will transition to SBB inflation at late times

Universe 'expands' in the Einstein frame :

$$\frac{\tilde{a}_c}{\tilde{a}} = e^{\tilde{N}}$$

$$\tilde{N} = \frac{(6A^2 - \eta/\alpha)^{1/2}}{(4A-B)} (\chi - \chi_c)$$

Einstein frame expansion = Jordan frame contraction

$$\text{Conformal factor ratio : } \frac{\Omega}{\Omega_c} = \exp\left(-\frac{A(\chi - \chi_c)}{(6A^2 - \eta/\alpha)^{1/2}}\right)$$

$$\Rightarrow \text{Ratio of scale factor in Jordan frame: } \frac{a_c}{a} = \frac{\Omega}{\Omega_c} e^{\tilde{N}} = \exp(-\Delta(\chi - \chi_c))$$

$$\Delta = \frac{A}{(6A^2 - \frac{\eta}{\alpha})^{1/2}} - \frac{(6A^2 - \frac{\eta}{\alpha})^{1/2}}{(4A - B)}$$


$$\text{This is a contraction if } \Delta > 0 \quad \Rightarrow \quad \boxed{\frac{\eta}{\alpha} > 2A^2 + AB}$$

This can only be satisfied if $\eta > 0$.


$$\Rightarrow \text{“Wrong sign” kinetic term in the Jordan frame } \frac{\eta e^{-2A\phi}}{2} \partial_\mu \phi \partial^\mu \phi$$



But no ghosts, as the kinetic term has the right sign in the Einstein frame, so no problem

Jordan
frame



Einstein
frame





Therefore there exists a consistent slow-roll solution of the interpolation model such that the Universe contracts at early times when $\phi < \phi_c$ and smoothly transitions to SBB inflation at late times when $\phi > \phi_c$

=> Non-singular transition from early contracting era to late-time inflating era


Smooth bounce without generation/amplification of inhomogeneities

[Unlike quantum or ghost-condensate bounces]

Xue, Steinhardt
PRL 105 (2010) 261301

Condition for smooth initial conditions and the onset of expansion

Contraction => sub-horizon modes naturally become super-horizon in the Jordan frame
 Horizon contracts more rapidly than length scale if $\ddot{a} < 0$



Therefore a sub-horizon potential-dominated patch containing the observed Universe can become super-horizon => Can enter Anamorphic contraction

Anamorphic contraction allows a non-singular transition to expansion

=> Preserves smoothness on superhorizon scales at the onset of expansion and inflation

Condition for smoothness $\ddot{a} < 0 \Rightarrow 4A - B > 0$

Stronger condition for smoothness – metric anisotropies should not dominate the potential in the Friedmann equation

$$ds^2 = dt^2 - \sum_{i=1}^3 X_i^2(t) dx_i^2$$

Contribution to Jordan-frame Friedmann equation grows as a^{-6}

Colley
hep-th/0110049

Potential contribution : $V_J / M_{Pl}^2 \text{eff} = \tilde{V}_J / \Omega^2$

$$\frac{V_J}{\Omega^2} = \frac{\beta}{\alpha} \exp\left(-\frac{(B-2A)\chi}{(6A^2 - \frac{\eta}{\alpha})^{1/2}}\right)$$

$$\frac{V_J}{\Omega^2} \propto a^{-r} ; \quad r = \frac{(B-2A)}{\Delta (6A^2 - \frac{\eta}{\alpha})^{1/2}}$$

$r > 6$ then requires: $\frac{16}{5}A > B$

Smoothness + anamorphic slow-roll $\Rightarrow A \sim B$ and $A \approx \frac{1}{\sqrt{6}} \left(\frac{2\gamma}{\xi}\right)^{1/2}$

Observable consequences?

The interpolation model predicts small deviations from the standard SBB inflation model:

$$\Omega^2(\phi) \approx 1 + \xi\phi^2 \boxed{-\xi\gamma\phi^4 + \xi\gamma^2\phi^6}$$

$$k(\phi) \approx 1 \boxed{-2\gamma\phi^2 + 2\gamma^2\phi^4}$$

$$V_J \approx \frac{\lambda}{4}\phi^4 \boxed{-\frac{\lambda\gamma}{2}\phi^6 + \frac{3\lambda\gamma^2}{4}\phi^8}$$

$$\Rightarrow n_s = 1 - \frac{2}{\tilde{N}} + \boxed{\frac{1}{3} \left(\frac{32\gamma\tilde{N}}{3\xi} \right)^2} \quad \text{Correction to spectral index}$$

$$\Delta n_s \leq 0.01 \quad \Rightarrow \quad \gamma \lesssim 27 \left(\frac{60}{\tilde{N}} \right) \left(\frac{\xi}{10^5} \right)$$

$$\phi_c = 1/\sqrt{\gamma} \quad \Rightarrow \quad \phi_c \gtrsim 0.2 \left(\frac{\tilde{N}}{60} \right)^{1/2} \left(\frac{10^5}{\xi} \right)^{1/2} M_{Pl}$$

If the bounce occurs when $\phi_c \sim M_{pl}$, the deviation of n_s from the SBB prediction can be large enough to be observable

Conclusions

Inflation and the expanding Universe require initial conditions

- Inflation requires a potential-dominated superhorizon patch, which requires a solution to the horizon-problem
- Initial singularity unphysical – best to avoid

Both problems can be consistently solved via a non-minimally coupled model interpolating between SBB inflation and anamorphic contraction

Predicts deviation from the SBB inflation model parameters

=> Increase in the spectral index

Potentially observable if bounce is at $\phi \sim M_{pl}$

End