



Contribution ID: 207

Type: **not specified**

Probing classically conformal B-L model with gravitational waves

Tuesday, 12 July 2016 12:45 (20 minutes)

We study the cosmological history of the classical conformal B-L gauge extension of the standard model, in which the physical scales are generated via the Coleman-Weinberg-type symmetry breaking. Especially, we consider the thermal phase transition of the U(1)B-L symmetry in the early universe and resulting gravitational-wave production. Due to the classical conformal invariance, the phase transition tends to be a first-order one with ultra-supercooling, which enhances the strength of the produced gravitational waves. We show that, requiring (1) U(1)B-L is broken after the reheating, (2) the B-L gauge coupling does not blow up below the Planck scale, (3) the thermal phase transition completes in almost all the patches in the universe, the gravitational wave spectrum can be as large as $\Omega_{\text{GW}} \sim 10^{-8}$ at the frequency $f \sim 0.01$ -1Hz for some model parameters, and a vast parameter region can be tested by future interferometer experiments such as eLISA, LISA, BBO and DECIGO.

Summary

We study the gravitational wave production in so-called “classically-conformal” type models at the time of cosmic phase transition, taking the classically-conformal B-L model as an example.

Primary author: Dr JINNO, Ryusuke (High Energy Accelerator Research Organization (KEK))

Presenter: Dr JINNO, Ryusuke (High Energy Accelerator Research Organization (KEK))

Session Classification: Parallel V

Track Classification: Inflation and alternatives, Strings, Cosmology