

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons

Photon +
jet

Jet-gap-jet
processes

Exclusive
jets

BSM
physics

Conclusions

Physics plans with AFP0+2 and AFP2+2 in Run 2

Rafał Staszewski

on behalf of ATLAS/AFP

Henryk Niewodniczański
Institute of Nuclear Physics
Polish Academy of Sciences
(IFJ PAN Cracow)



LHC Working Group on Forward Physics and Diffraction
27, 28 October 2015

Contents

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons

Photon +
jet

Jet-gap-jet
processes

Exclusive
jets

BSM
physics

Conclusions

1 AFP detectors

2 Soft processes

3 Jet production

4 Electroweak bosons

5 Photon + jet

6 Jet-gap-jet processes

7 Exclusive jets

8 BSM physics

9 Conclusions

AFP Detector

Physics plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons


Photon +
jet

Jet-gap-jet
processes

Exclusive
jets

BSM
physics

Conclusions




Near station (205 m from ATLAS IP):

- tracking detectors: 4 layers, staggered

Far station (217 m from ATLAS IP):

- tracking detectors
- ToF detectors: 4 x 4 bars



The AFP Detector for Run 2

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

APP
detectors

Soft
processes

Jet
production

Electroweak
bosons

Photon +
jet

Jet-gap-jet
processes

Exclusive
jets

BSM
physics

Conclusions

Goals:

- commission the detector; explore the environment close to the LHC beam
- special runs at low- μ , focusing on high-rate diffractive physics processes
- staged installation:
 - Winter 2015-2016 shutdown – installation of a single AFP ‘arm’ with two Roman pot stations, the ‘0+2’ AFP configuration (AFP0+2)
 - Winter 2016-2017 shutdown – installation of the second detector arm

APP 0+2:

- two silicon tracking detectors and a Level-1 Trigger
- physics: soft single diffraction, single diffractive jets, W , jet-gap-jet, exclusive jet production (one tag)

APP 2+2:

- two silicon tracking detectors on second arm and time-of-flight detectors on both far stations
- physics: soft central diffraction, central diffractive jets, jet-gap-jet, γ +jet, exclusive jet production, anomalous couplings

AFP Testbeams

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons

Photon +
jet



Jet-gap-jet
processes

Exclusive
jets

BSM
physics

Conclusions

First integrated AFP was tested in November 2014 and September 2015 at CERN-SPS (120 GeV pions).
AFP prototype: five tracking planes and a Quartic timing system.



- AFP beam tests 2014+15 with first AFP prototype successfully finished
- Tracking + timing integrated into RCE readout
- Integration into ATLAS TDAQ system tested
- Good performance of pixel tracker and LQbar timing detectors
- Analysis efforts of 2015 data on-going
- Preparing for installation

Machine optics (collision optics)

Physics plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons



Photon +
jet

Jet-gap-jet
processes

Exclusive
jets

BSM
physics

Conclusions



Geometric Acceptance

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons

Photon +
jet

Jet-gap-jet
processes

Exclusive
jets

BSM
physics


Conclusions

optics


ALFA

AFP





$\beta^* = 0.55 \text{ m}$
nominal (*collision*)



$\beta^* = 90 \text{ m}$
special (*high- β^**)



$\beta^* = 1000 \text{ m}$
special (*high- β^**)



Reconstruction resolution

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons



Photon +
jet

Jet-gap-jet
processes

Exclusive
jets

BSM
physics

Conclusions



Contents

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons

Photon +
jet

Jet-gap-jet
processes

Exclusive
jets

BSM
physics

Conclusions

1 AFP detectors

2 Soft processes

3 Jet production

4 Electroweak bosons

5 Photon + jet

6 Jet-gap-jet processes

7 Exclusive jets

8 BSM physics

9 Conclusions

Soft Processes

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons


Photon +
jet

Jet-gap-jet
processes


Exclusive
jets

BSM
physics

Conclusions



- Gap measurement in ATLAS does not distinguish SD from DD
- More information about events with forward proton tagging
- High cross sections → low lumi needed → low pile-up possible
- AFP 0+2 – single diffraction
AFP 2+2 – central diffraction
- Goal for 2016 running



Origin of forward protons

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons

Photon +
jet


Jet-gap-jet
processes

Exclusive
jets


BSM
physics

Conclusions


Single Diffraction




Double Diffraction




Non-diffractive



Pythia



Phojet



- High- ξ protons in ND and DD due to hadronisation
- Significant differences between MC generators
- Important also for simulating cosmic air showers

Contents

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons

Photon +
jet

Jet-gap-jet
processes

Exclusive
jets

BSM
physics

Conclusions

1 AFP detectors

2 Soft processes

3 Jet production

4 Electroweak bosons

5 Photon + jet

6 Jet-gap-jet processes

7 Exclusive jets

8 BSM physics

9 Conclusions

Single Diffractive Jet Production

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons


Photon +
jet

Jet-gap-jet
processes

Exclusive
jets



BSM
physics

Conclusions



Motivation:

- gap survival probability
- Pomeron structure studies
- Reggeon contribution
- Pomeron universality between ep and pp



- CERN-PH-LPCC-2015-001
- Goal for 2016 running

Central diffractive jets

Physics plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons


Photon +
jet

Jet-gap-jet
processes

Exclusive
jets


BSM
physics

Conclusions





Motivation:

- measure cross section and gap survival probability
- search for the presence of an additional contribution from Reggeon exchange
- Pomeron structure



Example: purity and statistical significance for AFP and $\beta^* = 0.55$ m.



Contents

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons

Photon +
jet

Jet-gap-jet
processes

Exclusive
jets

BSM
physics

Conclusions

1 AFP detectors

2 Soft processes

3 Jet production

4 Electroweak bosons

5 Photon + jet

6 Jet-gap-jet processes

7 Exclusive jets

8 BSM physics

9 Conclusions

Single Diffractive W Production

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons


Photon +
jet

Jet-gap-jet
processes

Exclusive
jets



BSM
physics

Conclusions



Motivation:

- gap survival probability
- Pomeron structure
- Pomeron flavour composition
- possible also in central diffractive events, but cross section probably too small



Charge asymmetry of diffractive W

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons

Photon +
jet

Jet-gap-jet
processes

Exclusive
jets

BSM
physics

Conclusions

- Definition: $\mathcal{A} = (\sigma_{W^+} - \sigma_{W^-})/(\sigma_{W^+} + \sigma_{W^-})$
- Some experimental systematic uncertainties cancel
- 4 processes (neglecting Cabibbo suppressed ones)

$$u_p + \bar{d}_p \rightarrow W^+, d_p + \bar{u}_p \rightarrow W^-$$
$$\bar{u}_p + d_p \rightarrow W^-, \bar{d}_p + u_p \rightarrow W^+$$

$$u_p \neq d_p, u_p \neq \bar{u}_p, d_p \neq \bar{d}_p$$




\mathcal{A} is sensitive to R_{ud}

- FPMC results:

R_{ud}	\mathcal{A}
$\frac{1}{2}$	0.185
1	0.096
2	0.019

arxiv: 1510.04218



Contents

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons

Photon +
jet

Jet-gap-jet
processes

Exclusive
jets

BSM
physics

Conclusions

1 AFP detectors

2 Soft processes

3 Jet production

4 Electroweak bosons

5 Photon + jet

6 Jet-gap-jet processes

7 Exclusive jets

8 BSM physics

9 Conclusions

Diffractive $\gamma + \text{Jet}$

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons


Photon +
jet

Jet-gap-jet
processes

Exclusive
jets



BSM
physics

Conclusions



Motivation:

- process observation
- gap survival probability
- Pomeron structure
- Pomeron flavour composition
- possible also in single diffractive processes



Contents

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons

Photon +
jet

Jet-gap-jet
processes

Exclusive
jets

BSM
physics

Conclusions

1 AFP detectors

2 Soft processes

3 Jet production

4 Electroweak bosons

5 Photon + jet

6 Jet-gap-jet processes

7 Exclusive jets

8 BSM physics

9 Conclusions

Diffractive Jet-Gap-Jet

Physics plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons



Photon +
jet

Jet-gap-jet
processes

Exclusive
jets


BSM
physics

Conclusions



Motivation:

- process observation
- gap survival probability
- BFKL effects
- possible also in single diffractive processes
- Phys. Rev. D 87 (2013) 3, 034010



Contents

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons

Photon +
jet

Jet-gap-jet
processes

Exclusive
jets

BSM
physics

Conclusions

1 AFP detectors

2 Soft processes

3 Jet production

4 Electroweak bosons

5 Photon + jet

6 Jet-gap-jet processes

7 Exclusive jets

8 BSM physics

9 Conclusions

Exclusive Jets

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons

Photon +
jet

Jet-gap-jet
processes


Exclusive
jets

BSM
physics

Conclusions


Exclusive jets

- Two intact protons
- No Pomeron remnants
- All particles measured



For comparison: CD (DPE) jets

- Two intact protons
- Pomeron remnants
- Remnants escape



- Motivation: verification of QCD production models, unintegrated gluon PDFs
- Small cross section for exclusive processes → measurement with two proton tags needs high luminosity
- Low luminosity – use only single tag events, but less pile-up background
- All particles measured → strong kinematic constraints between central state and each of the forward protons

Exclusive jets measurement

Physics plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons



Photon +
jet

Jet-gap-jet
processes


Exclusive
jets

BSM
physics

Conclusions



- Low cross section → high pile-up conditions → large backgrounds
- Background reduction possible due to kinematic correlations
- Data-driven background estimation needed



Exclusive Jet Production (Single Tag)

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons


Photon +
jet

Jet-gap-jet
processes

Exclusive
jets



BSM
physics

Conclusions



Motivation:

- bigger cross section → lower luminosity necessary
- less background reduction possibilities → low pile-up
- possible contribution from *semi-exclusive* processes (remnant on one side)



Eur. Phys. J. C 75 (2015) 320

Contents

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons

Photon +
jet

Jet-gap-jet
processes

Exclusive
jets

BSM
physics

Conclusions

1 AFP detectors

2 Soft processes

3 Jet production

4 Electroweak bosons

5 Photon + jet

6 Jet-gap-jet processes

7 Exclusive jets

8 BSM physics

9 Conclusions

Anomalous couplings

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons

Photon +
jet


Jet-gap-jet
processes

Exclusive
jets

BSM
physics

Conclusions

- $\gamma\gamma WW$, $\gamma\gamma ZZ$, $\gamma\gamma\gamma\gamma$ quartic couplings
- Testing BSM models
- Constrained kinematics \rightarrow low background
- Reaching limits predicted by string theory and grand unification models ($10^{-14} - 10^{-13}$ for $\gamma\gamma\gamma\gamma$)



$\gamma\gamma WW$ and $\gamma\gamma ZZ$

Coupling	OPAL limits [GeV $^{-2}$]	Sensitivity for 200 fb^{-1} 5σ	Sensitivity for 200 fb^{-1} 95% CL
a_0^W/Λ^2	[-0.020, 0.020]	$2.7 \cdot 10^{-6}$	$1.4 \cdot 10^{-6}$
a_C^W/Λ^2	[-0.052, 0.037]	$9.6 \cdot 10^{-6}$	$5.2 \cdot 10^{-6}$
a_0^Z/Λ^2	[-0.007, 0.023]	$5.5 \cdot 10^{-6}$	$2.5 \cdot 10^{-6}$
a_C^Z/Λ^2	[-0.029, 0.029]	$2.0 \cdot 10^{-5}$	$9.2 \cdot 10^{-6}$

$\gamma\gamma\gamma\gamma$

Coupling (GeV $^{-4}$)	1 conv. 5σ	1 conv. 95% CL	all 95% CL
ζ_1 f.f.	$1 \cdot 10^{-13}$	$7 \cdot 10^{-14}$	$4 \cdot 10^{-14}$
ζ_1 no f.f.	$3 \cdot 10^{-14}$	$2 \cdot 10^{-14}$	$1 \cdot 10^{-14}$
ζ_2 f.f.	$3 \cdot 10^{-13}$	$1.5 \cdot 10^{-13}$	$8 \cdot 10^{-14}$
ζ_2 no f.f.	$7 \cdot 10^{-14}$	$2 \cdot 10^{-14}$	$2 \cdot 10^{-14}$

Phys.Rev. D81 (2010) 074003, Phys.Rev. D88 (2013) 7, 074029

Contents

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons

Photon +
jet

Jet-gap-jet
processes

Exclusive
jets

BSM
physics

Conclusions

1 AFP detectors

2 Soft processes

3 Jet production

4 Electroweak bosons

5 Photon + jet

6 Jet-gap-jet processes

7 Exclusive jets

8 BSM physics

9 Conclusions

Physics plans for AFP

Physics
plans with
AFP0+2
and
AFP2+2
in Run 2

Rafał
Staszewski

on behalf of
AT-
LAS/AFP

AFP
detectors

Soft
processes

Jet
production

Electroweak
bosons

Photon +
jet

Jet-gap-jet
processes

Exclusive
jets

BSM
physics

Conclusions

- Understanding of soft processes
 - AFP 0+2: single diffraction at high ξ , Reggeon vs Pomeron, non-diffractive forward protons
 - AFP 2+2: central diffraction
- Diffractive factorisation breaking
 - all hard diffractive processes
 - AFP 0+2 – single diffractive
 - AFP 2+2 – central diffractive
- Pomeron structure
 - gluon – diffractive jets (SD and CD)
 - quark (and flavour) – diffractive W, photon + jet
- BFKL
 - jet-gap-jet processes (SD and CD)
- Exclusive processes
 - AFP 0+2 – single tag, *semi-exclusive*
 - AFP 2+2 – using full AFP potential
- BSM studies:
 - AFP 2+2: anomalous $\gamma\gamma WW$, $\gamma\gamma ZZ$, $\gamma\gamma\gamma\gamma$ couplings