

Dark matter searches in ATLAS

Thursday, 28 July 2016 09:30 (30 minutes)

Dark matter particles may be produced at the LHC in combination with other particles, typically from initial state radiation. We present results from the ATLAS experiment from searches for phenomena with jets, photons, heavy quarks, electroweak gauge bosons, or Higgs bosons recoiling against large missing transverse momentum. The measurements are interpreted using several theoretical frameworks including simplified models with pair production of Weakly Interacting Massive Particles, effective field theories, and other beyond the Standard Model scenarios. Constraints from dijet searches are compared with results from the “Mono-X” searches to provide a combined interpretation in the context of simplified models.

Summary

Dark matter particles may be produced at the LHC in combination with other particles, typically from initial state radiation. We present results from the ATLAS experiment from searches for phenomena with jets, photons, heavy quarks, electroweak gauge bosons, or Higgs bosons recoiling against large missing transverse momentum. The measurements are interpreted using several theoretical frameworks including simplified models with pair production of Weakly Interacting Massive Particles, effective field theories, and other beyond the Standard Model scenarios. Constraints from dijet searches are compared with results from the “Mono-X” searches to provide a combined interpretation in the context of simplified models.

Based on (arXiv number)

Primary author: DIEHL, Edward (University of Michigan (US))

Presenter: DIEHL, Edward (University of Michigan (US))

Track Classification: Dark Matter at Particle Colliders