


New Results from RENO


Hyunkwan Seo for the RENO Collaboration
Seoul National University

EPS Conference on High Energy Physics
Venice, Italy, 5-12 July 2017



KNRC
Korea Neutrino Research Center

RENO Collaboration



Reactor Experiment for Neutrino Oscillation

(8 institutions and 40 physicists)


- Chonnam National University
- Dongshin University
- GIST
- Gyeongsang National University
- Kyungpook National University
- Seoul National University
- Seoyeong University
- Sungkyunkwan University

- Total cost : **\$10M**
- Start of project : **2006**
- The first experiment running with both near & far detectors from **Aug. 2011**

YongGwang (靈光) :



RENO Experimental Set-up



RENO Data-taking Status



- Data taking began on Aug. 1, 2011 with both near and far detectors.
(DAQ efficiency : ~95%)

- **A (220 days) : First θ_{13} result**
[11 Aug, 2011~26 Mar, 2012]
PRL 108, 191802 (2012)


- **B (403 days) : Improved θ_{13} result**
[11 Aug, 2011~13 Oct, 2012]
NuTel 2013, TAUP 2013, WIN 2013

- **C (500 days) : First $|\Delta m_{ee}^2|$ result**
Rate+shape analysis (θ_{13} and $|\Delta m_{ee}^2|$)
[11 Aug, 2011 ~ 21 Jan, 2013]
PRL 116, 211801 (2016)
submitted to PRD (arXiv:1610.04326)

- **D (1500 days) : New results**
[11 Aug, 2011 ~ Sep, 2015]





New Results from RENO

- 
- Observation of energy dependent disappearance of reactor neutrinos to measure Δm_{ee}^2 and θ_{13} using 1500 live days of data (Aug. 2011 ~ Sep. 2015)
 - Observation of an excess at ~ 5 MeV in reactor neutrino spectrum using 1500 days of data
 - Measurement of absolute reactor neutrino flux using 1500 days

Reduction of background rates & uncertainties

Allows precise measurements of $\sin^2 2\theta_{13}$ and Δm_{ee}^2




- Accidentals : Additional cuts and improved flashing-PMT removal algorithms
- Cosmogenic $^9\text{Li}/^8\text{He}$: Optimized muon veto criteria
- ^{252}Cf contamination : Improved multiple-neutron removal algorithms

Measured Spectra of IBD Prompt Signal

Preliminary

RENO 1500 days




RENO's observation of 5 MeV excess


Clear excess at 5 MeV

Preliminary

RENO 1500 days



Correlation of 5 MeV Excess with Reactor Power




**5 MeV excess
has a clear
correlation
with reactor
thermal power !**

**The 5 MeV excess
comes from reactors!**

Correlation of 5 MeV excess with ^{235}U isotope fraction

^{235}U fraction corresponds to freshness of reactor fuel




Measurement of Absolute Reactor Neutrino Flux

RENO 1500 days
at near (411 m)

Data / Prediction (Huber + Mueller)


0.946 \pm 0.021



Deficit of observed reactor neutrino fluxes relative to the prediction (Huber + Mueller model) indicates an overestimated flux or possible oscillation to sterile neutrinos

Results from Spectral Fit


Energy-dependent disappearance of reactor antineutrinos




$$\sin^2 2\theta_{13} = 0.086 \pm 0.006(\text{stat.}) \pm 0.005(\text{syst.}) \quad (\pm 9 \%)$$

$$|\Delta m_{ee}^2| = 2.61_{-0.16}^{+0.15} (\text{stat.})_{-0.09}^{+0.09} (\text{syst.}) (\times 10^{-3} \text{eV}^2) \quad (\pm 7 \%)$$

Allowed regions in $|\Delta m_{ee}^2|$ and $\sin^2 2\theta_{13}$



Observed L/E Dependent Oscillation



$$P(\bar{\nu}_e \rightarrow \bar{\nu}_e) \approx 1 - \sin^2 2\theta_{13} \sin^2 \left(\Delta m_{ee}^2 \frac{L}{4E_\nu} \right)$$

More precise measurement of θ_{13} and $|\Delta m_{ee}^2|$

PRL 116, 211801 (2016), Submitted to PRD (arXiv:1610.04326)

500 days	Mean	Stat.	Sys.	Precision
$\sin^2 2\theta_{13}$	0.082	+0.009 -0.009	+0.006 -0.006	12 %
$ \Delta m_{ee}^2 $ ($\times 10^{-3} \text{ eV}^2$)	2.62	+0.21 -0.23	+0.12 -0.13	10 %




New results (preliminary)

1500 days	Mean	Stat.	Sys.	Precision
$\sin^2 2\theta_{13}$	0.086	+0.006 -0.006	+0.005 -0.005	9 %
$ \Delta m_{ee}^2 $ ($\times 10^{-3} \text{ eV}^2$)	2.61	+0.15 -0.16	+0.09 -0.09	7 %


Systematic errors are reduced due to background reduction and larger statistics of control samples

RENO : Plan and Prospects

Plan for RENO data taking



RENO data will be taken for 2 more years from now and it will take 3 additional years for the analysis.



$\sin^2 2\theta_{13}$ and $|\Delta m_{ee}^2|$ will approach to **~6% precision** (our design goal).

According to our recent study, the systematic error of $|\Delta m_{ee}^2|$ is smaller than the statistical error.

	500 days Measured	1500 days Measured (preliminary)	~3500 days Expected
$\sin^2 2\theta_{13}$	12 %	9 %	6 ~ 7 %
$ \Delta m_{ee}^2 $	10 %	7 %	4 ~ 5 %

Summary

- More precise measurements of θ_{13} and Δm_{ee}^2 energy dependent disappearance of reactor neutrinos

(Preliminary)

$$\sin^2 2\theta_{13} = 0.086 \pm 0.006(\text{stat.}) \pm 0.005(\text{syst.}) \quad \pm 0.008 (9 \%)$$


$$|\Delta m_{ee}^2| = 2.61_{-0.16}^{+0.15} (\text{stat.})_{-0.09}^{+0.09} (\text{syst.}) (\times 10^{-3} \text{eV}^2) \quad \pm 0.18 (7 \%)$$

(Preliminary)

- Measured absolute reactor neutrino flux : $R = 0.946 \pm 0.021$
- Observed an excess at 5 MeV in reactor neutrino spectrum
- $\sin^2(2\theta_{13})$ and Δm_{ee}^2 to 6% accuracy after 2 more years data taking
- Additional 2~3 years of data taking under consideration to improve Δm_{ee}^2 accuracy

Thanks for your attention!

RENO Detector




- 354 ID 10" PMTs
- 67 OD 10" PMTs



- Target : **16.5 ton Gd-LS**
($R=1.4\text{m}$, $H=3.2\text{m}$)
- Gamma Catcher : 30 ton LS
($R=2.0\text{m}$, $H=4.4\text{m}$)
- Buffer : 65 ton mineral oil
($R=2.7\text{m}$, $H=5.8\text{m}$)
- Veto : 350 ton water
($R=4.2\text{m}$, $H=8.8\text{m}$)

Detection of Reactor Antineutrinos

(prompt signal)



(delayed signal)

$\sim 180 \mu\text{s}$



$\sim 28 \mu\text{s}$

(0.1% Gd)






▪ Neutrino energy measurement

$$E_{\bar{\nu}} \equiv T_{e^+} + T_n + (M_n - M_p) + m_{e^+}$$

10-40 keV

1.8 MeV



Linear Alkyl Benzene (LAB)

$\gamma (0.511 \text{ MeV})$

$\gamma (0.511 \text{ MeV})$

prompt signal


$30 \mu\text{s}$

$$\sum E_{\gamma} \sim 8 \text{ MeV}$$


Delayed signal

Coincidence of prompt and delayed signals

(prompt signal)




Prompt signal



n-Gd IBD


$\sim 30 \mu\text{s}$

Delayed signal






$\sim 200 \mu\text{s}$


n-H IBD



Delayed Signals from Neutron Capture by Gd



Correlation of 5 MeV Excess with Reactor Power




5 MeV excess
has a clear
correlation
with reactor
thermal power !

The 5 MeV excess
comes from reactors!

Far/Near Shape Analysis for $|\Delta m_{ee}^2|$

Preliminary



Energy-dependent disappearance of reactor antineutrinos

Fit using far-to-near ratio

Observed
Far/Near

Expected
Far/Near

χ^2 fitter


Minimize χ^2 Function

$$\chi^2 = \sum_{P=\text{before, After}} \left\{ \sum_{i=1 \sim N_b} \frac{\left(\frac{N_{obs}^{F,P,i}}{N_{obs}^{N,P,i}} - \frac{N_{Exp}^{F,P,i}}{N_{Exp}^{N,P,i}} \right)^2}{(U_i)^2} \right\} + \text{Pull_Terms}$$


$$U_i = \frac{N_{obs}^{F,i}}{N_{obs}^{N,i}} \cdot \sqrt{\frac{N_{obs}^{F,i} + N_{bkg}^{F,i}}{(N_{obs}^{F,i})^2} + \frac{N_{obs}^{N,i} + N_{bkg}^{N,i}}{(N_{obs}^{N,i})^2}}$$

n-H IBD Analysis

1. Independent measurement of θ_{13} value.
2. Consistency and systematic check on reactor neutrinos.



n-H IBD Event Vertex Distribution




(Work in progress) 400 days of data before ^{252}Cf contamination

$$\sin^2 2\theta_{13} = 0.097 \pm 0.013(\text{stat.}) \pm 0.015(\text{syst.})$$

Light Sterile Neutrino Search Results

- All 500 days of RENO data
- Consistent with standard 3-flavor neutrino oscillation model
- Able to set stringent limits in the region $10^{-3} < \Delta m_{ee}^2 < 0.1 \text{ eV}^2$

(Preliminary)



full curves assumes $\sin^2 2\theta_{14} = 0.1$