

Flavour Physics meets Heavy Higgs Searches

Ayan Paul

ERC Ideas: NPFlavour

INFN, Sezione di Roma. Roma, Italy.



In collaboration with Stefania Gori, Christophe Grojean and Aurelio Juste
1707.0XXXX

Venezia. July 6th 2017.

this is how it all began...



not just a fairytale

- ✓ The flavour paradigm of models with an extra Higgs doublet is often limited to escape flavour bounds. But there there are the recent results for $h \rightarrow \tau\mu$ and $t \rightarrow ch$.
- ✓ Stringent bounds on the masses of the expanded Higgs sector can be avoided by proposing certain flavour textures for the Yukawa interactions.
- ✓ We show that we can go beyond the flavour diagonal regime for the couplings of the SM fermions to the neutral Higgs states, yet respect bounds from flavour physics.
- ✓ Once we allow for one or more of the expanded Higgs family to have lower masses, interesting and yet unexplored collider signatures can arise.
- ✓ We show this with a axion variant model with the right handed top quark charged -1 two Higgs doublets charged 0 and -1 under a Peccei-Quinn symmetry.
- ✓ We also introduce a top-charm mixing between right handed up-quark sector. We implement a similar structure in the lepton sector too.

$$U_R \equiv \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \frac{\rho_u}{2} & \sin \frac{\rho_u}{2} \\ 0 & -\sin \frac{\rho_u}{2} & \cos \frac{\rho_u}{2} \end{pmatrix}, \quad L_R \equiv \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \frac{\rho_\ell}{2} & \sin \frac{\rho_\ell}{2} \\ 0 & -\sin \frac{\rho_\ell}{2} & \cos \frac{\rho_\ell}{2} \end{pmatrix}$$

$$\begin{aligned} \mathcal{L}_Y^u &= -\Phi_1 \bar{u}_{Ra} [Y_{u1}]_{ai} Q_i - \Phi_2 \bar{u}_{R3} [Y_{u2}]_i Q_i + \text{h.c.} \\ &= -\Phi^{\text{SM}} \bar{u}_{Ri} [Y_u^{\text{SM}}]_{ij} Q_j - \Phi' \bar{u}_{Ri} [Y'_u]_{ij} Q_j + \text{h.c.} \end{aligned}$$

Some details

$$Y_{u1} = \begin{pmatrix} * & * & * \\ * & * & * \\ 0 & 0 & 0 \end{pmatrix}, \quad Y_{u2} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ * & * & * \end{pmatrix}$$

neutral Higgs couplings ($H_k^0 = H, h, A$)

$$\Gamma_{u_L^f u_R^i}^{H_k^0} = x_u^k \left(\frac{m_{u_i}}{v_u} \delta_{fi} - \epsilon_{fi}^u \cot \beta \right) + x_d^{k*} \epsilon_{fi}^u$$

$$\Gamma_{d_L^f d_R^i}^{H_k^0} = x_d^k \left(\frac{m_{d_i}}{v_d} \delta_{fi} - \epsilon_{fi}^d \tan \beta \right) + x_u^{k*} \epsilon_{fi}^d$$

$$x_u^k = \left(-\frac{1}{\sqrt{2}} \sin \alpha, -\frac{1}{\sqrt{2}} \cos \alpha, \frac{i}{\sqrt{2}} \cos \beta \right)$$

$$x_d^k = \left(-\frac{1}{\sqrt{2}} \cos \alpha, \frac{1}{\sqrt{2}} \sin \alpha, \frac{i}{\sqrt{2}} \sin \beta \right)$$

charged Higgs couplings

$$\Gamma_{u_L^f d_R^i}^{H^\pm} = \sin \beta \sum_{j=1}^3 V_{fj} \left(\frac{m_{d_i}}{v_d} \delta_{ji} - \epsilon_{ji}^d (\tan \beta + \cot \beta) \right)$$

$$\Gamma_{d_L^f u_R^i}^{H^\pm} = \cos \beta \sum_{j=1}^3 V_{jf}^* \left(\frac{m_{u_i}}{v_u} \delta_{ji} - \epsilon_{ji}^u (\tan \beta + \cot \beta) \right)$$

$\epsilon^d = 0_{3 \times 3}$ \longleftarrow **THDM type II structure**

$$\epsilon^u = \begin{pmatrix} \frac{m_u}{v \cos \beta} & 0 & 0 \\ 0 & \frac{m_c}{v \cos \beta} \frac{1 + \cos \rho_u}{2} & -\frac{m_c \sin \rho_u}{2v \cos \beta} \\ 0 & -\frac{m_t \sin \rho_u}{2v \cos \beta} & \frac{m_t}{v \cos \beta} \frac{1 - \cos \rho_u}{2} \end{pmatrix}$$

$$\epsilon^\ell = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \frac{m_\mu}{v \sin \beta} \frac{1 - \cos \rho_\ell}{2} & \frac{m_\mu \sin \rho_\ell}{2v \sin \beta} \\ 0 & \frac{m_\tau \sin \rho_\ell}{2v \sin \beta} & \frac{m_\tau}{v \sin \beta} \frac{1 + \cos \rho_\ell}{2} \end{pmatrix}$$

\swarrow **Goes beyond the THDM type II structure**
 \swarrow


$$\boxed{\rho_u = \rho_\ell \equiv \rho} \longleftarrow \text{model parameter}$$

lepton - Higgs couplings (charged and neutral)

$$\Gamma_{\ell_L^f \ell_R^i}^{H_k^0} = x_d^k \left(\frac{m_{\ell_i}}{v_d} \delta_{fi} - \epsilon_{fi}^\ell \tan \beta \right) + x_u^{k*} \epsilon_{fi}^\ell,$$


$$\Gamma_{\nu_L \ell_R^i}^{H^\pm} = \sin \beta \sum_{j=1}^3 \left(\frac{m_{\ell_i}}{v_d} \delta_{ji} - \epsilon_{ji}^\ell (\tan \beta + \cot \beta) \right)$$

HEPfit: a Code for the Combination of Indirect and Direct Constraints on High Energy Physics Models.




Higgs Physics

HEPfit can be used to study Higgs couplings and analyze data on signal strengths.




Precision Electroweak

Electroweak precision observables are included in HEPfit



Flavour Physics

The Flavour Physics menu in HEPfit includes both quark and lepton flavour dynamics.



BSM Physics

Dynamics beyond the Standard Model can be studied by adding models in HEPfit.

fits to the Higgs couplings

$$\kappa_{gZ} = \frac{\kappa_g \kappa_Z}{\kappa_h} \quad \text{and} \quad \lambda_{ij} = \frac{\kappa_i}{\kappa_j}, \quad (i, j) = (Z, g), (t, g), (W, Z), (\gamma, Z), (\tau, Z), (b, Z)$$

Higgs width modifier:

$$\kappa_h^2 \simeq 0.57\kappa_b^2 + 0.22\kappa_W^2 + 0.09\kappa_g^2 + 0.06\kappa_t^2 + 0.03\kappa_Z^2 + 0.03\kappa_c^2 \\ + 2.3 \times 10^{-3}\kappa_\gamma^2 + 1.6 \times 10^{-3}\kappa_{Z\gamma}^2 + 10^{-4}\kappa_s^2 + 2.2 \times 10^{-4}\kappa_\mu^2$$

	Mean	RMS
κ_{gZ}	1.090	0.110
λ_{Zg}	1.285	0.215
λ_{tg}	1.795	0.285
λ_{WZ}	0.885	0.095
$ \lambda_{\gamma Z} $	0.895	0.105
$ \lambda_{\tau Z} $	0.855	0.125
$ \lambda_{bZ} $	0.565	0.175

	κ_{gZ}	λ_{Zg}	λ_{tg}	λ_{WZ}	$ \lambda_{\gamma Z} $	$ \lambda_{\tau Z} $	$ \lambda_{bZ} $
κ_{gZ}	1.00	-0.03	-0.24	-0.62	-0.57	-0.38	-0.34
λ_{Zg}	-0.03	1.00	0.51	-0.59	-0.51	-0.62	-0.54
λ_{tg}	-0.24	0.51	1.00	-0.21	-0.23	-0.28	-0.35
λ_{WZ}	-0.62	-0.59	-0.21	1.00	0.66	0.55	0.55
$ \lambda_{\gamma Z} $	-0.57	-0.51	-0.23	0.66	1.00	0.58	0.51
$ \lambda_{\tau Z} $	-0.38	-0.62	-0.28	0.55	0.58	1.00	0.49
$ \lambda_{bZ} $	-0.34	-0.54	-0.35	0.55	0.51	0.49	1.00



Higgs-gauge field coupling modifier:

$$\kappa_W = \kappa_Z = \sin(\beta - \alpha), \\ \kappa_{Z\gamma}^2 = 0.00348\kappa_t^2 + 1.121\kappa_W^2 - 0.1249\kappa_t\kappa_W, \\ \kappa_g^2 = 1.06\kappa_t^2 + 0.01\kappa_b^2 - 0.07\kappa_b\kappa_t, \\ \kappa_\gamma^2 = 1.59\kappa_W^2 + 0.07\kappa_t^2 - 0.66\kappa_W\kappa_t,$$


Higgs-fermion coupling modifier:

$$\kappa_f = \frac{\sqrt{2}v}{m_f} c_f^h$$

Run 1 ATLAS-CMS combination
arXiv:1606.02266



Ayan Paul -- EPS 2017




68.2 %
95.4 %
99.7 %


fits to flavour violating Higgs and top decays

Experiment	$\text{BR}(h \rightarrow \tau\mu)$	$\text{BR}(t \rightarrow ch)$
ATLAS 8 TeV 20.3 fb ⁻¹	$(0.53 \pm 0.51)\%$	$(0.22 \pm 0.14)\%$
CMS 8 TeV 19.7 fb ⁻¹	$(0.84^{+0.39}_{-0.37})\%$	$< 0.40\% \text{ @ } 95\% \text{ CL}^\dagger$
ATLAS 13 TeV 36.1 fb ⁻¹	—	$(0.069^{+0.075}_{-0.054})\%$
CMS 13 TeV 35.9 fb ⁻¹	$(0.00 \pm 0.12)\%$	—
Average	$(0.10 \pm 0.11)\%$	$(0.109 \pm 0.061)\%$


$$\text{BR}(t \rightarrow ch) \simeq 3.24 \times 10^{-2} a^2 \sin^2 \rho.$$



(a) $\text{BR}(t \rightarrow ch)$




(b) $\text{BR}(t \rightarrow ch) \ \& \ \text{BR}(h \rightarrow \tau\mu)$




(c) $\text{BR}(h \rightarrow \tau\mu)$

$$\text{BR}_{\text{exp}}(h \rightarrow \tau\mu) = \frac{\sigma^{pp \rightarrow h}}{\sigma_{\text{SM}}^{pp \rightarrow h}} \text{BR}_{\text{th}}(h \rightarrow \tau\mu) \simeq \frac{(\kappa_g)^2 a^2 \sin^2 \rho}{36.5(\kappa_b)^2 + 14.64 \sin^2(\beta - \alpha) + 5.44(\kappa_g)^2 + 4(\kappa_\tau)^2}$$

$$a = (\tan \beta + \cot \beta) \cos(\beta - \alpha)$$




**Combined with Higgs
couplings data**



68.2 %
95.4 %
99.7 %

Ayan Paul -- EPS 2017



Higgs couplings

combined

average

fits to low energy FCNC and charged current decays

Process	Measurement	SM Prediction
$\text{BR}(b \rightarrow s\gamma)$	$(3.32 \pm 0.15) \times 10^{-4}$	$(3.36 \pm 0.23) \times 10^{-4}$
$\text{BR}(B \rightarrow \tau\nu)$	$(1.06 \pm 0.19) \times 10^{-4}$	$(0.807 \pm 0.061) \times 10^{-4}$
R_D	0.403 ± 0.47	0.299 ± 0.003
R_{D^*}	0.310 ± 0.17	0.257 ± 0.003

$$\text{BR}(B \rightarrow \tau\nu) = \frac{G_F^2 |V_{ub}|^2}{8\pi} m_\tau^2 f_B^2 m_B \left(1 - \frac{m_\tau^2}{m_B^2}\right)^2 \tau_B \left|1 + \frac{m_B^2}{m_b m_\tau} \frac{C_R^{ub} - C_L^{ub}}{C_{\text{SM}}^{ub}}\right|^2$$


$$C_R^{ub} = -\frac{1}{m_{H^\pm}^2} \Gamma_{b_R u_L}^{H^\pm} \Gamma_{\nu_L \tau_R}^{H^\pm} \text{ and } C_L^{ub} = -\frac{1}{m_{H^\pm}^2} \Gamma_{b_L u_R}^{H^\pm} \Gamma_{\nu_L \tau_R}^{H^\pm}$$

Large contributions to $B \rightarrow \tau\nu$ are not generated by this model

$$R_D = R_D^{\text{SM}} \left(1 + 1.5 \Re \left(\frac{C_R^{cb} + C_L^{cb}}{C_{\text{SM}}^{cb}} \right) + 1.0 \left| \frac{C_R^{cb} + C_L^{cb}}{C_{\text{SM}}^{cb}} \right|^2 \right),$$

$$R_{D^*} = R_{D^*}^{\text{SM}} \left(1 + 0.12 \Re \left(\frac{C_R^{cb} - C_L^{cb}}{C_{\text{SM}}^{cb}} \right) + 0.05 \left| \frac{C_R^{cb} - C_L^{cb}}{C_{\text{SM}}^{cb}} \right|^2 \right),$$

R_D and R_{D^*} are not explained by this model but the fit to the parameter space is affected by these measurements



$b \rightarrow s\gamma$

$$\delta C_7^0 = \frac{v^2}{\lambda_t m_b} \sum_{j=1}^3 \Gamma_{u_R^j s_L}^{H^{\pm*}} \Gamma_{u_L^j b_R}^{H^\pm} \frac{C_{7,XY}^0(y_j)}{m_{u_j}} + \frac{v^2}{\lambda_t} \sum_{j=1}^3 \Gamma_{u_R^j s_L}^{H^{\pm*}} \Gamma_{u_R^j b_L}^{H^\pm} \frac{C_{7,YY}^0(y_j)}{m_{u_j}^2},$$

$$\delta C_8^0 = \frac{v^2}{\lambda_t m_b} \sum_{j=1}^3 \Gamma_{u_R^j s_L}^{H^{\pm*}} \Gamma_{u_L^j b_R}^{H^\pm} \frac{C_{8,XY}^0(y_j)}{m_{u_j}} + \frac{v^2}{\lambda_t} \sum_{j=1}^3 \Gamma_{u_R^j s_L}^{H^{\pm*}} \Gamma_{u_R^j b_L}^{H^\pm} \frac{C_{8,YY}^0(y_j)}{m_{u_j}^2}$$


strong bound on charged Higgs mass (typical of THDM type II) is alleviated because of cancellations with the SM contributions at low $\tan\beta$

$$m_{H^\pm} \gtrsim 580 \text{ GeV @ 95\% CL in THDM type II}$$




combining all constraints

The picture is not only hopeful but quite promising!!

a preference for $\rho \neq 0$





a preference for low $\tan\beta$





The black dots mark the benchmark point with discuss in our study of collider phenomenology

collider phenomenology of the heavy neutral Higgs

$$\cos(\beta-\alpha)=0.125, \tan\beta=1, \rho=1$$



$$\cos(\beta-\alpha)=0.125, \tan\beta=1, \rho=1$$




excluded by 13 TeV
 $gg \rightarrow H \rightarrow ZZ^* \rightarrow 4l$

$$\cos(\beta-\alpha) = 0.125 \text{ and } \tan\beta = 1$$



$$m_{H^\pm} = m_{H^0}$$

marginalized over
 m_{H^\pm}





(a) $\sigma_{gg} \cdot \text{BR}(H \rightarrow tc)(\text{fb})$

(d) $\sigma_{tc} \cdot \text{BR}(H \rightarrow tc)(\text{fb})$

collider phenomenology of the charged Higgs


$$\cos(\beta-\alpha)=0.125, \tan\beta=1, \rho=1$$



(a) $\sigma_{cb} \cdot BR(H^\pm \rightarrow cb)$ (fb)



$$\cos(\beta - \alpha) = 0.125 \text{ and } \tan \beta = 1$$

Already probed by light dijet searches



(b) $\sigma_{cb} \cdot BR(H^\pm \rightarrow tb)$ (fb)

$$\cos(\beta-\alpha)=0.125, \tan\beta=1, \rho=1$$



(c) $\sigma_{cb} \cdot BR(H^\pm \rightarrow Wh)$ (fb)

summary

- ✓ The intricate alleys of a general THDM are not often navigated leaving interesting phenomenology untouched.
- ✓ The (pseudo)scalar family is awaiting the arrival of other members for which we must search in the right place.
- ✓ We also show that these degrees of freedom leave collider signatures that remain unsearched for.
- ✓ At times, these collider signatures can be quite bold and easily searched for.
- ✓ Stringent lower bounds on the mass of the charged Higgs can be alleviated by a more intricate flavour structure of the Yukawa interactions.

The lower bounds on new (pseudo)scalar states, both neutral and charged, should be reconsidered and collider searches should be open to the possibility of production and decays of these states.

Out beyond the ideas of right and
wrong there is a field. I will meet
you there. - Rumi

Thank you...!!



the model

$$\epsilon^d = 0_{3 \times 3} \quad U_R \equiv \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \frac{\rho_u}{2} & \sin \frac{\rho_u}{2} \\ 0 & -\sin \frac{\rho_u}{2} & \cos \frac{\rho_u}{2} \end{pmatrix}, \quad L_R \equiv \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \frac{\rho_\ell}{2} & \sin \frac{\rho_\ell}{2} \\ 0 & -\sin \frac{\rho_\ell}{2} & \cos \frac{\rho_\ell}{2} \end{pmatrix} \quad \epsilon^\ell = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \frac{m_\mu}{v \sin \beta} \frac{1 - \cos \rho_\ell}{2} & \frac{m_\mu \sin \rho_\ell}{2v \sin \beta} \\ 0 & \frac{m_\tau \sin \rho_\ell}{2v \sin \beta} & \frac{m_\tau}{v \sin \beta} \frac{1 + \cos \rho_\ell}{2} \end{pmatrix}$$

$$c_f^h = \frac{m_f}{\sqrt{2}v} \begin{cases} \sin(\beta - \alpha) + \left(\cot \beta - \frac{1 - \cos \rho_u}{2} (\tan \beta + \cot \beta) \right) \cos(\beta - \alpha) & (\text{for } f = t), \\ \sin(\beta - \alpha) - \left(\tan \beta - \frac{1 - \cos \rho_u}{2} (\tan \beta + \cot \beta) \right) \cos(\beta - \alpha) & (\text{for } f = c), \\ \sin(\beta - \alpha) - \tan \beta \cos(\beta - \alpha) & (\text{for the others}) \end{cases}$$

$$\epsilon^u = \begin{pmatrix} \frac{m_u}{v \cos \beta} & 0 & 0 \\ 0 & \frac{m_c}{v \cos \beta} \frac{1 + \cos \rho_u}{2} & -\frac{m_c \sin \rho_u}{2v \cos \beta} \\ 0 & -\frac{m_t \sin \rho_u}{2v \cos \beta} & \frac{m_t}{v \cos \beta} \frac{1 - \cos \rho_u}{2} \end{pmatrix}$$

$$c_f^H = \frac{m_f}{\sqrt{2}v} \begin{cases} \cos(\beta - \alpha) - \left(\cot \beta - \frac{1 - \cos \rho_u}{2} (\tan \beta + \cot \beta) \right) \sin(\beta - \alpha) & (\text{for } f = t), \\ \cos(\beta - \alpha) + \left(\tan \beta - \frac{1 - \cos \rho_u}{2} (\tan \beta + \cot \beta) \right) \sin(\beta - \alpha) & (\text{for } f = c), \\ \cos(\beta - \alpha) + \tan \beta \sin(\beta - \alpha) & (\text{for the others}) \end{cases}$$

$$\begin{aligned} c_{23}^h &= \frac{m_t}{2\sqrt{2}v} (\cot \beta + \tan \beta) \cos(\beta - \alpha) \sin \rho_u, \\ c_{32}^h &= \frac{m_c}{2\sqrt{2}v} (\cot \beta + \tan \beta) \cos(\beta - \alpha) \sin \rho_u, \\ c_{23}^H &= -\frac{m_t}{2\sqrt{2}v} (\cot \beta + \tan \beta) \sin(\beta - \alpha) \sin \rho_u, \\ c_{32}^H &= -\frac{m_c}{2\sqrt{2}v} (\cot \beta + \tan \beta) \sin(\beta - \alpha) \sin \rho_u, \\ c_{23}^A &= \frac{m_t}{2\sqrt{2}v} (\cot \beta + \tan \beta) \sin \rho_u, \\ c_{32}^A &= \frac{m_c}{2\sqrt{2}v} (\cot \beta + \tan \beta) \sin \rho_u. \end{aligned}$$

$$c_f^A = \frac{m_f}{\sqrt{2}v} \begin{cases} -\cot \beta + \frac{1 - \cos \rho_u}{2} (\tan \beta + \cot \beta) & (\text{for } f = t), \\ \tan \beta - \frac{1 - \cos \rho_u}{2} (\tan \beta + \cot \beta) & (\text{for } f = c), \\ \tan \beta & (\text{for the others}) \end{cases}$$

To my Mother and Father, who showed me what I could do,
and to Ikaros, who showed me what I could not.

“To know what no one else does, what a pleasure it can be!”

– adopted from the words of
Eugene Wigner.

