

Contribution ID: 522

Type: Poster Presentation

On the coherent inelastic binary and multiparticle processes in ultrarelativistic hadron-nucleus, photon-nucleus and nucleus-nucleus collisions

The coherent inelastic processes of the type $a \to b$, which may take place in the interaction of hadrons and γ quanta with nuclei at very high energies (the nucleus remains the same), are theoretically investigated. For taking into account the influence of matter inside the nucleus, the optical model based on the concept of refraction index is applied. Analytical formulas for the effective cross section $\sigma_{\rm coh}(a \to b)$ are obtained, taking into account that at ultrarelativistic energies the main contribution into $\sigma_{\rm coh}$ is provided by very small transferred momenta in the vicinity of the minimum longitudinal momentum transferred to the nucleus. It is shown that the cross section $\sigma_{\rm coh}$ may be expressed through the "forward" amplitudes of inelastic scattering $f_{a+N\to b+N}(0)$ and elastic scattering

 $f_{a+N \rightarrow a+N}(0), f_{b+N \rightarrow b+N}(0)$ on a separate nucleon,

and it depends on the ratios L_a/R , L_b/R , where L_a and L_b are the respective mean free paths in the nucleus matter for the particles a, b and R is the nuclear radius.

In doing so, several characteristic cases are considered in detail. In particular, when $L_a/R \gg 1$, but $L_b/R \ll 1$ (or $L_a/R \ll 1$, but $L_b/R \gg 1$), then $\sigma_{\rm coh}$ is equal to the ratio of the "forward" cross sections of inelastic scattering

 $a+N \rightarrow b+N$ and elastic

scattering of the particle \boldsymbol{b} (or $\boldsymbol{a})$ on a nucleon, multiplied by

the cross section of scattering on the "black" nucleus πR^2 .

If $L_a/R \gg 1$ and $L_b/R \gg 1$,

then $\sigma_{\rm coh} \sim R^4/k^2$, where k is the initial energy of the particle a in the laboratory frame.

The above formalism is generalized also for coherent inelastic multiparticle processes on a nucleus of the type $a \rightarrow \{b_1, b_2, b_3...b_i\}$ and for the case of coherent processes in collisions of two ultrarelativistic nuclei.

Experimental Collaboration

Authors: Dr LYUBOSHITZ, Valery (Joint Institute for Nuclear Research, Dubna); Dr LYUBOSHITZ, Vladimir (Joint Institute for Nuclear Research, Dubna)

Presenter: Dr LYUBOSHITZ, Valery (Joint Institute for Nuclear Research, Dubna)

Session Classification: Poster session

Track Classification: Heavy Ion Physics