



Contribution ID: 567

Type: **Poster Presentation**

## Generalized Gounaris-Sakurai formula and $\rho(770)$ , $\rho'(1450)$ and $\rho''(1700)$ masses and widths

It is demonstrated that Gounaris-Sakurai model of the pion electromagnetic form factor is based on the P-wave iso-vector  $\pi\pi$  scattering phase-shift given by a generalized effective-range formula of the Chew-Mandelstam type, valid exclusively only at the elastic region up to  $1GeV^2$ . Therefore the Gounaris-Sakurai model is justified to be used in a determination of the  $\rho(770)$  meson parameters from existing data, however, in no case in a determination of the inelastic  $\rho'(1450)$  and  $\rho''(1700)$  resonance parameters.

We propose the pion electromagnetic form factor model found on the analyticity in the complex energy plane in which all three resonances  $\rho(770)$ ,  $\rho'(1450)$ ,  $\rho''(1700)$  are defined on equal level as poles on unphysical sheets of the corresponding Riemann surface. The  $\rho(770)$  meson parameters obtained in a such way coincide with the parameters obtained in the framework of the GOPY Roy-like equations analysis..

### Experimental Collaboration

**Authors:** BARTOS, Erik (Institute of Physics, Slovak Academy of Sciences); DUBNICKA, Stanislav (Institute of Physics, Slovak Academy of Sciences); DUBNICKOVA, Anna Zuzana (Comenius University (SK))

**Presenter:** BARTOS, Erik (Institute of Physics, Slovak Academy of Sciences)

**Session Classification:** Poster session

**Track Classification:** QCD and Hadronic Physics