



Contribution ID: 534

Type: **Poster Presentation**

## Theoretical prediction of $\Lambda$ , $\Sigma$ , $\Xi$ hyperon magnetic form factors $|G_M^H(t)|$ behaviours and also ratio $|G_E^H(t)|/|G_M^H(t)|$ in time-like region.

The advanced  $\Lambda$ ,  $\Sigma$ ,  $\Xi$  hyperon Unitary and Analytic electromagnetic structure models are constructed explicitly and behaviours of  $|G_M^H(t)|$  and ratios  $|G_E^H(t)|/|G_M^H(t)|$  are predicted in time-like region as functions of the total energy squared  $t = W^2$  in the c.m. system of  $H\bar{H}$ , which will be found out in intended measurements of the  $\Lambda$ ,  $\Sigma$ ,  $\Xi$  hyperon polar angle  $\Theta_H$  distributions  $F(\cos \Theta_H)$  in  $e^+e^- \rightarrow H\bar{H}$  processes.

### Experimental Collaboration

**Author:** Prof. DUBNICKA, Stanislav (Inst. of physics SAS)

**Co-authors:** Prof. DUBNICKOVA, Anna Zuzana (Dept.of Theor.Physics, Comenius University); Dr ADAMUS-CIN, Cyril (Slovak Academy of Sciences (SK)); Dr BARTOV S, Erik (Inst.of Physics SAS)

**Presenter:** Prof. DUBNICKA, Stanislav (Inst. of physics SAS)

**Session Classification:** Poster session

**Track Classification:** Top and Electroweak Physics