



Contribution ID: 353

Type: **Talk**

NuSTAR Discovery of Galactic Center Hard X-ray Emission

Tuesday, 8 December 2015 15:03 (21 minutes)

The Galactic Center region contains one of the highest concentration of X-ray sources in the Milky Way. Recently NuSTAR, with its sub-arcminute spatial resolution, has discovered an unresolved hard (20-40 keV) X-ray emission within the central 10 pc. This emission is consistent with either stellar origins, such as large populations of intermediate polars, low-mass X-ray binaries, or millisecond pulsars, or diffuse origins, such as cosmic-ray outflows from the supermassive black hole Sagittarius A*. However, each of these explanations implies source properties peculiar to this central region. In particular, the implied average white dwarf mass for the intermediate polar population is $> 0.8M_{\odot}$. The relation of this population to the much lighter $\sim 0.5M_{\odot}$ population implied by previous Chandra and XMM-Newton measurements of the surrounding ~ 50 pc, or to the ~ 0.5 - $0.8M_{\odot}$ populations implied by previous Galactic bulge and ridge, nearby field, or SDSS survey measurements, is unclear. We present in this contribution details of the central hard X-ray discovery, as well as follow-up work on a possible intermediate polar interpretation. Using the broad-band (3-79 keV) energy resolution of NuSTAR and observations of the intermediate polars TV Columbae and IGR J17303-2601, we argue that both the broad-band NuSTAR measurement of the central 10 pc and the low-energy measurements of the surrounding ~ 50 pc could be attributed to a single population of intermediate polars with mean white dwarf mass $\sim 0.85M_{\odot}$.

Primary author: PEREZ, Kerstin

Presenter: PEREZ, Kerstin

Session Classification: 17 - Activity at the galactic center