

Small-x evolution beyond the eikonal approximation

Fabio Dominguez
Universidade de Santiago de Compostela

In collaboration with Tolga Altinoluk


3rd International Conference on the Initial Stages in High-
Energy Nuclear Collisions
Lisbon, Portugal
May 24th, 2016



Recent developments

- Beyond eikonal expansion for finite target thickness
 - Next-to-eikonal corrections for gluon production Altinoluk, Armesto, Beuf, Martinez, Salgado: 1404.2219
 - Next-to-next-to-eikonal corrections Altinoluk, Armesto, Beuf, Moscoso: 1505.01400
 - Lipatov vertex and numerical estimates Altinoluk, Dumitru: 1512.00279
- TMDs and Helicity observables
 - Evolution of gluon TMD Balitsky, Tarasov: 1505.02151
 - Definition and evolution of helicity distributions in CGC Kovchegov, Pitonyak, Sievert: 1511.06737

Eikonal approximation



- Transverse coordinates of partons in the projectile remain frozen during multiple interaction with target
- No emissions inside the target
- Helicity is unchanged along the multiple scatterings


Formally valid for partons with infinite energy

$$\text{Corrections} \sim \frac{1}{s}$$

TMDs


- Evolution of gluon TMD
 - Vertices inside the target

Balitsky's talk today



Balitsky, Tarasov: 1505.02151


- Helicity observables
 - Quark exchanges
 - Helicity flips



Kovchegov, Pitonyak, Sievert:
1511.06737

Finite thickness corrections

- Allow partons to move in transverse coordinate space while traversing the target
- Use in-medium propagator



$$\mathcal{G}_{p^+}^{ab}(x, y) = \int_{y_\perp}^{x_\perp} \mathcal{D}r_\perp \exp \left\{ \frac{ip^+}{2} \int_{y^+}^{x^+} dt \dot{r}_\perp^2(t) \right\} \mathcal{U}^{ab}(x^+, y^+; [r_\perp])$$

$$\xrightarrow{p^+ \rightarrow \infty} \delta^{(2)}(x_\perp - y_\perp) \mathcal{U}^{ab}(x^+, y^+; x_\perp)$$

Eikonal

Expand around classical trajectory

Next-to-eikonal expansion

Altinoluk, Armesto, Beuf,
Martinez, Salgado: 1404.2219

$$\int d^2x_\perp e^{-ik_\perp \cdot x_\perp} \mathcal{G}_{k^+}^{ab}(x, y) \simeq \theta(x^+ - y^+) e^{-ik_\perp \cdot y_\perp} e^{-ik^-(x^+ - y^+)} \left\{ \mathcal{U}(x^+, y^+, y_\perp) + \frac{x^+ - y^+}{k^+} k_\perp^i \mathcal{U}_{(1)}^i(x^+, y^+, y_\perp) + i \frac{x^+ - y^+}{2k^+} \mathcal{U}_{(2)}(x^+, y^+, y_\perp) \right\}^{ab}$$

Decorated Wilson lines:

$$\mathcal{U}_{(1)}^i(x^+, y^+, y_\perp) = \int_{y^+}^{x^+} dz^+ \frac{z^+ - y^+}{x^+ - y^+} \mathcal{U}(x^+, z^+, y_\perp) [igT \cdot \partial_{y_\perp^i} A^-(z^+, y_\perp)] \mathcal{U}(z^+, y^+, y_\perp)$$

Similarly for the other one but with two field insertions

When plugging this into expressions for observables, we get new operators

Decorated dipoles

$$\mathcal{O}_{(1)}^j(x_\perp, y_\perp) = \frac{1}{N_c^2 - 1} \left\langle \text{Tr} \left[\mathcal{U}^\dagger(x_\perp) \mathcal{U}_{(1)}^j(y_\perp) \right] \right\rangle$$


$$\mathcal{O}_{(2)}(x_\perp, y_\perp) = \frac{1}{N_c^2 - 1} \left\langle \text{Tr} \left[\mathcal{U}^\dagger(x_\perp) \mathcal{U}_{(2)}(y_\perp) \right] \right\rangle$$

What about evolution?

- New (and old) operators have to be defined inside a factorization scheme to regulate rapidity divergences
- Current calculations only LO
- Diagrams for the rapidity evolution have to be considered at the same level of accuracy (next-to-eikonal)
- Small- x evolution is driven by emission of soft gluons, which are more likely to be in a region of phase space where the eikonal approximation breaks down
- It has already been established that finite energy considerations play an important role in determining the value of the rapidity to which quantities should be evolved in NLO calculations

Next-to-eikonal evolution for regular dipoles

- Modify derivation of the BK equation
 - Insert expansion of the in-medium propagator for terms where the soft gluon interacts with the medium
 - Include diagrams with the soft emission inside the target
- Analog in theory of jet quenching
 - Hamiltonian formulation for evolution in extended media
 - Motivated by double log contributions to momentum broadening and energy loss



Iancu: 1403.1996

Liou, Mueller, Wu: 1304.7677
Blaizot, Dominguez, Iancu,
Mehtar-Tani: 1311.5823
Blaizot, Mehtar-Tani: 1403.2323

Next-to-eikonal evolution for regular dipoles


Schematically

$$\partial_Y S \sim \int \frac{d\omega}{\omega} \left\{ \mathcal{K} \otimes (SS - S) + \frac{L}{\omega} \mathcal{K}' \otimes \mathcal{O} + \dots \right\}$$

Next-to-eikonal evolution for regular dipoles

Schematically

$$\partial_Y S \sim \int \frac{d\omega}{\omega} \left\{ \mathcal{K} \otimes (SS - S) + \frac{L}{\omega} \mathcal{K}' \otimes \mathcal{O} + \dots \right\}$$


$$\int \frac{d\omega}{\omega^2}$$

Power divergence

Next-to-eikonal evolution for regular dipoles

Schematically

$$\partial_Y S \sim \int \frac{d\omega}{\omega} \left\{ \mathcal{K} \otimes (SS - S) + \frac{L}{\omega} \mathcal{K}' \otimes \mathcal{O} + \dots \right\}$$

$\int \frac{d\omega}{\omega^2}$ Power divergence

- Divergence comes from calculating the kernel for all possible emissions. It can be solved by restricting phase space to the region where next-to-eikonal corrections are relevant
- Once this is done the divergence goes away, including the logarithmic divergence responsible for the evolution

Regulating the kernels

Look at the BK kernel first:

$$\mathcal{K}_{xyz} \propto (A_{xz}^i - A_{yz}^i)^2 \qquad A_{xz}^i = \frac{(x-z)^i}{(x-z)^2} = \int \frac{d^2 k}{2\pi i} e^{ik \cdot (x-z)} \frac{k^i}{k^2}$$

WW Field

\mathcal{K}' is a cumbersome combination of derivatives of WW fields

The boundaries of the phase space for which next-to-eikonal corrections are relevant are put in the momentum integral in the WW field

Similar to kinematical improvement of BK

Kinematical improvement of BK

Beuf: 1401.0313

- It has been shown that finite energy corrections are relevant for NLO calculations in the CGC context
- One of the proposed ways of incorporating these effects in the calculations is to impose a kinematical constraint which is equivalent to ordering in p_- to avoid an over subtraction of the rapidity divergence
- Such approach cuts off the phase space where the next-to-eikonal corrections become relevant, in agreement with our result of no log enhancement from next-to-eikonal terms

Ducloué's talk today

JIMWLK evolution for decorated dipoles

- The eikonal evolution of the decorated dipoles found in calculations for particle production at next-to-eikonal accuracy can be evolved using JIMWLK

$$\begin{aligned} H_{\text{JIMWLK}} \mathcal{O}_{(1)}^i(x_\perp, y_\perp) &= \frac{\alpha_s}{\pi^2} \int_{z_\perp} \mathcal{K}_{xyz} \left\{ \frac{1}{N_c^2 - 1} \left\langle \text{Tr} \left[T^b \mathcal{U}^\dagger(x_\perp) T^a \mathcal{U}_{(1)}^i(y_\perp) \right] \mathcal{U}^{ab}(z_\perp) \right\rangle - N_c \mathcal{O}_{(1)}^i(x_\perp, y_\perp) \right\} \\ &+ \frac{\alpha_s}{\pi^2} \int_{z_\perp} (\partial_{y^i} \mathcal{K}_{xyz}) \left\{ \frac{1}{N_c^2 - 1} \left\langle \text{Tr} \left[T^b \mathcal{U}^\dagger(x_\perp) T^a \mathcal{U}(y_\perp) \right] \mathcal{U}^{ab}(z_\perp) \right\rangle - N_c S(x_\perp, y_\perp) \right\} \end{aligned}$$

Conclusions

- Next-to-eikonal corrections do not have rapidity logs and therefore do not change LL small- x evolution
- This is consistent with previous observations about finite energy considerations for NLO calculations
- Even though the formalism used is the same used in jet quenching calculations, the results are very different since the relevant regions of phase space are very different. The double log enhancement in jet quenching comes from very soft gluons for which the medium is effectively infinite
- JIMWLK can be (formally) used to derive small- x evolution for the new operators involved in next-to-eikonal corrections