

# A global fit of the gamma-ray galactic center excess within the scalar Higgs portal model



[A. Cuoco, B. Eiteneuer, JH, M. Krämer; JCAP 1606 (2016) 050, 1603.08228]

Jan Heisig (RWTH Aachen)



TeVPA2016  
CERN, September 15



# Fermi GeV Galactic Center Excess



[Goodenough, Hooper '09, '10; Vitale, Morselli '09; Hooper, Linden '11; Abazajian, Kaplinghat '12; Hooper, Slatyer '13; Macias, Gordan '13; Huang *et al.* '13; Abazajian *et al.* '14; Daylan *et al.* '14; Zhou *et al.* '14; Calore *et al.* '14; Gaggero *et al.* '15; Cholis *et al.* '15; Bartels *et al.* '15; Lee *et al.* '15; Ajello *et al.* (Fermi-LAT) '15; ...]

⇒ Excess over the known foregrounds in *Fermi*-LAT data

# Fermi GeV Galactic Center Excess



[Goodenough, Hooper '09, '10; Vitale, Morselli '09; Hooper, Linden '11; Abazajian, Kaplinghat '12; Hooper, Slatyer '13; Macias, Gordan '13; Huang *et al.* '13; Abazajian *et al.* '14; Daylan *et al.* '14; Zhou *et al.* '14; Calore *et al.* '14; *et al.* '15; '15; Ajello *et al.* (Fermi-LAT) '15; ...]

**Origine controversially discussed**

⇒ Excess over the known foregrounds in *Fermi*-LAT data



# ~~Fermi GC Excess~~ → Galactic bulge emission




[Christoph's talk]




# Fermi GeV Galactic Center Excess

Astrophysical sources




?




⇒ Excess over the known foregrounds in *Fermi*-LAT data

# Fermi GeV Galactic Center Excess

Astrophysical sources





WIMP Dark Matter



⇒ Excess over the known foregrounds in *Fermi*-LAT data

# Fermi GeV Galactic Center Excess

## WIMP Dark Matter





⇒ Excess over the known foregrounds in *Fermi*-LAT data





# Fermi GeV Galactic Center Excess



## WIMP Dark Matter



⇒ Excess over the known foregrounds in *Fermi*-LAT data

# This work:

- Very simple Dark Matter model  
(singlet scalar Higgs portal)
- Detailed numerical fit involving further constraints  
(invisible Higgs width, LUX, relic density,...)
- Allow for additional non-WIMP DM component  
(PBHs, axions,...)

$$R = \rho_{\text{WIMP}} / \rho_{\text{DM, total}}$$

→ Interesting implications

# Scalar Singlet Higgs Portal Model

[Silveira, Zee '85; McDonald '94; Burgess, Pospelov, Veldhuis: '01; ...]

- Higgs bilinear  $H^\dagger H$  unique (renormalizable) way to directly couple DM to the SM
- Add Singlet Scalar  $S$  with  $Z_2$ -symmetry:

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \frac{1}{2} \partial_\mu S \partial^\mu S - \frac{1}{2} m_{S,0}^2 S^2 - \frac{1}{4} \lambda_S S^4 - \frac{1}{2} \lambda_{HS} S^2 H^\dagger H$$

(before EWSB)



# Scalar Singlet Higgs Portal Model

[Silveira, Zee '85; McDonald '94; Burgess, Pospelov, Veldhuis: '01; ...]

- Higgs bilinear  $H^\dagger H$  unique (renormalizable) way to directly couple DM to the SM
- Add Singlet Scalar  $S$  with  $Z_2$ -symmetry:

$$\mathcal{L} \supset -\frac{1}{2}m_S^2 S^2 - \frac{1}{4}\lambda_S S^4 - \frac{1}{4}\lambda_{HS} h^2 S^2 - \frac{1}{2}\lambda_{HS} v h S^2,$$

where  $m_S^2 = m_{S,0}^2 + \lambda_{HS} v^2/2$ . (after EWSB)

# Scalar Singlet Higgs Portal Model

[Silveira, Zee '85; McDonald '94; Burgess, Pospelov, Veldhuis: '01; ...]

- Higgs bilinear  $H^\dagger H$  unique (renormalizable) way to directly couple DM to the SM
- Add Singlet Scalar  $S$  with  $Z_2$ -symmetry:

$$\mathcal{L} \supset -\frac{1}{2}m_S^2 S^2 - \frac{1}{4}\lambda_S S^4 - \frac{1}{4}\lambda_{HS} h^2 S^2 - \frac{1}{2}\lambda_{HS} v h S^2,$$


where  $m_S^2 = m_{S,0}^2 + \lambda_{HS} v^2/2$ . (after EWSB)

Important for this work

$\Rightarrow$  Only two parameters:  $m_S, \lambda_{HS}$


# Dark Matter annihilation

## ■ Annihilation processes:



SM =  $t, h, Z, W, b, \tau, c, g, \gamma$

Only present above Higgs threshold




[see also e.g.  
Cline, Scott, Kainulainen, Weniger '13;  
Duerr, Pérez, Smirnov '15;  
Beniwal *et al.* '15]




# Dark Matter annihilation

## ■ Annihilation processes:




$$\propto v\lambda_{HS}$$

$$\text{SM} = t, h, Z, W, b, \tau, c, g, \gamma$$




$$\propto (v\lambda_{HS})^2$$




$$\propto \lambda_{HS}$$

Only present above Higgs threshold




[see also e.g.  
Cline, Scott, Kainulainen, Weniger '13;  
Duerr, Pérez, Smirnov '15;  
Beniwal *et al.* '15]

# Gamma-ray spectrum




- Continuous photon spectrum
- Slow in fit  
 $\Rightarrow$  Precompute spectra for all channels with MadGraph/Pythia 8
- During fit: Combine spectra according to contribution

Photon spectra for several masses/couplings:



# $\chi^2$ -computation for the GCE


- Take measured spectrum  $d_i$  and covariance matrix  $\Sigma_{ij}$  from [Calore, Cholis, Weniger: 1409.0042]
- Additional uncertainty on the theoretical prediction of the spectrum  $\Sigma_{ij} \rightarrow \Sigma_{ij} + \Sigma_{ij} \delta_{ij} t_i^2 \sigma_t^2$ ,  $\sigma_t = 10\%$  [Achterberg et al. 1502.05703]
- Large theoretical uncertainties on DM distribution in galaxy:



- Take NFWc profile
- Vary around best fit parameters with MC [from Calore, Cholis, Weniger: 1409.0042]
- ⇒ Distribution for  $J$ -factor
- Determine  $\sigma_\xi$  for  $\xi = \ln(\bar{J}/\bar{J}_{\text{nom}})$

- Compute  $\chi^2$ :

$$\chi^2 = \sum_{i,j} (d_i - e^\xi t_i) (\Sigma_{ij})^{-1} (d_j - e^\xi t_j) + \frac{\xi^2}{(\sigma_\xi)^2}$$






# Constraints on the parameter space

# Constraints on the parameter space

- (i) Collider constraints:  
Higgs invisible BR
- (ii) Direct detection  
constraints: LUX '13  
log-likelihood from  
LUXCalc [Savage et al. 1502.02667]
- (iii) Dwarf Spheroidal Galaxies  
[Fermi-LAT: 1503.02641]




# Constraints on the parameter space

## (iv) Gamma-lines:

[Fermi-LAT: 1506.00013]

$J$ -factor different from GCE  
almost 100% correlation




## (v) Relic density constraint

[Planck: 2013]

Apply 10% theoretical  
uncertainty

[computed with micrOMEGAs]



# Fit parameters and tools

- Allow for additional unspecified DM component

→ WIMP fraction:  $R = \rho_{\text{WIMP}} / \rho_{\text{DM, total}}$

- 4 scan parameters:

$$\begin{aligned} m_S: & \quad 5 \dots 220 \text{ GeV} \\ \lambda_{HS}: & \quad 3 \times 10^{-5} \dots 4\pi \\ \ln(\bar{J} / \bar{J}_{\text{nom}}): & \quad -4\sigma_\xi \dots 4\sigma_\xi \\ R: & \quad 10^{-3} \dots 1 \end{aligned}$$







- Use MultiNest (nested sampling algorithm) [Feroz *et al.* '13]
- Annihilation cross sections and BRs: micrOMEGAs  
[Bélanger *et al.* '14]
- Frequentist interpretation

# Results









GCE only

$$\chi^2_{\text{GCE}} = 19.3$$



GCE+BR<sub>inv</sub>







$$\chi^2_{\text{GCE}} = 25.3$$



# GCE+BR<sub>inv</sub>+LUX





$$\chi^2_{\text{GCE}} = 25.6$$

After LUX: only Higgs-resonant region,  $m_S \approx m_h/2$ , remains





# GCE+BR<sub>inv</sub>+LUX+dwarfs+ $\gamma$ -lines

$$\chi^2_{\text{GCE}} = 26.0$$






Limits from dwarf spheroidal galaxies and gamma lines tighten range for  $\ln(\bar{J} / \bar{J}_{\text{nom}})$






# GCE+BR<sub>inv</sub>+LUX+dwarfs+ $\gamma$ -lines +relic density

$$\chi^2_{\text{GCE}} = 26.8$$



← Interesting structure in  $R$   
Two distinct regions








GCE+BR<sub>inv</sub>+LUX+dwarfs+ $\gamma$ -lines  
+relic density

$$\chi^2_{\text{GCE}} = 26.8$$

$\lambda_{HS}$






$R = \rho_{\text{WIMP}} / \rho_{\text{total}}$



Interesting structure in  $R$   
Two distinct regions

$\log(J_{40^\circ} / J_{40^\circ, \text{nom}})$



$m_S [\text{GeV}]$

$\lambda_{HS}$

$R = \rho_{\text{WIMP}} / \rho_{\text{total}}$





# GCE+BR<sub>inv</sub>+LUX+dwarfs+ $\gamma$ -lines+relic density

- Large velocity dependence around Higgs resonance

- $\sigma v \propto \frac{1}{(m_h^2 - s)^2 + m_h^2 \Gamma_h^2} \simeq \frac{1/m_h^2}{(\delta^2 - v_{\text{rel}}^2)^2 + \Gamma_h^2}, \quad \delta^2 \equiv \frac{m_h^2 - 4m_S^2}{m_h^2}$

- annihilation today:  $v_{\text{rel}} \simeq 10^{-3}$ , freeze-out:  $v_{\text{rel}} \lesssim 0.3$





# GCE+BR<sub>inv</sub>+LUX+dwarfs+ $\gamma$ -lines+relic density

- Large velocity dependence around Higgs resonance

$$\sigma v \propto \frac{1}{(m_h^2 - s)^2 + m_h^2 \Gamma_h^2} \simeq \frac{1/m_h^2}{(\delta^2 - v_{\text{rel}}^2)^2 + \Gamma_h^2}, \quad \delta^2 \equiv \frac{m_h^2 - 4m_S^2}{m_h^2}$$

- annihilation today:  $v_{\text{rel}} \simeq 10^{-3}$ , freeze-out:  $v_{\text{rel}} \lesssim 0.3$





# GCE+BR<sub>inv</sub>+LUX+dwarfs+ $\gamma$ -lines+relic density

- Large velocity dependence around Higgs resonance

- $$\sigma v \propto \frac{1}{(m_h^2 - s)^2 + m_h^2 \Gamma_h^2} \simeq \frac{1/m_h^2}{(\delta^2 - v_{\text{rel}}^2)^2 + \Gamma_h^2}, \quad \delta^2 \equiv \frac{m_h^2 - 4m_S^2}{m_h^2}$$

- annihilation today:  $v_{\text{rel}} \simeq 10^{-3}$ , freeze-out:  $v_{\text{rel}} \lesssim 0.3$





# GCE+BR<sub>inv</sub>+LUX+dwarfs+ $\gamma$ -lines+relic density

- For  $R < 1$ :

→ Relic density:  $\Omega_{\text{DM, total}} = \frac{\Omega_{\text{WIMP}}}{R} \propto \frac{1}{R \langle \sigma v \rangle_{\text{f.o.}}}$

→ GCE flux:  $\phi \propto R^2 \langle \sigma v \rangle_{\text{today}}$





# GCE+BR<sub>inv</sub>+LUX+dwarfs+ $\gamma$ -lines+relic density

- For  $R < 1$ :

→ Relic density:  $\Omega_{\text{DM, total}} = \frac{\Omega_{\text{WIMP}}}{R} \propto \frac{1}{R \langle \sigma v \rangle_{\text{f.o.}}}$

→ GCE flux:  $\phi \propto R^2 \langle \sigma v \rangle_{\text{today}}$







# GCE+BR<sub>inv</sub>+LUX+dwarfs+ $\gamma$ -lines+relic density

- For  $R < 1$ :

→ Relic density:  $\Omega_{\text{DM, total}} = \frac{\Omega_{\text{WIMP}}}{R} \propto \frac{1}{R \langle \sigma v \rangle_{\text{f.o.}}}$

→ GCE flux:  $\phi \propto R^2 \langle \sigma v \rangle_{\text{today}}$





# GCE+BR<sub>inv</sub>+LUX+dwarfs+ $\gamma$ -lines+relic density

- For  $R < 1$ :

→ Relic density:  $\Omega_{\text{DM, total}} = \frac{\Omega_{\text{WIMP}}}{R} \propto \frac{1}{R \langle \sigma v \rangle_{\text{f.o.}}}$

→ GCE flux:  $\phi \propto R^2 \langle \sigma v \rangle_{\text{today}}$





# GCE+BR<sub>inv</sub>+LUX+dwarfs+ $\gamma$ -lines+relic density

- For  $R < 1$ :

→ Relic density:  $\Omega_{\text{DM, total}} = \frac{\Omega_{\text{WIMP}}}{R} \propto \frac{1}{R \langle \sigma v \rangle_{\text{f.o.}}}$

→ GCE flux:  $\phi \propto R^2 \langle \sigma v \rangle_{\text{today}}$





# GCE+BR<sub>inv</sub>+LUX+dwarfs+ $\gamma$ -lines+relic density

- For  $R < 1$ :

→ Relic density:  $\Omega_{\text{DM, total}} = \frac{\Omega_{\text{WIMP}}}{R} \propto \frac{1}{R \langle \sigma v \rangle_{\text{f.o.}}}$


→ GCE flux:  $\phi \propto R^2 \langle \sigma v \rangle_{\text{today}}$




→  $R=1$ : only one spot

→  $R<1$ : two regions remain

■ Consistent fit with per mille WIMP fraction!



$m_S$  [GeV]



$R = \rho_{\text{WIMP}} / \rho_{\text{total}}$

---

# Summary


---

- GCE: Astrophysics of WIMPs?
- Higgs Portal: Unique coupling to minimal DM
- Singlet Scalar Model: Good fit!
- After constraints: Only Higgs-resonance remains
- Allow for additional non-WIMP DM component
- Non-trivial implications for WIMP fraction near resonance (large velocity dependence)




# Back-up I: Future experimental prospects

- Collider constraints: virtually unchanged
- Constraints from dwarfs: General challenge for GCE
- Direct detection projections:



# Back-up II: Photon spectra for best-fit points

GCE only (blue) and after all constraints (red):



# Back-up III: Table with best-fit points

log $L$ contribution	GCE	+BR <sub>inv</sub>	+LUX	+dwarfs	+lines	+relic den.	2nd region
$m_S$ [GeV]	$45.50^{+5.98}_{-5.36}$	$61.07^{+2.65}_{-1.98}$	$61.55^{+1.78}_{-0.85}$	$61.35^{+1.90}_{-0.79}$	$61.46^{+1.87}_{-0.85}$	$62.70^{+0.57}_{-0.18}$	$62.52^{+0.02}_{-0.01}$
$\lambda_{HS}$	$0.17^{+11.67}_{-0.09}$	$0.0125^{+7.31}_{-0.0125}$	$0.0082^{+0.317}_{-0.0082}$	$0.0087^{+0.312}_{-0.0087}$	$0.0082^{+0.315}_{-0.0082}$	$0.022^{+0.015}_{-0.013}$	$0.00029^{+0.0078}_{-0.00010}$
$R$	$0.68^{+0.32}_{-0.65}$	$1.0^{+0.0}_{-1.0}$	$0.99^{+0.01}_{-0.99}$	$1.0^{+0.0}_{-1.0}$	$1.0^{+0.0}_{-1.0}$	$0.054^{+0.141}_{-0.053}$	$0.498^{+0.502}_{-0.496}$
$\log J/J_{\text{nom}}$	$0.0^{+0.44}_{-0.44}$	$-0.05^{+0.48}_{-0.36}$	$0.02^{+0.42}_{-0.43}$	$0.22^{+0.36}_{-0.35}$	$0.12^{+0.31}_{-0.29}$	$0.13^{+0.30}_{-0.32}$	$0.13^{+0.32}_{-0.31}$
$\sigma v$ [ $10^{-26}$ cm <sup>3</sup> /s]	$1.97^{+1034}_{-1.38}$	$1.28^{+4.1\text{e}6}_{-0.61}$	$1.23^{+1.7\text{e}6}_{-0.55}$	$0.96^{+1.3\text{e}6}_{-0.37}$	$1.04^{+1.3\text{e}6}_{-0.42}$	$359^{+9.7\text{e}5}_{-327}$	$4.3^{+1.6\text{e}5}_{-0.9}$
$\sigma v R^2$ [ $10^{-26}$ cm <sup>3</sup> /s]	$0.91^{+0.53}_{-0.35}$	$1.28^{+2.02}_{-0.53}$	$1.21^{+0.68}_{-0.45}$	$0.96^{+0.43}_{-0.31}$	$1.04^{+0.39}_{-0.32}$	$1.06^{+0.42}_{-0.32}$	$1.06^{+0.43}_{-0.31}$
$\chi^2_{\text{GCE}}$	19.3	25.3	25.6	26.0	26.0	26.8	26.7
$p(\chi^2_{\text{GCE}})$	0.57	0.20	0.24	0.22	0.21	0.18	0.18
$p(\text{BR}_{\text{inv}})$	0.0	0.90	0.97	0.97	0.97	1.0	1.0
$p(\text{LUX})$	0.0	0.32	0.62	0.58	0.62	0.84	1.0
$p(\text{dwarfs})$	0.18	0.16	0.18	0.24	0.22	0.22	0.22
$p(\text{lines R3})$	0.5	0.5	0.5	0.5	0.5	0.5	0.5
$p(\text{relic den.})$	0.03	0.0	0.0	0.0	0.0	0.99	1.0