

A photograph of the Alpha Magnetic Spectrometer (AMS-02) experiment mounted on the International Space Station. The AMS-02 logo is prominently displayed on the side of the instrument. The logo is circular with a red border. Inside, the words "Alpha Magnetic Spectrometer" are written in a yellow arc at the top, and "AMS-02" is written in large white letters in the center. Below "AMS-02", it says "Europe • Asia" and "North America". The bottom part of the logo features a stylized illustration of a satellite in space with various scientific symbols like atoms and particles.

Protons and helium in cosmic rays

AMS results and interpretations

Nicola Tomassetti

on behalf of the AMS Collaboration

*Perugia University & INFN Perugia, Italy
LPSC & CNRS/IN2P3 Grenoble, France*

TeV Particle Astrophysics conference “TeVPA-2016”
12 – 16 September 2016 – CERN, Switzerland

The AMS Project



Particle physics detector for high precision CR measurements at TeV energy

[→ see presentation by S. Shael]

Physics goals

- ✓ Antimatter search ($|Z|>1$ anti-nuclei)
- ✓ Dark Matter (light anti-matter & γ -rays)
- ✓ Exotic signals?
- ✓ **Galactic CR astrophysics & γ -rays**
- ✓ Heliophysics (long-term modulation & SEP)
- ✓ Magnetospheric physics & space radiation studies



How it will fulfill these goals?

- **Large collaboration: 16 Countries, 60 Institutes and ~500+ Physicists**
- **Same concept (precision & capability) as the large state-of-the-art HEP detectors**
[but: fitting into the space shuttle & no human intervention after installation]
- **Operation in space, ISS, at 400km, no backgrounds from atmospheric interactions**
[extensive multi-step space qualification tests]
- **Collection power: geometrical factor ($\approx 0.5 \text{ m}^2\text{sr}$) X exposure time (= ISS lifetime)**
[extensive calibration campaigns on ground]

Flux Measurement

Differential flux ($\text{m}^{-2} \text{sr}^{-1} \text{s}^{-1} \text{GV}^{-1}$)

$$\Phi(R) = \frac{N(R, R + \Delta R)}{\epsilon_{Trig}(R) \times A_{Tot}(R) \times T(R) \times \Delta R}$$

- R = p/Z, rigidity; important in magnetic spectrometry & CR astrophysics
N = Number of selected protons (helium) events in R,R+ΔR
T = Effective *exposure time* above geomagnetic cut-off (s)
 A_{Tot} = Total *acceptance* ($\text{m}^2 \text{sr}$) including geom factor + efficiencies
 ϵ_{Trig} = Trigger efficiency


DATA/MC check

- Compute efficiencies
- Interaction studies

SPECTRAL UNFOLDING

- Resolution modeling
- Deconvolution algorithm

Multiple measurements of energy



Tracker, $R = p/Z$
 $MDR \approx 2TV (p); 3TV (He)$

TOF, β
 $\Delta\beta/\beta \approx 1\%$



ECAL, E
 $\Delta E/E (\text{TeV } e^\pm) \sim 2\%$
 $\Delta E/E (\text{TeV } p) \sim 50\%$

RICH, β
 $\Delta\beta/\beta \approx 0.05\%$

Geomagnetic cutoff, R
 $\Delta R/R \approx 10\% \text{ up } \sim 25 \text{ GV}$

Selection of Proton and Helium signals

$$\Phi_i(R_i) = \frac{N_i}{T_i \varepsilon_i A_i \Delta R_i}$$




Acceptance

- ✓ Based on our MC simulation program.
- ✓ Detector response, signal digitization, and full analysis chain simulated.
- ✓ Data/MC corrections and several data-driven crosschecks performed.
- ✓ Role of interactions: flux attenuation in the detector material (C, Al)

[→ see presentation by Q. Yan]

Proton acceptance

Cross-sections for proton interactions off detector material (C, Al) known to few percent at 1 GV and 1.8 TV.




~0.6 – 1 % systematic errors at GV – 2 TV

$$\Phi_i(R_i) = \frac{N_i}{T_i \varepsilon_i A_i \Delta R_i}$$


Helium acceptance

Helium collisions off C and Al: cross section data exist only below 10 GV

New method to determine interactions from ISS data with AMS pointing in horizontal direction



Survival prob. L8 → L9



Survival prob. L2 → L1

~1% < 200 GV increasing to ~2% at highest rigidities

p/He fragmentation studies: 3D CAT

AMS Hadronic Tomography

with the cosmic-ray p/He ratio

Exposure Time: May 20 2011 – May 20 2012


Number of Protons: 3,676,863,217

Number of Helium nuclei: 620,303,906

Rigidity range: 2 GV – 2000 GV

Tomographic plane: $Z = +165 \text{ cm}$


XY pixel area: 1 cm^2



Unfolding

Correction of bin-to-bin migration
is needed due to the finite
tracker resolution

$$\Phi_i(R_i) = \frac{N_i}{T_i \varepsilon_i A_i \Delta R_i}$$




Difference between different unfolding algorithms gives a systematic error $\sim 0.5\%$

Tracker resolution


Protons:

- Resolution function from MC simulation
- Verified with:
 - 400 GeV/c Test Beams data
 - ISS data: tracker residuals, rigidity reconstruction (L1-L8) vs. (L2-L9)



Helium:

- Resolution function from MC simulation
- Verified with ISS data:
 - Tracker residuals
 - Rigidity reconstruction (L1-L8) vs. (L2-L9)





Uncertainty on the flux < 1% below 300 GV rising to 3% at 2 TV


Absolute Rigidity Scale

Two contributions to the uncertainty:

1. **Residual tracker misalignment ($1/\Delta$):**
checked with $E_{ECAL}/R_{Tracker}$ ratio for **electrons and positrons**, limited by the current high energy positron statistics.
Corresponding flux error: 2.5% @1 TV.
2. **Magnetic field:**
Mapping measurement (0.25%) and temperature corrections (0.1%).
Flux error: less than 0.5% at all rigidity




Proton Flux




Helium Flux


Phys. Rev. Lett. 115, 211101



Helium Flux VS Kinetic energy per nucleon



p/He ratio as function of rigidity



Physics behind p-HE anomalies...?


Basic predictions

DSA@SNRs: power-law ($\alpha \sim 2.0 - 2.2$) $Q(E) \approx E^{-\nu}$
QLT: power-law diffusivity ($\delta \sim 0.3 - 0.6$) $K(E) \approx E^{\delta}$
Equilibrium spectra ($E \gg \text{GeV}$) $\phi(E) \sim Q / K \approx E^{-(\nu+\delta)}$


homogeneity
isotropy
stationarity
linearity

Proton and He data cannot be explained by standard models of DSA acceleration and CR transport.


Multi-component flux



Propagation, $K(E)$



Acceleration, $Q(E)$




- Non-linear DSA
- Mach number time-evolution
- Transport in CR-induced turbulence
- Spatial dependent diffusion $K(z, E)$
- Local SNR + Galactic ensemble
- Reaccelerated CRs in weak shocks

p/He ratio: violation of universality in CR acceleration?

- Particle-dependent injection
- Non-uniform He distribution .
- Non-DSA acceleration superbubbles
- Strong unaccounted spallation

$Q(E)$: Revisited CR acceleration



- Non-linear DSA: concavity from backreaction of CRs to the shock structure. Ptuskin 2013 [1212.0381]
 - Time-dependent DSA with decreasing speed of the shock and decreasing $E_{\text{max}} \rightarrow$ low-energy steepening.
- Decreasing p/He ratio from inhomogeneous H/He background density around the SN



- Re-acceleration of pre-existing CRs from (weak) expanding shockwaves. → steep LE flux contribution


Thoudam & Horan 2013 [1308.1357]

Ptuskin et al 2013 [1212.0381]




K(E): Revisited CR propagation

CR propagation under self-generated
(+ pre-existing) turbulence



The transition between CR propagation in their own generated turbulence and pre-existing is seen as a feature at $E \sim 300$ GeV.

Spatial dependent propagation:
Non-separable $K(Z,E) \rightarrow$ two halos




Shallower CR propagation in proximity of the Galactic plane. Two diffusive propagation zones characterized by different diffusion properties.

- ✓ Universal features in all primary and in S/P ratios (p/\bar{p} & $B/C, Li/C$ ratios)
- ✓ Smaller anisotropy amplitude. Connections w/ γ -ray gradient [Gaggero et al 1411.7623]
- ✗ p/He ratio anomaly not addressed. [→ ascribed to acceleration]
- ✗ No solution for the positron excess [→ nearby source of HE $e+e-$]

Nearby sources, but where?

GeV-TeV: Steep flux of Galactic <SNRs>

TeV-PeV: Hard flux from local young SNR




Bernard et al 2013 [1207.4670]

Thoudam & Horandel 2013 [1304.1400]

Erlinin & Wolfendale 2012 [.]

GeV-TeV Steep spectrum from local SNRs

TeV-PeV: Hard flux of Galactic <SNR> ensemble



Connection w/ e+ excess

Explanation for p/He

Implications for O/Fe

TeV anisotropy signatures

Kachelriess et al 1504.06472

NT 2015, 1511.04460

NT 2015, 1509.05774

Kachelriess et al 1505.02720

With extra sources, all anomalies in charged CR spectra can be explained.

No clear predictions for secondary/primary ratios. Implications for DGE emissions.

Many unknowns --- Fine tuning --- Loose predictivity

Conclusions

AMS-02 has measured proton and Helium at 0.5 GeV - 2 TeV/nucleon of kinetic energy
High statistics, extensive studies of the systematics, cross-checks, independent analyses.

- ✓ A smooth spectral change in both fluxes has been reported (“happy ending”)
- ✓ The p/He ratio above \sim 40 GV of rigidity decreases steadily as power-law

Explanations may require a revisit of CR acceleration, diffusive propagation, or the inclusion of multiple sources contributing to the observed flux.

Secondary/primary data from AMS-02: [\rightarrow talks by S. Shael, Q. Yan, A. Bachlechner]

- B/C ratio does *not* harden
- Li/C ratio *seems* to harden
- Pbar/p ratio *flattens*

:-/

It is not clear if a consistent picture is emerging. Multi-TeV nuclear data may be crucial

Protons and helium fluxes in cosmic rays

AMS-02 results and interpretations

Nicola Tomassetti

on behalf of the AMS Collaboration

Perugia University & INFN Perugia, Italy

LPSC & CNRS/IN2P3 Grenoble, France

thank you

This work is supported by all organizations and individuals acknowledged in PRL 115(2015)211101



backup slides

The AMS Project




AMS Collaboration

- 16 countries
- 60 institutes
- 500+ physicists
- 20 years

Project timeline

1994 CONCEPT



1997
AMS-01
PROTOTYPE




1998: STS-91




2008
@CERN
SC MAGNET
BEAM TEST

2000 @CERN
AMS-02 CONSTRUCTION


→ Steadily taking data on the ISS since May 19th 2011



2011
@KSC
AMS-02
INTEGRATION & CR- μ RUN





Multiple measurements of charge



Trigger Efficiency

$$\Phi_i(R_i) = \frac{N_i}{T_i \varepsilon_i A_i \Delta R_i}$$





Trigger efficiency estimation can be done using flight data, thanks to a event sample collected with a dedicated minimum-bias trigger.

$$\varepsilon_{Trig} = \frac{N_{Phys}}{N_{Phys} + 100 \cdot N_{MinBias}}$$

Exposure Time


$$\Phi_i(R_i) = \frac{N_i}{T_i \varepsilon_i A_i \Delta R_i}$$

Exposure time “above cutoff” is function of rigidity
It depends on the ISS orbit along the geomagnetic field



Verifications: Protons


Angular dependence of measured flux at R>45 GV: to verify the systematic error assigned to the **acceptance**.



Time dependence of the high-energy flux: at R>45 GV no observable effects from solar modulation. This verifies that the **detector performance is stable**.


Flux reconstruction in different TOI entry regions of the acceptance. Verification of errors assigned to the tracker alignment.

Measured flux using **inner tracker**, i.e., with a different resolution and MDR. This verifies the errors on **rigidity resolution function and unfolding procedure**



Verifications: Helium


Angular dependence of measured flux at R>45 GV: to verify the systematic error assigned to the **acceptance**.




Time dependence of the high-energy flux: at R>45 GV no observable effects from solar modulation. This verifies that the **detector performance is stable**.

Measured flux using different tracker patterns (**inner-tracker+L1**) with a different resolution and MDR. This verifies the errors on **rigidity resolution function and unfolding procedure**


Measured flux using **inner tracker**, i.e., with a tracker pattern of different rigidity resolution and MDR. This verifies the errors on **rigidity resolution function and unfolding procedure**



Proton Flux VS Kinetic energy



Proton Flux



The spectrum cannot be described by a single power-law function. We obtain a good description using a double power-law:

$$\Phi = C \left(\frac{R}{45 \text{ GV}} \right)^\gamma \left[1 + \left(\frac{R}{R_0} \right)^{\Delta\gamma/s} \right]^s$$

$$\gamma = -2.849_{-0.002}^{+0.002} (\text{fit})_{-0.003}^{+0.004} (\text{sys}) \quad \text{low-rigidity slope}$$

$$\Delta\gamma = 0.133_{-0.021}^{+0.032} (\text{fit})_{-0.030}^{+0.046} (\text{sys}) \quad \text{delta-slope}$$

$$R_0 = 336_{-44}^{+68} (\text{fit})_{-28}^{+66} (\text{sys}) [\text{GV}] \quad \text{critical rigidity}$$

The detailed variation of the high-energy flux can be characterized by measuring the log-slope. As shown, the proton flux experiences a progressive hardening above ~ 100 GV of rigidity.






Figure 3. Proton to Helium ratio as a function of rigidity.

R. Webber et al. 1979


Proton/Helium Flux Ratio




Helium Spectral Index




Proton & Helium Spectral Indices



Proton/Helium Ratio Spectral Index





Helium Flux VS rigidity



Low-Energy He spectrum and solar modulation


Helium spectrum from AMS data

Modulation strength from neutron monitor data (OULU station)




Non-linear CR transport in self-induced turbulence

Self-induced (CR-driven) and pre-existing (SNR-generated) turbulence




R. Aloisio et al. 1507.00594; P. Blasi et al. 1207.3706

- Diffusion to CR-induced turbulence at $E \sim 1\text{-}300 \text{ GeV}$
- Advection to CR-generated Alfvèn waves at $E < 1 \text{ GeV}$
- Diffusion to pre-existing turbulence at $> 300 \text{ GeV}$
- ✓ Flattening in all nuclei and sec/pri ratios
- ✓ Low energy Voyager-1 data.
- ✓ B/C seems to require an additional primary component




New phenomena in cosmic-ray propagation?

Diffusion coefficient is not separable into energy and space coordinates → no power-law
Shallower diffusivity in the region close to the Galactic disk → high-energy flattening



NT, 1204.4492 (2012); NT, 1509.05775 (2015)

- ✓ Predicted flattening in all nuclei and sec/pri ratios
- ✓ Enhanced antimatter production at high energy
- ✓ Connection with gamma-rays and anisotropy





New [nearby-SNR] components in the CR spectrum?

two SNR components in the CR flux

- *Nearby source component for the GeV-TeV flux (and e+ excess)*
Must have $R_{max} \sim TeV$ and secondary production ($B \sim 1uG$, $n > n_{ISM}$, $v \sim 10^7 cm/s \rightarrow OLD\ SNR$)
- *Galactic SNR ensemble*
Must have $R_{max} \sim PeV$: $B \sim 500uG$, $v \sim 10^9 m/s \rightarrow YOUNG\ SNRs$


- ✓ Connection between hadron spectra and positron excess [NT & F.Donato, 1502.06150 (2015)]
- ✓ Predicted features in heavy nuclei: explanation of C/Fe and O/Fe [NT, 1509.05774 (2015)]
- ✓ Connection with p/He ratio anomaly? [NT in preparation] [Kachelriess et al. 1504.06472 (2015)]



p/He ratio: two general considerations on the CR spectrum


1. The data at multi-TeV energy do *not* show evidence of p/He decreasing. The p/He ratio at high-energy is essentially constant.

This trend, if confirmed, will invalidate the existing explanations based on intrinsic properties of the DSA mechanism




2. The AMS data indicate that the proton spectral change is much *smoother* than that reported by PAMELA. The differential slope hardens progressively at $E > 100$ GV, without any spectral kink.

This invalidates the usual considerations that the p/He ratio “cancels out” the features of the single p and He spectra.



The p/He ratio seems rather a ~10-1000 GV feature, vanishing at higher energies

p/He ratio anomaly as signature of a nearby source



The idea. Two different classes of sources contribute to the CR flux. Each class has different spectra and composition.

Low-energy flux (GeV-TeV energies)

Nearby old SNRs. The shock is weaker, the B-field are damping, the DSA is not efficient

→ The injection spectra may be steeper

Higher background density for p+p interactions (to explain the e+ excess), e.g. due a molecular cloud

→ It may be well a hydrogen-rich source

High-energy flux (TeV-PeV energies)

Galactic SNR ensemble. Younger SNRs, with stronger shock and B-field amplification

→ Efficient DSA working up to PV rigidities.

→ Hard acceleration spectra: slope ~2-2.1.

Composition = average Galactic SNRs properties

- ✓ Single p & He spectral hardening → signature of transition to different DSA spectra
- ✓ pHe ratio → signature of transition to different composition of H & He in the medium

Each class of source has elemental-independent spectra → the DSA universality is preserved

p/He ratio: two general considerations on the CR spectrum

