

Study of VBF/VBS in the LHC at 13 TeV and available tools

Raquel Gómez Ambrosio
Università & INFN @Torino

Second Annual Meeting of HiggsTools ITN

April 13, 2016



Outline

Introduction:

VBF/VBS motivation

Tools used for simulation: Monte Carlo generators

PHANTOM

MadGraph5_aMC@NLO

Search for new physics: EFT

Top-down approach

Bottom-up approach


Dim-6 LO EFT

NLO EFT

Next steps



Vector Boson Fusion & Vector Boson Scattering



$$V = W^+, W^-, Z, \gamma$$


Why look at this production channel? Mainly 2 reasons:

- ▶ Subleading channel for H production at LHC.
- ▶ Most important channel for EWSB and unitarity studies.

Plus: “Someone has to do it” : LHC will reach % accuracy \rightarrow EW corrections relevant

Why study VBF at LHC RUN-II ?

- ▶ σ_{VBF} grows with \sqrt{s} .
- ▶ Also the obvious: big luminosity \rightarrow big statistics.



Delayed unitarity

Facts of perturbative unitarity

- ▶ In principle $\sigma_{VBF} \propto s$, i.e. it is divergent when $s \rightarrow \infty$
- ▶ The Higgs boson stabilizes σ_{VBF} , making it converge for $s \rightarrow \infty$


But ...

- ▶ *What happens between the SM scale and the scale for new physics?*
(Recall, VBS is the portal to triple and quartic gauge and Higgs couplings)



Importance of NLO EW corrections

For $M_H = 125$ GeV, in gluon fusion: $\delta_{EW} \approx 5\% \approx \mathcal{O}$ (LHC RUN2)



Tools



PHANTOM: Monte Carlo Event Generator for Six Parton Final States

Scope:

Six fermion final states: VBF, VBS, $t\bar{t}$ production ...

PHANTOM output

Unweighted events (LHA) } Lowest Order: $\mathcal{O}(\alpha^6) + \mathcal{O}(\alpha^4\alpha_s^2)$ + interferences
 Cross sections

Features

- ▶ Uses the “modular helicity formalism” for the calculation.
- ▶ Does **not** use narrow width approximations ($\sigma \times \text{BR}$)
- ▶ Faster than other MC generators, since it has all the master amplitudes implemented.
- ▶ Need to run parton shower (PYTHIA) in order to get final events.

A. Ballestrero, A. Belhouari, G. Bevilacqua, V. Kashkan, E. Maina

MadGraph5_aMC@NLO

“The MC generator that will calculate any amplitude”


- ▶ Predecessors: MadGraph, MadEvent, MadGraph5, MC@NLO
- ▶ Also Lowest Order in the EW sector
- ▶ An interface to PYTHIA is already implemented → good for theorists
- ▶ Different algorithm (QCD inspired)
- ▶ Handicap: Rather difficult to master
- ▶ Highlight: Can accommodate Dim-6 EFT operators

J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H.-S. Shao, T. Stelzer, P. Torrielli, M. Zaro

Q: MADGRAPH or PHANTOM?

A: Rivet

Make sure you understand your MC generators before you claim New Physics!



SEARCHING FOR NEW PHYSICS




Effective field theory

Top-down approach

In the top down approach, we integrate out the heavy fields. The propagator becomes a point.

Recall Fermi theory:



Effective field theory

Top-Down approach

- ▶ Model-dependent, therefore not very efficient . . .
- ▶ Useful for Dark Matter and SUSY searches

Low energy behaviour of standard model extensions

Michele Boggia^{a,1}, Raquel Gomez-Ambrosio^{b,1}, Giampiero Passarino^{b,1}


^a*Albert-Ludwigs-Universität Freiburg, Physikalisches Institut
D-79104, Freiburg, Germany*

^b*Dipartimento di Fisica Teorica, Università di Torino, Italy
INFN, Sezione di Torino, Italy*


arXiv: 1603.03660



Bottom-up approach



- ▶ In the bottom up approach we do exactly the opposite:



- ▶ **Observe:** Unlike in the top-down, the relation is not bijective: one low-energy operator might come from many different UV completions

SM EFT (bottom-up approach)

$$\mathcal{L}_{eff} = \underbrace{\mathcal{L}_{SM}}_{\text{dim 4}} + \underbrace{\frac{1}{\Lambda^2} \sum_k \alpha_k \mathcal{O}_k^{(6)}}_{\text{dim 6}} + \underbrace{\dots}_{\text{higher dim. operators}}$$

- ▶ α_k is the Wilson coefficient of the k^{th} operator \rightarrow needs to be calculated
- ▶ One can build 80 dim-6 operators (compatible with $SU(2) \times SU(3) \times U(1)$ and lepton/baryon conservation)
- ▶ Eqs. of motion, reduce this set to a 59-operator basis
(for one generation of particles! for three \rightarrow 2499 operators)
- ▶ “Warsaw Basis” \rightarrow [arXiv: 1008.4884](https://arxiv.org/abs/1008.4884)



Warsaw Basis: 34 Bosonic + 25 (+5) fermionic operators

X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_{\mu\nu}^A G_{\nu\rho}^B G_{\rho\mu}^C$	Q_φ	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_{\mu\nu}^A G_{\nu\rho}^B G_{\rho\mu}^C$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{w\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_{\mu\nu}^I W_{\nu\rho}^J W_{\rho\mu}^K$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_{\mu\nu}^I W_{\nu\rho}^J W_{\rho\mu}^K$				


$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi e}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$

$(\bar{L}L)(\bar{L}L)$		$(\bar{R}R)(\bar{R}R)$		$(\bar{L}L)(\bar{R}R)$	
Q_{ll}	$(\bar{l}_p \gamma_\mu l_r)(\bar{l}_s \gamma^\mu l_t)$	Q_{ee}	$(\bar{e}_p \gamma_\mu e_r)(\bar{e}_s \gamma^\mu e_t)$	Q_{le}	$(\bar{l}_p \gamma_\mu l_r)(\bar{e}_s \gamma^\mu e_t)$
$Q_{qq}^{(1)}$	$(\bar{q}_p \gamma_\mu q_r)(\bar{q}_s \gamma^\mu q_t)$	Q_{uu}	$(\bar{u}_p \gamma_\mu u_r)(\bar{u}_s \gamma^\mu u_t)$	Q_{lu}	$(\bar{l}_p \gamma_\mu l_r)(\bar{u}_s \gamma^\mu u_t)$
$Q_{qq}^{(3)}$	$(\bar{q}_p \gamma_\mu \tau^I q_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	Q_{dd}	$(\bar{d}_p \gamma_\mu d_r)(\bar{d}_s \gamma^\mu d_t)$	Q_{ld}	$(\bar{l}_p \gamma_\mu l_r)(\bar{d}_s \gamma^\mu d_t)$
$Q_{lq}^{(1)}$	$(\bar{l}_p \gamma_\mu l_r)(\bar{q}_s \gamma^\mu q_t)$	Q_{eu}	$(\bar{e}_p \gamma_\mu e_r)(\bar{u}_s \gamma^\mu u_t)$	Q_{qe}	$(\bar{q}_p \gamma_\mu q_r)(\bar{e}_s \gamma^\mu e_t)$
$Q_{lq}^{(3)}$	$(\bar{l}_p \gamma_\mu \tau^I l_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	Q_{ed}	$(\bar{e}_p \gamma_\mu e_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{qu}^{(1)}$	$(\bar{q}_p \gamma_\mu q_r)(\bar{u}_s \gamma^\mu u_t)$
		$Q_{ud}^{(1)}$	$(\bar{u}_p \gamma_\mu u_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{qu}^{(8)}$	$(\bar{q}_p \gamma_\mu T^A q_r)(\bar{u}_s \gamma^\mu T^A u_t)$
		$Q_{ud}^{(8)}$	$(\bar{u}_p \gamma_\mu T^A u_r)(\bar{d}_s \gamma^\mu T^A d_t)$	$Q_{qd}^{(1)}$	$(\bar{q}_p \gamma_\mu q_r)(\bar{d}_s \gamma^\mu d_t)$
				$Q_{qd}^{(8)}$	$(\bar{q}_p \gamma_\mu T^A q_r)(\bar{d}_s \gamma^\mu T^A d_t)$

$(\bar{L}R)(\bar{R}L)$ and $(\bar{L}R)(\bar{L}R)$		B-violating	
Q_{ledq}	$(\bar{l}_p e_r)(\bar{d}_s q_t^c)$	Q_{duq}	$\varepsilon^{\alpha\beta\gamma} \varepsilon_{jk} [(d_p^\alpha)^T C u_r^\beta] [(q_s^\gamma)^T C l_t^k]$
$Q_{qqqd}^{(1)}$	$(\bar{q}_p^i u_r) \varepsilon_{jk} (\bar{q}_s^k d_t)$	Q_{quq}	$\varepsilon^{\alpha\beta\gamma} \varepsilon_{jk} [(q_p^\alpha)^T C q_r^{\beta k}] [(u_s^\gamma)^T C e_t]$
$Q_{qqqd}^{(8)}$	$(\bar{q}_p^i T^A u_r) \varepsilon_{jk} (\bar{q}_s^k T^A d_t)$	$Q_{quq}^{(1)}$	$\varepsilon^{\alpha\beta\gamma} \varepsilon_{jk} \varepsilon_{mn} [(q_p^\alpha)^T C q_r^{\beta k}] [(q_s^\gamma)^T C l_t^m]$
$Q_{lquq}^{(1)}$	$(\bar{l}_p e_r) \varepsilon_{jk} (\bar{q}_s^k u_t)$	$Q_{quq}^{(3)}$	$\varepsilon^{\alpha\beta\gamma} (\tau^I \varepsilon)_{jk} (\tau^I \varepsilon)_{mn} [(q_p^\alpha)^T C q_r^{\beta k}] [(u_s^\gamma)^T C l_t^m]$
$Q_{lquq}^{(3)}$	$(\bar{l}_p \sigma_{\mu\nu} e_r) \varepsilon_{jk} (\bar{q}_s^k \sigma^{\mu\nu} u_t)$	Q_{duu}	$\varepsilon^{\alpha\beta\gamma} [(d_p^\alpha)^T C u_r^\beta] [(u_s^\gamma)^T C e_t]$

Warsaw Basis: Purely Fermionic Operators

At lowest order in VBF, these operators do not contribute. At NLO they should be taken on account during the renormalization process



Warsaw Basis: Bosonic Operators

X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	Q_φ	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$

Implementing EFT in Vector Boson Fusion

What is LO EFT? is it more relevant than NLO-SM? and NLO-EFT?

$$\mathcal{M} = \mathcal{M}_{SM}^{\text{LO}} + \underbrace{\mathcal{M}_{SM}^{\text{NLO}}}_{\substack{\text{all available} \\ \text{QCD, EW corrections}}} + \underbrace{\mathcal{M}_{\text{dim}=6}^{\text{LO}} + \mathcal{M}_{\text{dim}=6}^{\text{NLO}}}_{\substack{\kappa\text{-framework} \\ \text{and dim 6 EFT}}}$$

Implementing EFT in Vector Boson Fusion

What is LO EFT? is it more relevant than NLO-SM? and NLO-EFT?

$$\mathcal{M} = \mathcal{M}_{SM}^{LO} + \underbrace{\mathcal{M}_{SM}^{NLO}}_{\substack{\text{all available} \\ \text{QCD, EW corrections}}} + \underbrace{\mathcal{M}_{dim=6}^{LO} + \mathcal{M}_{dim=6}^{NLO}}_{\substack{\kappa\text{-framework} \\ \text{and dim 6 EFT}}}$$

The answer depends on the scale of new physics Λ , the only way to be sure is to calculate the contributions and include them in the analysis.


Vector Boson Fusion Lowest Order EFT

Amplitude with dim-6 insertions:


$$\mathcal{A} = \sum_{n=N}^{\infty} g^n \mathcal{A}_n^{(4)} + g_6 \sum_{n=N_6}^{\infty} \sum_{l=0}^n g^n g_6^l \mathcal{A}_{nl}^{(6)}, \quad \text{with : } g_6 = \frac{1}{\sqrt{2}G_F\Lambda^2}.$$

VBF Signal


At lowest order we don't have to consider loops originated by dimension 6 operators. Therefore only corrections to the actual SM vertices



Feynman Diagrams



Vector Boson Fusion: Results



Vertices we have to calculate:

$\bar{u}uZ$, $\bar{d}dZ$, hZZ , hW^+W^- , $W^-u\bar{d}$, $W^+d\bar{u}$

Life is easy at LO ...

Bosonic dim-6 operators

Triple and quartic
gauge couplings

X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_\mu^A G_\nu^B G_\rho^C$	Q_φ	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_\mu^A G_\nu^B G_\rho^C$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\epsilon^{IJK} W_\mu^I W_\nu^J W_\rho^K$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\epsilon^{IJK} \tilde{W}_\mu^I W_\nu^J W_\rho^K$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{e\varphi}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$

Bosonic dim-6 operators

Purely Higgs

Triple and quartic gauge couplings

X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	Q_φ	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_{\mu\nu}^A G_\rho^{B\mu} G_\rho^{C\nu}$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_{\mu\nu}^I W_\nu^{J\rho} W_\rho^{K\mu}$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r)^\dagger \varphi W_{\mu\nu}^I$	$Q_{\varphi d}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi d}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\varphi^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$



Bosonic dim-6 operators

Purely Higgs

Triple and quartic gauge couplings

X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_\mu^A G_\nu^B G_\rho^C$	Q_{φ^6}	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_r e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_\mu^A G_\nu^B G_\rho^C$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\epsilon^{IJK} W_\mu^I W_\nu^J W_\rho^K$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\epsilon^{IJK} \tilde{W}_\mu^I W_\nu^J W_\rho^K$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_\mu^A G^{A\mu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_\mu^A G^{A\mu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_\mu^I W^{I\mu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_\mu^I W^{I\mu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_\mu^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_\mu^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$

Leptonic vertices



Bosonic dim-6 operators

Purely Higgs

Triple and quartic
gauge couplings

	X^3	φ^6 and $\varphi^4 D^2$	$\psi^2 \varphi^3$
Q_G	$f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	$Q_\varphi (\varphi^\dagger \varphi)^3$	$Q_{e\varphi} (\varphi^\dagger \varphi) (\bar{l}_p e_p \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_{\mu\nu}^A G_\rho^{B\mu} G_\rho^{C\nu}$	$Q_{\varphi\Box} (\varphi^\dagger \varphi) \Box (\varphi^\dagger \varphi)$	$Q_{u\varphi} (\varphi^\dagger \varphi) (\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\epsilon^{IJK} W_\mu^I W_\nu^J W_\rho^{K\mu}$	$Q_{\varphi D} (\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi} (\varphi^\dagger \varphi) (\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\epsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$		
	$X^2 \varphi^2$	$\psi^2 X \varphi$	$\psi^2 \varphi^2 D$
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_\mu^A G^{A\mu}$	$Q_{eW} (\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)} (\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi) (\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	$Q_{eB} (\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)} (\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi) (\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_\mu^I W^{I\mu}$	$Q_{uG} (\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e} (\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi) (\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_\mu^I W^{I\mu}$	$Q_{uW} (\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)} (\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi) (\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	$Q_{uB} (\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)} (\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi) (\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	$Q_{dG} (\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u} (\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi) (\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_\mu^I B^{\mu\nu}$	$Q_{dW} (\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d} (\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi) (\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_\mu^I B^{\mu\nu}$	$Q_{dB} (\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud} i (\tilde{\varphi}^\dagger D_\mu \varphi) (\bar{u}_p \gamma^\mu d_r)$

Gluon vertices

Leptonic
vertices

Bosonic dim-6 operators

Purely Higgs

Triple and quartic gauge couplings

	X^3	φ^6 and $\varphi^4 D^2$	$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^C$	Q_φ	$(\varphi^\dagger \varphi)^3$	
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}^{A\nu} G_\nu^{B\rho} G_\rho^C$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	
Q_W	$\varepsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$		$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\tilde{t}_R e_\mu \varphi)$
	$X^2 \varphi^2$	$\psi^2 X \varphi$	$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_\mu^A G^{A\mu}$	Q_{eW}	$(\tilde{t}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_\mu^A G^{A\mu}$	Q_{eB}	$(\tilde{t}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_\mu^I W^{I\mu}$	Q_{uG}	$(\tilde{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_\mu^I W^{I\mu}$	Q_{uW}	$(\tilde{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\tilde{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\tilde{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_\mu^I B^{\mu\nu}$	Q_{dW}	$(\tilde{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_\mu^I B^{\mu\nu}$	Q_{dB}	$(\tilde{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	
			$Q_{\varphi d}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi) (\tilde{t}_R \gamma^\mu t_r)$
			$Q_{\varphi d}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^2 \varphi) (\tilde{t}_R \tau^I \gamma^\mu t_r)$
			$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi) (\tilde{e}_R \gamma^\mu e_r)$
			$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi) (\tilde{q}_p \gamma^\mu q_r)$
			$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^2 \varphi) (\tilde{q}_p \tau^I \gamma^\mu q_r)$
			$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi) (\tilde{u}_p \gamma^\mu u_r)$
			$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi) (\tilde{d}_p \gamma^\mu d_r)$
			$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi) (\tilde{u}_p \gamma^\mu d_r)$

Gluon vertices

Leptonic vertices

CP-odd

Only 17 of the bosonic operators contribute to the signal vertices

X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_\mu^{AV} G_\nu^{B\rho} G_\rho^{C\mu}$	Q_φ	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_\mu^{AV} G_\nu^{B\rho} G_\rho^{C\mu}$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{JK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$



This is not the end of the story:

Dim-6 operators also induce corrections to kinetic terms and not only to vertices. If we want to be consistent we have to restore the canonical normalization of the \mathcal{L}_{SM} in the first place.

Even at Lowest Order ...

One example: $(\phi^\dagger D^\mu \phi)^*(\phi^\dagger D_\mu \phi)$

Choose one paradigmatic operator from the Warsaw basis: $(\phi^\dagger D^\mu \phi)^*(\phi^\dagger D_\mu \phi)$. Expand it in terms of physical fields:

$$\phi = \frac{1}{\sqrt{2}} \begin{pmatrix} h + \sqrt{2}F \\ 0 \end{pmatrix}, \quad D^\mu = \partial_\mu + \frac{ig}{2} B_\mu^a \tau^a + \frac{ig'}{2} B_\mu^0 \tau^0$$


$$(\phi^\dagger D_\mu \phi)^2 = \frac{1}{4} \left((h + \sqrt{2}F)(\partial_\mu h + i \left(\frac{gB_\mu^3 + g'B_\mu^0}{2} \right) (h + \sqrt{2}F)) \right)^2$$

- ▶ Corrections to the h propagator: $\frac{F^2(\partial_\mu h)^2}{2}$
- ▶ Corrections to the Z propagator: $\frac{F^4}{4} (g^2 \cos^2 \theta + g'^2 \sin^2 \theta + 2gg' \sin \cos \theta) Z^2$
- ▶ Correction to the hZZ vertex: $2\sqrt{2}F^3 (g^2 \cos^2 \theta + g'^2 \sin^2 \theta + 2gg' \sin \cos \theta) Z^2 h$

Some words on NLO and renormalization

At NLO, the procedure is the same, with two added difficulties:

1. Need to renormalize the new \mathcal{L}_{eff} → *The SM renormalization is not recyclable.*
2. Many new contributions will appear, through non-SM loops:



Ongoing work ... More in future presentations

Final words: Next steps

Now that the Higgs EFT and SMEFT have been formally developed and the first beams of RUN-2 are circulating, it's time to design the experimental analyses.

Focusing on two tasks:

1. Implementation of dim-6 operators in the MC generators and analysis of cross-sections for \mathcal{L}_{eff} (currently addressed in MadGraph)
2. Definition of pseudo-observables (PO's) to be measured: partial decay widths, effective couplings that might hint at New Physics

A little taste of PO's




Figure : $p_T(j_1)$ Vs. $p_T(j_2)$. Left plot: SM. Right plot: Prediction for $\epsilon_{WdL} = 0.05$

A little taste of PO's




Figure : M_Z Vs. $p_T(Z)$. Left plot: SM. Right plot: Prediction for $\kappa_{ZZ} = 1$ and $\epsilon_{ZuL} = 0.1$.

Preliminary plots from HXSWG WG2

Thank You!

... Any Questions?

