

ATTRACT TWD Symposium: Trends, Wishes and Dreams in Detection and Imaging Technologies



Contribution ID: 63

Type: **not specified**

STAX. Axion-like particle searches with sub-THz photons

We discuss an improved detection scheme for a light-shining-through-wall (LSW) experiment for axion-like particle searches [Capparelli, Cavoto, Ferretti, Giazotto, Polosa and Spagnolo, arXiv: 1510.06892]. We propose to use: extremely intense photon fluxes (from 100 kW to 1MW) from gyrotron sources at frequencies around 30 GHz; single photon detectors in this frequency domain, with efficiency ≈ 1 , based on transition-edge-sensors (TES); high quality factor Fabry-Perot cavities in the microwave domain ($Q \approx 10^4 - 10^5$), both on the photon-axion conversion and photon regeneration sides.

We compute that present laboratory exclusion limits on axion-like particles might be improved by at least four orders of magnitude for axion masses ≤ 0.01 meV.

Summary

Authors: POLOSA, Antonio (Universita' La Sapienza, Roma - Italy); Dr GIAZOTTO, Francesco (NEST, Istituto Nanoscienze-CNR and Scuola Normale Superiore); CAVOTO, Gianluca (Universita e INFN, Roma I (IT)); FERRETTI, Jacopo (INFN - National Institute for Nuclear Physics); Dr CAPPARELLI, Ludovico (UCLA); SPAGNOLO, Paolo (Universita di Pisa & INFN (IT))

Presenters: POLOSA, Antonio (Universita' La Sapienza, Roma - Italy); Dr GIAZOTTO, Francesco (NEST, Istituto Nanoscienze-CNR and Scuola Normale Superiore); CAVOTO, Gianluca (Universita e INFN, Roma I (IT)); FERRETTI, Jacopo (INFN - National Institute for Nuclear Physics); Dr CAPPARELLI, Ludovico (UCLA); SPAGNOLO, Paolo (Universita di Pisa & INFN (IT))