

# Overview of the CERN Accelerator Complex

CERN Lab 1954

1952: Geneva selected by the provisional Council as site for CERN  
1953: approved by referendum in Canton Genève  
1954: the first shovel of earth was dug on the Meyrin site

JULY 2016 - 10 January 2015

**Reyes Alemany, Beams Department, CERN**



# CERN Lab 2016

JURA

CMS

ALICE

SPS  
ATLAS


LHCb

BOOSTER  
PS

GENEVA

LEMANS





IR3

LHC

IR7

ALICE

SPS

LHC

LHCb

ATLAS

BOOSTER

PS



ALICE

SPS

ATLAS

BOOSTER

PS





LHC

SPS

ATLAS

ISR

AD

ENTRANCE B

BOOSTER

IT

PS

MAIN

RESTO 2

LINAC II

LEIR

CTF3

LINAC III









# The Proton Beam Starts Here ...

- The source cage houses the HV platform at 90 kV.







# Duoplasmatron Proton Source



Protons (at 90 keV) are produced by creating a plasma using  $H_2$  which is charged due to interaction with free electrons from the cathode. The plasma is then accelerated and becomes an ion beam.



# Duoplasmatron Proton Source







Protons (at 90 keV) are produced by creating a plasma using H<sub>2</sub> which is charged due to interaction with free electrons from the cathode. The plasma is then accelerated and becomes an ion beam.




# Radio Frequency Quadrupole

- RFQ is a linear accelerator that **FOCUSES, BUNCHES & ACCELERATES** with **HIGH EFFICIENCY** (90% w.r.t. 50% of conventional accelerators) and **PRESERVES THE EMITTANCE**
- The whole beam dynamics depends upon the shape of the vane tips






# (Phase space and emittance)



Analysis of  $x=f(t) \rightarrow$  provides information about the **path** taken by the system **BUT NOT** about the **energy**.

Analysis of  $v=f(t) \rightarrow$  provides information about the **energy** of the system **BUT NOT** about the **trajectory** taken.

... **Let's be inventive and try to analyse the evolution of the velocity as a function of position  $v=f(x)$**




A beam of charged particles in an accelerator subjected to **focusing and defocusing forces** have the same dynamics as the system above. The beam dynamics also **reproduces the same ellipse in phase space**

...

# (Phase space and emittance)

All particles with the same initial betatron amplitude (equivalent to  $x$ ) at a given position in the accelerator (or time) but different phases or momentum due to momentum spread (equivalent to  $v$ ), describe the same ellipse turn after turn




Along a beam line, the orientation and aspect ratio of the ellipse varies, **BUT THE AREA** remains **CONSTANT** in the absence of non-linear forces or acceleration

**AREA  $\approx$  EMITTANCE ( $\mathcal{E}$ )**


**Beam size  $\rightarrow \sigma = \sqrt{\mathcal{E}\beta}$  (in places without dispersion)**



# Linac 2




**DTL (Alvarez structure 1945)**



**Drift tubes and spacing become larger as the energy increases**  
**Focusing quads inside drift tubes**


**Pulse of beam ~100 μs long every 1.2 s**



$$V_A < V_B \rightarrow L_B > L_A$$

$$\rightarrow L = vT_{rf} = \beta\lambda_0$$

# PS Booster






$C = 154 \text{ m}$   
Commissioned in 1972

- **Synchrotron with 4 vertically stacked rings, each  $\frac{1}{4}$  of PS Circumference**
- **Duty cycle 1.2 s  $\rightarrow$  two cycles needed to fill the PS with protons for LHC**






# PS Booster: $E_{inj}=50\text{MeV}$ , $C=154\text{ m}$

Pulse from LINAC2 =  $100\ \mu\text{s}$



# PS Booster: $E_{inj}=50\text{MeV}$ , $C=154\text{ m}$




Pulse from LINAC2 =  $100\ \mu\text{s}$









# PS Booster: $E_{inj}=50\text{MeV}$ , $C=154\text{ m}$

Pulse from LINAC2 =  $100\ \mu\text{s}$



# PS Booster: $E_{inj}=50\text{MeV}$ , $C=154\text{ m}$


Pulse from LINAC2 =  $100\ \mu\text{s}$





- The bigger the number of turns the more intensity we can accumulate
- The problem is that the longer the injection takes, the more time the particles have to fill the whole available phase space + SPACE CHARGE  $\rightarrow$  emittance increases  $\rightarrow$  beam size increases
- **The Booster is the machine in the LHC Injector Chain where the transverse brightness of the LHC beam is determined**



# (Space Charge in One Slide)



+




$$\beta = \frac{v}{c}$$

Particles in the beam feel a strong repulsive force  $\rightarrow$


**change in tune**

# Proton Synchrotron (PS)



Combined-function magnets (QFQDB)

oldest functioning machine at CERN  
the first Alternating Gradient Machine!




1970-1976



GARGAMELLE

First evidence of weak neutral currents ( $Z^0$ )






# Proton Synchrotron (PS)


BOOSTER (1.4 GeV) → PS (26 GeV) → SPS (450 GeV) → LHC

BOOSTER (4 rings)



$h=1$

Two injections from  
BOOSTER to PS  
(2 x 1.2 s)




$h=7$  (6 buckets filled + 1 empty)

All operational beams cross **transition**  
(Transition energy 6.1 GeV)

JUAS 2016 26 January 2016

# Proton Synchrotron (PS)

Longitudinal bunch splitting → Reduce voltage on principal RF harmonic and simultaneously rise voltage on multiple RF harmonic → several type of RF cavities needed




**BOOSTER**

Nominal 25 ns beam production 26 January 2016



# Super Proton Synchrotron (SPS)


North area



Sp $\bar{p}$ S




- has probed the inner structure of protons
- investigated matter antimatter asymmetry
- searched for exotic forms of matter





# Large Hadron Collider (LHC)






# Large Hadron Collider (LHC)

## Geometry of the main dipoles (Total of 1232 cryodipoles)




LHC DIPOLE : STANDARD CROSS-SECTION

13894 AL-CD-AM - PB 137 - 9/94 DPH



Heat exchanger

Beam pipe (Ultrahigh beam vacuum  $10^{-10}$  Torr like at 1000 km over sea)



Cold bore non-magnetic austenitic steel

Beam Screen (Stainless Steel + Cu)

36.9 mm  
46.5 mm

Collars


He Vessel

Thermal shield

Iron yoke

Vacuum vessel ( $10^{-6}$  mbar)

Superconducting coils





L ~ 15 m  
8.3 T, 11.87 kA  
T = 1.9 K, ~27.5 ton



# Large Hadron Collider (LHC)

LHC arc cells = FoDo lattice\* with  
 ~ 90° phase advance per cell in the V & H plane



- MB:** main dipole
- MQ:** main quadrupole
- MQT:** Trim quadrupole
- MQS:** Skew trim quadrupole
- MO:** Lattice octupole (Landau damping)
- MSCB:** Skew sextupole + Orbit corrector (lattice chroma+orbit)
- MCS:** Spool piece sextupole
- MCDO:** Spool piece octupole + Decapole
- BPM:** Beam position monitor


## The FoDo-Lattice

A magnet structure consisting of focusing and defocusing quadrupole lenses in alternating order with **nothing** in between.  
 (Nothing = elements that can be neglected on first sight: drift, bending magnets, RF structures ... **and especially experiments...**)

# Linac4 : Replacing Linac2

## Linac4 : Approved in 2007 as a replacement to Linac2

- Energy 160 MeV (cf 50 MeV in Linac2) Doubles the space charge tune shift limit at injection into the PS Booster
- H- Injection : CERN is one of the few labs still using p<sup>+</sup>
- Connection to PSB LS2 (~ 2019)



Delivers 40 mA, 400 μs pulses at 2 Hz



50 MeV → 160 MeV

0.31\*1.12=0.35 → 0.52\*1.37=0.70

$$\Delta Q_{\text{LINAC4}} \approx 0.5 \Delta Q_{\text{LINAC2}}$$


$$\Delta Q_{SC} \propto \frac{N_b}{\epsilon_{X,Y}} \cdot \frac{R}{\beta\gamma^2}$$

with  $N_b$  : number of protons/bunch


$\epsilon_{X,Y}$  : norm. transverse emittances

$R$  : mean radius of the accelerator



# H<sup>-</sup> Injection







# H<sup>-</sup> Injection



Injection in the same phase space region!!!  
Emittance better preserved  
Not possible with LINAC2

The most important plus! → since we can afford a SPACE CHARGE  $\Delta Q_{50\text{MeV}}$  →



But  $\Delta Q_{\text{LINAC4}(160\text{MeV})} \approx 0.5 \Delta Q_{\text{LINAC2}(50\text{MeV})}$

$$\Delta Q_{SC} \propto \frac{N_b}{\epsilon_{X,Y}} \cdot \frac{R}{\beta\gamma^2}$$

$N_b^{\text{LINAC4}} \approx 2 N_b^{\text{LINAC2}}!!!!$

# Ion Chain

Detlef Küchler



Small sliver of solid isotopically pure  $^{208}\text{Pb}$  is placed in a ceramic crucible that sits in an "oven"


The metal is heated to around  $800^\circ\text{C}$  and ionized to become plasma. Ions are then extracted from the plasma and accelerated up to  $2.5 \text{ keV/nucleon}$ .



The source can also be set up to deliver other species... Ar and Xe being prepared for the SPS Physics programme



# Linac 3



Stripping foil Pb<sub>29</sub><sup>+</sup> → Pb<sub>54</sub><sup>+</sup>  
Stripping Efficiency is 20%

Interdigital-H (IH) linac  
4.2 MeV/nucleon


RFQ

Pb<sub>29</sub><sup>+</sup> 2.5 keV/nucleon

Spectrometer to select Pb<sub>29</sub><sup>+</sup>



# Ion Chain : Low Energy Ion Ring (LEIR)



LEIR Accumulates the 200 ms pulses from Linac3 into 2 bunches  
 Electron Cooling is used to achieve the required brightness  
 Acceleration to 72 MeV/nucleon before transfer to the PS  
 LEIR Cycle is 3.6 s

The Pb54+ is finally fully stripped to Pb82+ in the transfer line from PS to SPS


ANTIMATTER'S  
GONE MISSING ...

WHEN DID THIS  
HAPPEN, SIR?

ABOUT 15  
BILLION YEARS  
AGO ...



# History of the Antiproton Decelerator Chain












# Antiproton Decelerator : AD

Built in 1999 (from the old AC)  
 26 GeV/c PS Proton beam produces  $\bar{p}$   
 (1 in  $10^7$ ) which are focused and  
 captured in the AD and decelerated to  
 100 MeV/c (5.3 MeV)



# AD Layout




**Electron Cooling** **Experimental Area**

- ASACUSA**  
Antiprotonic helium  $\rightarrow m_{\bar{p}}$
- ALPHA**  
2011  $\bar{H}$  trapped for 16'
- AEGIS**  
1<sup>st</sup> meas. of gravitational effect on  $\bar{H}$
- GBAR<sup>(1)</sup>**  
Gravitational effect on  $\bar{H}$
- ATRAP**  
2002 first glimpse inside  $\bar{H}$
- BASE**  
 $\bar{p}$  magnetic moment

(1) In construction

# Elena ... More Deceleration


Today's set-up



ELENA will overcome this problem + will be able to deliver beams almost simultaneously to all four experiments resulting in an essential gain in total beam time for each experiment. This also opens up the possibility to accommodate an extra experimental zone.

**Under Construction**

**A second stage of deceleration after  
AD Momentum: 100 – 13.7 MeV/c  
Kinetic : 5.3 – 0.1 MeV**






**Commissioning in 2016  
Operation 2017**



ISOLDE SC in 1967 (until 1990)  
 ISOLDE PSB in 1992

# PSB Experimental Areas: ISOLDE



## Next generation of nuclear physics:

**HIE-ISOLDE (+SC RF):**  $E_{kin} \leq 10 \text{ MeV/n} \rightarrow A \leq 200$   
**PHASE 1a:** 2015  $\rightarrow 4.8 \text{ MeV/u}$  **PHASE 1b:** 2016  $\rightarrow 5.5 \text{ MeV/u}$   
**PHASE 2:** 2017?  $\rightarrow 10 \text{ MeV/u}$  **PHASE 3** (+ chooper)

GPS: Global Purpose Separator  
 HRS: High Resolution Separator  
 HIE-ISOLDE: High Intensity and Energy ISOLDE

# PS Experimental Areas: East Hall




**Study** the influence of galactic cosmic rays on the **Earth's climate** through the media of aerosols and clouds

Detector Calibration

CLIMATE

Proton & neutron irradiation facilities



$p^+$  (26 GeV)

Bdg 157



SATELLITE EXPERIMENTS

**Secondary Beams:**  
 Momentum range 1-15 GeV/c  
 Electrons, Hadrons & Muons  
 Max  $1-2E^+6$  particles per spill

Resonant Extraction with a spill length of  $\sim 400$ ms

East Hall

de

PS

CTF3

n-ToF

LEIR

Linac2


JUA



# PS Experimental Areas: n-TOF




Study of neutron-induced reactions



- Transmutation of nuclear waste
- Stellar Nucleosynthesis
- Symmetry Breaking in compound nuclei

**Each primary proton produces ~300 neutrons**  
**Neutron → meV - GeV**


The neutron kinetic energy is determined by **time-of-flight**






# SPS Experimental Areas: North Area

**CALET: Calorimetric Electron Telescope**



High energy astroparticle physics of the International Space Station

**NA61/SHINE (QCD experiment)**



**Study of hadron structure and hadron spectroscopy** with high intensity muon and hadron beams

- 7 beam lines (tot:5.8 km)
- 3 experimental halls
- ~ 2000 scientist/year
- Slow extraction
- 3 primary targets
- Ion physics program: (Be, Ar, Xe)
- ~ 50 different clients/year

**NEUTRON**




Russian regular satellite Clarify the Cosmic Rays origin


**North Experimental Area**



**Physics Beyond the Standard Model**



**HiRadMat**



**SPS**

**Awake (ex CNGS)**

JUAS 2016



COMPASS: Common Muon and Proton Apparatus for Structure and Spectroscopy

# SPS Experimental Areas: *AWAKE*

Proof-of-principle:

- Inject 10-20 MeV electron beam
- acceleration of electrons to **multi-GeV energy range** in the wakefield driven by protons.


→ first proton driven PWA experiment world-wide



# SPS Experimental Areas:


Current and Future Accelerators operate with higher energy, higher intensity, smaller size beams.

LHC nominal beam (2808 bunches with 1.5  $10^{11}$  p+/b at 7 TeV) energy = **362 MJ/beam**  
→ energy equivalent to



**HiRadMat is a facility designed, to study the impact of intense pulsed beam on materials**

- Thermal management
- Radiation Damage to materials
- Thermal shock – beam induced pressure waves





# SPS Experimental Areas:


Current and Future Accelerators operate with higher energy, higher intensity, smaller size beams.

LHC nominal beam (2808 bunches with 1.5 10<sup>11</sup> p<sup>+</sup>/b at 7 TeV) energy = **362 MJ/beam**  
 → energy equivalent to




Elkin (@155 km/h) ≈ 360 MJ

Simulation: 8 LHC bunches @5 TeV impacting a Tungsten collimator jaw



HiRadMat is a facility designed, to study the impact of intense pulsed beam on materials

- Thermal management
- Radiation Damage to materials
- Thermal shock – beam induced pressure waves



# *Spare Slides*

# Further Reading

**The LHC Design Report Volume 1: The LHC Main Ring**, CERN-2004-003-V-1,  
<http://cds.cern.ch/record/782076/files/CERN-2004-003-V1.pdf>

**The LHC Design Report Volume 1: The LHC Infrastructure and Services**, CERN-2004-003-V-2,  
<http://cds.cern.ch/record/782076/files/CERN-2004-003-V2.pdf>

**The LHC Design Report Volume 3: The LHC Injector Chain** : CERN-2004-003-V-3:  
<http://cds.cern.ch/record/823808/files/CERN-2004-003-V3.pdf>

**Fifty years of the CERN Proton Synchrotron: Volume 1** :CERN-2011-004,  
<http://cds.cern.ch/record/1359959/files/cern-2011-004.pdf>

**Fifty years of the CERN Proton Synchrotron: Volume 2** :CERN-2013-005,  
<http://cds.cern.ch/record/1597087/files/CERN-2013-005.pdf>

**Linac4 Technical Design Report::**  
<http://cds.cern.ch/record/1004186/files/ab-2006-084.pdf>

**Elena Conceptual Design Report:**  
<http://cds.cern.ch/record/1309538/files/CERN-BE-2010-029.pdf>

**AWAKE Technical Design Report:**  
<http://cds.cern.ch/record/1537318/files/SPSC-TDR-003.pdf>


**HiRadMat:**  
<http://cds.cern.ch/record/1403043/files/CERN-ATS-2011-232.pdf>



# Generating a 25ns Bunch Train in the PS



- **Longitudinal bunch splitting (basic principle)**

- Reduce voltage on principal RF harmonic and simultaneously rise voltage on multiple harmonics (adiabatically with correct phase, etc.)



**Use double splitting at 25 GeV to generate 50ns bunch trains instead**

# Proton Synchrotron (PS)




The PS is the machine in the LHC Injector Chain where the Longitudinal characteristics of the LHC beam are determined

# Large Hadron Collider (LHC)

Golden formula (you should know by heart)


$$B\rho = \frac{p}{Ze}$$



$p$  = nucleon momentum → defined by the physics case → TeV range → **7 TeV**

$$B = \frac{p}{\rho Ze} \approx 3.33 \frac{p \left( \frac{\text{GeV}}{c} \right)}{\rho(\text{m})} = 8.39 \text{ T}$$

We need SUPERCONDUCTING technology







# Large Hadron Collider (LHC)

Production rate of events is determined by the cross section  $\Sigma_{\text{react}}$  and a parameter L that is given by the design of the accelerator:  
 ... the luminosity

$$R = L * \Sigma_{\text{react}} \approx 25 \frac{1}{10^{-15} b} 10^{-12} b = \text{some } 1000H$$



remember:  
 $1b = 10^{-24} \text{ cm}^2$



Integrated luminosity during RUN I

$$\int L dt \approx 25 \text{ fb}^{-1}$$

**Official number: 1400 clearly identified Higgs particles “on-tape”**

# Overall Protons Delivered in 2012

Facility	Protons Delivered	% of Total
Isolde	$1.15 \times 10^{20}$	63.8%
CNGS	$3.9 \times 10^{19}$	21.6%
n-TOF	$1.9 \times 10^{19}$	10.2%
The rest	$8.13 \times 10^{18}$	4.5%
<b>LHC</b>	<b><math>3.25 \times 10^{16}</math></b>	<b>0.018%</b>
<b>Total</b>	<b><math>1.81 \times 10^{20}</math></b>	


## Colliders are very Efficient!

**The LHC Physics Program Used 0.018% of the protons produced in CERN accelerators during 2012!**


- ❖ Intensities as delivered to the facility, upstream losses ignored,
- ❖ Beams for Machine Setup and Studies Excluded
- ❖ The total delivered protons represents roughly 0.27mg (rest mass!)

# Large Hadron Collider (LHC)

## Superconducting cables of Nb-Ti



LHC ~ 27 km circumf. with 20 km of superconducting magnets operating @8.3 T. An equivalent machine with normal conducting magnets would have a circumference of 100 km and would consume 1000 MW of power → we would need a dedicated nuclear power station for such a machine. LHC consumes ~ 10% nuclear power station



## LHC Requires


- 90,000 T of liquid Nitrogen
- 130 T of Liquid Helium to keep it cold



**June 1994**  
first full scale prototype dipole

**June 2007** First sector cold

**ECFA-CERN workshop**



**April 2008**  
Last dipole down



**1994 project approved by council (1-in-2)**



SSC

25 y

cancelled


Main contracts signed



**First set of twin 1 m prototypes Over 9 T**



**2002 String 2**





**November 2006**  
1232 delivered



**September 10, 2008**  
First beams around

**August 2008**


First injection test




**Sept. 10, 2008**  
First beams around

**Repair and Consolidation**

**November 29, 2009**  
Beam back




**October 14, 2010**  
 $L = 1 \times 10^{32}$   
248 bunches




**October, 2011**  
 $3.5 \times 10^{33}$ ,  $5.7 \text{ fb}^{-1}$   
**First Hints!!**

**June 28 2011**  
1380 bunches  
**1380**




**May 2012**  
Ramping  
Performance



**Feb. 2013**  
p-Pb<sup>82+</sup>  
New Operation  
Mode

**March 14<sup>th</sup> 2012**  
Restart  
with Beam



**Nov. 2012**  
End of p<sup>+</sup> Run I

2008

2009

2010

2011


2012

2013

**Sept. 19, 2008**  
Disaster




**March 30, 2010**  
First collisions at 3.5 TeV



**November 2010**  
Pb<sup>82+</sup> Ions




**November 2011**  
Second Ion Run



**LSI**

# Filling the LHC (2012)

	25 ns (design)	50 ns (2012)	25 ns (2012)#
Energy per beam [TeV]	7	<b>4</b>	4
Intensity per bunch [ $\times 10^{11}$ ]	1.15	<b>1.7</b>	1.2
Norm. Emittance H&V [ $\mu\text{m}$ ]	3.75	<b>1.8</b>	2.7
Number of bunches	2808	<b>1380</b>	N.A.#
$\beta^*$ [m]	0.55	<b>0.6</b>	N.A.#
Peak luminosity [ $\text{cm}^{-2}\text{s}^{-1}$ ]	$1 \times 10^{34}$	<b><math>7.7 \times 10^{33}</math></b>	N.A.#





→ Each bunch from the Booster divided by 6 →  $6 \times 3 \times 2 \times 2 = 72$   
 The 25 ns PS production scheme (2012)



# The 25 ns was only used for scrubbing and tests in 2012



# CTF 3 – CLIC Test Facility





CLIC goal:



**Drive Beam** 100 A, 239 ns  
2.38 GeV  $\rightarrow$  240 MeV

**Main Beam** 1.2 A, 156 ns  
**9 GeV  $\rightarrow$  1.5 TeV**

# CTF 3 – CLIC Test Facility




# CTF 3 – CLIC Test Facility






# *High Light Of HEP -Year*



***ATLAS event display: Higgs => two electrons & two muons***







A visualization of the cosmic web, showing a complex network of filaments and clusters of galaxies. The background is dark, with numerous bright orange and red points representing galaxies. The filaments are formed by these points connected by thin lines, creating a web-like structure. The overall appearance is that of a vast, interconnected network of matter in the universe.

*Reconstruction of Dark Matter distribution based on observations*

*Budget: Dark Matter: 33 %  
Dark Energy: 66 %  
Anything else (including us) 1%*




# CERN Lab 20??

HE\_LHC:  
27 km 33 TeV 20T

VHE\_LHC:  
100 km 100 TeV

- LHC Tunnel
- - - VHE\_LHC (80 km)
- VHE\_LHC (100 km)



LHC

Jura

Geneva

Salève

© 2012 Google  
Image © 2012 GeoEye  
Map data © 2012 IGN France