

Double-parton scattering effects in double charm production within gluon fragmentation scenario

Rafał Maciuta

Institute of Nuclear Physics (PAN), Kraków, Poland

Low-x Meeting 2016,

6th - 11th June 2016, Gyöngyös, Hungary



Outline

- 1 Motivation
- 2 Theoretical framework
 - Factorized Ansatz for DPS
 - Basics of the k_T -factorization approach
- 3 Numerical results vs. LHCb data
 - Standard fragmentation approach
 - New scenario with gluon fragmentation
- 4 Summary

Based on:

Maciuła, Saleev, Shipilova, Szczurek, Phys. Lett. B758, 458 (2016)

Hameren, Maciuła, Szczurek, Phys. Lett. B748, 167 (2015)

Hameren, Maciuła, Szczurek, Phys. Rev. D89, 094019 (2014)

Maciuła, Szczurek, Phys. Rev. D87, 074039 (2013)


Łuszczak, Maciuła, Szczurek, Phys. Rev. D79, 094034 (2012)




Why double charm?

Double charm \Rightarrow final state with two pairs of $c\bar{c}$


Single-parton scattering (SPS)



Double-parton scattering (DPS)



SINGLE vs. DOUBLE CHARM



- SPS $c\bar{c} \gg$ SPS $c\bar{c}c\bar{c}$: the SPS double charm is negligible higher-order correction only
- SPS $c\bar{c}$ vs. DPS $c\bar{c}c\bar{c}$: **comparable total cross sections at LHC energies!**



Best known place to study DPS effects at the LHC


How the DPS effects in charm sector can be investigated experimentally?

Study of **CHARM MESON-MESON pair** production:

DD pairs - both mesons containing c quarks or both containing \bar{c} antiquark

- impossible to produce within standard SPS single $c\bar{c}$ mechanism
- SPS double charm expected to be very small
- measurements of charm meson-meson pairs highly recommended at the LHC
- same-sign nonphotonic lepton pairs, e.g. $\mu^+\mu^+$ at ALICE

First measurement by LHCb: *Observation of double charm production involving open charm in pp collisions at $\sqrt{s} = 7$ TeV*, J. High Energy Phys. 06, 141 (2012)




Mode	σ [nb]
$D^0 D^0$	$690 \pm 40 \pm 70$
$D^0 \bar{D}^0$	$6230 \pm 120 \pm 630$

- very large cross section: $\frac{\sigma(DD)}{\sigma(D\bar{D})} \approx 10\%$
- various DD modes
- several differential distributions



SPS double charm definitely negligible





SPS double charm calculated in two different ways:

- LO collinear approach
- **NEW**: k_T -factorization approach (part of higher-order corrections included)
- **first time**: off-shell initial state partons ($2 \rightarrow 4$)

Monte Carlo: **AVHLIB** (A. van Hameren):


<https://bitbucket.org/hameren/avhlib>



Can the LHCb data be explained by the DPS?



Double charm within double-parton scattering (DPS)



DPS in general form:

$$d\sigma^{DPS}(pp \rightarrow c\bar{c}c\bar{c}X) = \frac{1}{2} \cdot \Gamma_{gg}(b, x_1, x_2; \mu_1^2, \mu_2^2) \Gamma_{gg}(b, x'_1, x'_2; \mu_1^2, \mu_2^2) \times d\sigma_{gg \rightarrow c\bar{c}}(x_1, x'_2, \mu_1^2) \cdot d\sigma_{gg \rightarrow c\bar{c}}(x'_1, x_2, \mu_2^2) dx_1 dx_2 dx'_1 dx'_2 d^2 b$$

DPDF - emission of one parton with assumption that second parton is also emitted

$$\Gamma_{ij}(b, x_1, x_2; \mu_1^2, \mu_2^2) = F_i(x_1, \mu_1^2) F_j(x_2, \mu_2^2) F(b; x_1, x_2, \mu_1^2, \mu_2^2)$$

- longitudinal and transverse correlations between two partons
- spin, flavour and color correlations
- well established theory: e.g. Diehl, Ostermeier, Schafer, JHEP 03, 089 (2012) but not yet available for phenomenological studies

Factorized ansatz (pocket-formula)

In a simple probabilistic picture process initiated by:

two simultaneous hard gluon-gluon scatterings in one proton-proton interaction

$$\sigma^{DPS}(pp \rightarrow c\bar{c}c\bar{c}X) = \frac{1}{2\sigma_{eff}} \cdot \sigma^{SPS}(pp \rightarrow c\bar{c}X_1) \cdot \sigma^{SPS}(pp \rightarrow c\bar{c}X_2)$$

two subprocesses are not correlated and do not interfere


- $\sigma_{eff} \Rightarrow$ model parameter \Rightarrow normalization of σ^{DPS}



Double charm within double-parton scattering (DPS)

Factorized ansatz (pocket-formula)

- a good approximation for **small- x partons** \Rightarrow charm at high energies
- **color/flavour correlations suppressed** in evolution (Kasemets et al., Phys. Rev. D91, 014015 (2015))
- **DPDFs in multiplicative form:** $\Gamma_{gg}(b; x_1, x_2, \mu_1^2, \mu_2^2) = F_g(x_1, \mu_1^2)F_g(x_2, \mu_2^2)F(b)$
- only transverse correlations taken into account
- $\sigma_{eff} = \left[\int d^2b (F(b))^2 \right]^{-1}$, $F(b)$ - overlap of the matter distribution in transverse plane where b is a distance between both gluons
- nonperturbative quantity with dimension of cross section, connected to transverse size of proton




- extracted from several experimental analyses
- in principle may not be universal
- detailed studies: Seymour, Siódmok, JHEP 10, 113 (2013)
- **world average:** $\sigma_{eff} \approx 15$ mb (large uncertainties)
- **LHCb double charm data:** $\sigma_{eff} \approx 15 - 30$ mb
- e.g. $D^0 D^0$: $\sigma_{eff} \approx 21_{-6}^{+7}$ mb (arXiv:1308.6749 (hep-ph))

- For charm: **spin (polarization) correlations very small** (Echevarria et al. JHEP 04, 034 (2015))



Each step of DPS: two single-parton scatterings



k_T -factorization $\rightarrow \kappa_{1,t}, \kappa_{2,t} \neq 0$ e.g. Collins-Ellis, Nucl. Phys. B360 (1991) 3


\Rightarrow very efficient approach for $Q\bar{Q}$ correlations

- multi-differential cross section


$$\frac{d\sigma}{dy_1 dy_2 d^2 p_{1,t} d^2 p_{2,t}} = \sum_{i,j} \int \frac{d^2 \kappa_{1,t}}{\pi} \frac{d^2 \kappa_{2,t}}{\pi} \frac{1}{16\pi^2 (x_1 x_2 s)^2} \overline{|\mathcal{M}_{g^*g^* \rightarrow Q\bar{Q}}|^2} \times \delta^2(\bar{\kappa}_{1,t} + \bar{\kappa}_{2,t} - \bar{p}_{1,t} - \bar{p}_{2,t}) \mathcal{F}_i(x_1, \kappa_{1,t}^2) \mathcal{F}_j(x_2, \kappa_{2,t}^2)$$

- **LO off-shell** $|\mathcal{M}_{g^*g^* \rightarrow Q\bar{Q}}|^2$: e.g. Catani-Ciafaloni-Hautmann, Nucl. Phys. B366 (1991) 135
- $\mathcal{F}_i(x_1, \kappa_{1,t}^2), \mathcal{F}_j(x_2, \kappa_{2,t}^2)$ - unintegrated (k_T -dependent) gluon distributions
- some part of very important **higher-order corrections effectively included**


pair creation
with gluon emission




flavour excitation



gluon splitting





Unintegrated gluon distribution functions (UGDFs)



most popular models:

- Kwieciński, Jung (CCFM, wide range of x)
- Kimber-Martin-Ryskin (DGLAP-BFKL, wide range of x)
- Kwieciński-Martin-Staśto (BFKL-DGLAP, small x -values)
- Kutak-Staśto (BK, saturation, only small x -values)




Lesson from **inclusive single D meson and $D\bar{D}$ pair** production at the LHC:



- **KMR UGDF works very well** (single particle spectra and correlation observables)



First DPS results vs. LHCb double charm data



- DPS mechanism describes the LHCb data
- clear evidence of DPS effects
- factorized ansatz: flat $\Delta\phi$ distribution
- LHCb data: some $\Delta\phi$ correlations suggested



New scenario with gluon fragmentation

Different fragmentation schemes



Standard approach (firs results):

- Peterson et al.:
scale-independent FFs
- only one component:
 $c \rightarrow D$
- k_T -factorization: Maciula, Szczurek,
Phys.Rev.D87, 094022 (2013)

New scenario:

- KKKS08 (Kniehl et al.): several FFs of a parton
(gluon, $u, d, s, c, \bar{u}, \bar{d}, \bar{s}, \bar{c}$) $\rightarrow D$ mesons
- scale-dependent FFs undergo DGLAP evolution
- $c \rightarrow D$ reduced with increasing p_T (scale)
- very important contribution from $g \rightarrow D$
- k_T -factorization: Karpishkov et al. Phys.Rev.D91, 054009 (2015)

Both prescriptions describe the LHCb data for single D meson production




What are consequences of the new scenario for DPS double charm?




New scenario with gluon fragmentation

New scenario for double charm production


DPS: $cc \rightarrow DD$




DPS: $gg \rightarrow DD$



DPS: $cg \rightarrow DD$



SPS: $gg \rightarrow DD$ (NEW!)






all diagrams calculated in the k_t -factorization approach



New scenario with gluon fragmentation

New scenario results vs. LHCb double charm data



- LHCb data overestimated
- SPS $gg \rightarrow DD$ contribution very important
- $\Delta\phi$ correlation as in the LHCb data




Possible problems:

- massless DGLAP evolution?
- larger σ_{eff} ?
- small- x (nonlinear) or large- x effects?



New scenario with gluon fragmentation

New scenario results vs. LHCb double charm data



- very large σ_{eff} needed to describe the LHCb data
- rather unrealistic and probably unacceptable by the MPI community
- one should rather concentrate on mass effects in evolution or small/large- x effects



Conclusions

Production of double charm - DD meson-meson pairs - at the LHC
is dominated by the double-parton scattering (DPS) mechanism

Standard approach:

- very good description of the LHCb data with the DPS factorized ansatz and k_T -factorization
- some small $\Delta\varphi$ correlations visible in the LHCb data difficult to obtain within the factorized DPS model, even when including spin correlations between two partons in DPDFs

New scenario:

- $g \rightarrow D$ component which appears in the DGLAP evolution of FFs change the overall picture
- new mechanisms appear: DPS $gg \rightarrow DD$, DPS $cg \rightarrow DD$ and SPS $gg \rightarrow DD$
- the LHCb data overestimated
BUT: $\Delta\varphi$ correlations arise due to the presence of the new SPS $gg \rightarrow DD$ component
- large contribution from the SPS $gg \rightarrow DD$ makes the extraction of σ_{eff} from the LHCb data more complex (revision of the existing extraction needed)

Thank You for attention!

