



Contribution ID: 1052

Type: Oral (Non-Student) / orale (non-étudiant)

## Determination of global-scale diffusion coefficients in the stratosphere using a new model of local mixing

*Monday, 13 June 2016 10:45 (15 minutes)*

Due to the intermittency and spatial distribution of small layers of turbulence in the stratosphere, determination of large-scale diffusion coefficients is a non-trivial process. Previous models have assumed that all layers are perfectly mixed, but generally spatio-temporally disconnected. Our new model allows for partially mixed layers, especially for large and intense layers, which has profound effects on the global-scale diffusion coefficient. We also better represent the spatio-temporal distribution using a 2-dimensional model, as distinct from the more common one-dimensional model. Our newer model is used in combination with balloon and aircraft measurements of layer thicknesses and distributions to place limits on the possible values of large-scale diffusion in the stratosphere and upper troposphere. The values in the stratosphere are particularly well confined.

**Primary author:** HOCKING, Wayne (University of Western Ontario)

**Co-authors:** Dr TARASICK, David (Environment Canada); Dr OSMAN, Mohammed (Environment Canada)

**Presenter:** HOCKING, Wayne (University of Western Ontario)

**Session Classification:** M1-3 Theory, Modelling, and Forecasting I (DASP) / Théorie, modélisation et prévisions I (DPAE)

**Track Classification:** Atmospheric and Space Physics / Physique atmosphérique et de l'espace (DASP-DPAE)