



Contribution ID: 1215

Type: **Oral (Non-Student) / orale (non-étudiant)**

## **Pseudospin representation of the two-site Anderson-Hubbard model**

*Thursday 16 June 2016 09:00 (15 minutes)*

The state of an Anderson localized system can be described in terms of the occupation of a set of single-particle wave functions which are localized in space. When interactions are added, single-particle wave functions are no longer well defined, so what is a useful description of the state of a many-body localized system and what about it is localized? Given that any system with Hilbert-space dimension  $2^N$  may be described by an Ising-type Hamiltonian, it has been proposed that in a fully many-body localized system the Ising pseudospins in this representation may be chosen to be local. Actually constructing these spins is non-trivial. While a number of approaches have been proposed, few explicit examples exist and almost all work has been on spin systems. Here we present the Hamiltonian of a two-site Hubbard model with disorder and nearest-neighbor interactions written in terms of pseudospins, and we explore the form of these pseudospins and their evolution as a function of hopping amplitude.

**Primary author:** WORTIS, Rachel (Trent University)

**Co-author:** KENNETT, Malcolm (Simon Fraser University)

**Presenter:** WORTIS, Rachel (Trent University)

**Session Classification:** R1-2 Strongly Correlated Systems (DCMMP) / Systèmes fortement corrélés (DPMCM)

**Track Classification:** Condensed Matter and Materials Physics / Physique de la matière condensée et matériaux (DCMMP-DPMCM)