

Contribution ID: 1447 Type: Poster (Student, Not in Competition) / Affiche (Étudiant(e), pas dans la compétition)

Colossal Permittivity in In + Nb co-doped TiO2

Tuesday 14 June 2016 19:46 (2 minutes)

In + Nb co-doped TiO₂ has been previously investigated and shown to display colossal permittivity ($\epsilon_1 > 10^3$) for frequencies less than or equal to 1 MHz using impedance spectroscopy. In this work the lattice vibrational contribution to the low frequency dielectric function is studied in ceramic samples at the 0\%,5\%, and 10\% doping level using temperature-dependent reflectance spectroscopy. It is found that the soft mode behaviour exhibited by the lowest frequency infrared ibrational mode in pure TiO₂ is not strongly affected by the codoping. There is a large discrepancy between the ϵ_1 measured at the lowest far infrared frequencies in this reflectance spectroscopy study (≈ 1.5 THz) and the value measured below 1 MHz.

Primary authors: CRANDLES, David (Brock University); YEE, Susan

Presenter: YEE, Susan

Session Classification: DCMMP Poster Session with beer / Session d'affiches, avec bière DPMCM

Track Classification: Condensed Matter and Materials Physics / Physique de la matière condensée et matériaux (DCMMP-DPMCM)