

Higgs to di-photons at ATLAS

B. Mellado


University of the Witwatersrand

On behalf of the ATLAS Collaboration




Higgs Couplings 2016, SLAC, 11/09/16

Higgs Cross-Sections at LHC




Main Decay Modes




B_b	B_τ	B_μ	B_c	B_g
$0.577^{+3.2\%}_{-3.3\%}$	$0.0632^{+5.7\%}_{-5.7\%}$	$(2.2 \times 10^{-4})^{+6.0\%}_{-5.9\%}$	$0.0291^{+12.2\%}_{-12.2\%}$	$0.0857^{+10.2\%}_{-10.0\%}$
B_γ	$B_{Z\gamma}$	B_W	B_Z	Γ_H [MeV]
$(2.28 \times 10^{-3})^{+5.0\%}_{-4.8\%}$	$(2.28 \times 10^{-3})^{+9.0\%}_{-8.8\%}$	$+4.3\%_{-4.2\%}$	$2.64^{+4.3\%}_{-4.2\%}$	$4.07^{+4.0\%}_{-3.9\%}$


Higgs decay to $\gamma\gamma$



$\gamma\gamma$ Backgrounds



Reducible γj and jj Backgrounds



$q \rightarrow \pi^0$

Different background components studied with data-driven methods

Inner detector :

Pixel detector + SCT + TRT



$$\frac{\sigma_{PT}}{PT} \approx 0.05\% \rho_T \oplus 1\%, |\eta| < 2.5$$

EM calorimeter :

Lead-LAr sampling calo. with accordion geometry

$$\frac{\sigma_E}{E} \approx \frac{10\%}{\sqrt{E}} \oplus 0.7\%, |\eta| < 3.2$$

The ATLAS Detector



Hadronic calorimeter :

Steel and scintillating tiles in the barrel, copper and liquid argon in end-caps

$$\frac{\sigma_E}{E} = \frac{50\%}{\sqrt{E}} \oplus 3\%, |\eta| < 3.2$$

$$\frac{\sigma_E}{E} = \frac{100\%}{\sqrt{E}} \oplus 10\%, 3.1 < |\eta| < 4.9$$

Muon spectrometer :

superconducting air-core toroid magnets, gas based muon chambers

$$\frac{\sigma_{PT}}{PT} \approx 2\% \text{ at } 50\text{GeV to } 10\% \text{ at } 1\text{TeV}, |\eta| < 2.7$$

Object and Event Selection

□ **Photons: EM clusters with calo + track isolation that:**

$$|\eta_\gamma| < 1.37 \text{ or } 1.52 < |\eta_\gamma| < 2.37$$

$$p_T^{\gamma 1} / m_{\gamma\gamma} > 0.35 \text{ and } p_T^{\gamma 2} / m_{\gamma\gamma} > 0.25$$

□ **Leptons: Calo. + track isolation with:**

$$|\eta_e| < 2.47, E_{Te} > 15 \text{ GeV}, |\eta_\mu| < 2.7, E_{T\mu} > 10 \text{ GeV}$$

□ **Jets: Anti- k_t with R=0.4, overlap removal:**




$$|\eta_j| < 4.4, p_{Tj} > 30 \text{ GeV}$$


□ **B-jets with working points**

$$\epsilon_b(R_j) : 60\%(450), 70\%(140), 80\%(29), 85\%(11) \quad 6$$


Diphoton Invariant Mass

- Primary vertex reconstruction :
 - Photon calorimeter pointing
 - Use conversion tracks when available
- Energy scale calibration from Z to electrons applied
- Crystal Ball + Gaussian model with narrow widths (of the core of the distribution) :





Introducing categories to enhance sensitivity to explore various production mechanisms



Effective signal mass resolutions σ_{68} and σ_{90} for the 7 TeV and 8 TeV data in each event category, where σ_{68} (σ_{90}) is defined as half of the smallest interval expected to contain 68% (90%) of the signal events


Category	$\sqrt{s}=7$ TeV		$\sqrt{s}=8$ TeV	
	σ_{68} [GeV]	σ_{90} [GeV]	σ_{68} [GeV]	σ_{90} [GeV]
Central - low p_{Tt}	1.36	2.32	1.47	2.50
Central - high p_{Tt}	1.21	2.04	1.32	2.21
Forward - low p_{Tt}	1.69	3.03	1.86	3.31
Forward - high p_{Tt}	1.48	2.59	1.64	2.88
VBF loose	1.43	2.53	1.57	2.78
VBF tight	1.37	2.39	1.47	2.61
VH hadronic	1.35	2.32	1.45	2.57
VH E_T^{miss}	1.41	2.44	1.56	2.74
VH one-lepton	1.48	2.55	1.61	2.80
VH dilepton	1.45	2.59	1.59	2.76
$t\bar{t}H$ hadronic	1.39	2.37	1.53	2.64
$t\bar{t}H$ leptonic	1.42	2.45	1.56	2.69



$$\mu_i = \frac{\sigma_i}{(\sigma_i)_{\text{SM}}} \quad \text{and} \quad \mu^f = \frac{B^f}{(B^f)_{\text{SM}}}$$

$$\mu_i^f = \frac{\sigma_i \cdot B^f}{(\sigma_i)_{\text{SM}} \cdot (B^f)_{\text{SM}}} = \mu_i \cdot \mu^f$$

Coupling Modifiers

$$\sigma_i \cdot B^f = \frac{\sigma_i(\vec{k}) \cdot \Gamma^f(\vec{k})}{\Gamma_H}, \quad K_j^2 = \Gamma^j / \Gamma_{\text{SM}}^j$$






ATLAS

$H \rightarrow \gamma\gamma, m_H = 125.4 \text{ GeV}$

Total uncertainty

■ $\pm 1\sigma$ ■ $\pm 2\sigma$



Systematics

Relative systematic uncertainties on the inclusive yields. Parentheses are for $t\bar{t}H$ and VH mechanisms

Uncertainty source	$\sqrt{s} = 7$ TeV	$\sqrt{s} = 8$ TeV
Luminosity	1.8	2.8
Trigger	0.2	0.2
Photon identification	8.4(9.3)	1.0(4.1)
Isolation	1.3–2.3(3.8)	1.3–2.3(3.8)


Systematic uncertainties on the diphoton mass resolution for the 8-TeV data. Used leading production mechanism in each category

Category	Constant term	Sampling term	Material modeling	Noise term
Central-low p_{Tt}	7.5	2.6	4.9	2.6
Central-high p_{Tt}	9.6	5.6	6.2	1.7
Forward-low p_{Tt}	9.9	1.3	6.0	2.1
Forward-high p_{Tt}	12	2.8	7.8	1.9
VBF loose	9.4	2.6	6.0	2.1
VBF tight	10	3.8	6.5	2.1
VH hadronic	11	4.0	7.2	1.6
$VH E_T^{\text{miss}}$	11	3.6	7.4	1.7
VH one-lepton	9.8	2.8	6.3	2.1
VH dilepton	9.5	2.7	6.2	2.1
$t\bar{t}H$ hadronic	9.6	3.6	6.3	1.9
$t\bar{t}H$ leptonic	9.5	3.4	6.2	2.1

Relative systematic uncertainties due to jet energy scale, jet energy resolution and jet vertex fraction

Category	ggF	VBF	$t\bar{t}H$	$WH + ZH$
Central + Forward-low p_{Tt}	0.1	2.9	4.0	0.1
Central + Forward-high p_{Tt}	1.1	4.5	3.5	1.4
VBF loose	12	4.4	7.6	13
VBF tight	13	9.1	6.3	17
VH hadronic	2.8	4.1	9.5	2.5
$VH E_T^{\text{miss}}$	2.6	9.0	1.2	0.2
VH one-lepton	4.9	6.2	2.8	0.5
VH dilepton	0	0	5.1	1.0
$t\bar{t}H$ hadronic	11	21	7.3	22
$t\bar{t}H$ leptonic	37	7.7	0.5	7.4

Signal strength with Run 2 data




For more details see Alain Bellerive's talk

Signal Strength


Fiducial and differential cross-sections

Measure detector-corrected event yields without imposing theory assumptions







JHEP09(2014)112
ATLAS-CONF-2016-067

The correction factors is the ratio of events reconstructed at detector-and particle-levels



Jet multiplicity

Transverse momentum



Effective Lagrangian approach

CP-even

$$\mathcal{L}_{\text{eff}} = \bar{c}_\gamma O_\gamma + \bar{c}_g O_g + \bar{c}_{HW} O_{HW} + \bar{c}_{HB} O_{HB} + \tilde{c}_\gamma \tilde{O}_\gamma + \tilde{c}_g \tilde{O}_g + \tilde{c}_{HW} \tilde{O}_{HW} + \tilde{c}_{HB} \tilde{O}_{HB}$$


New HWW, HZZ, HZ γ interactions

CP-odd

Sensitivity to signal strength and differential distributions

$$\frac{d\sigma}{dX} = \sum_j \left(\frac{d\sigma_j}{dX} \right)^{\text{ref}} \cdot \left(\frac{d\sigma_j}{dX} \right)_{c_i}^{\text{MG5}} / \left(\frac{d\sigma_j}{dX} \right)_{c_i=0}^{\text{MG5}}$$

LO Madgraph used to model effect of HD operators. Assume QCD/EW HO corrections factorize from NP



Observed allowed ranges at 95% CL

Coefficient	95% 1 - CL limit
\bar{c}_γ	$[-7.4, 5.7] \times 10^{-4} \cup [3.8, 5.1] \times 10^{-3}$
\tilde{c}_γ	$[-1.8, 1.8] \times 10^{-3}$
\bar{c}_g	$[-0.7, 1.3] \times 10^{-4} \cup [-5.8, -3.8] \times 10^{-4}$
\tilde{c}_g	$[-2.4, 2.4] \times 10^{-4}$
\bar{c}_{HW}	$[-8.6, 9.2] \times 10^{-2}$
\tilde{c}_{HW}	$[-0.23, 0.23]$

Outlook and Conclusions

- ❑ **The di-photon final state plays a critical role in the understanding of the Higgs boson**
- ❑ **Signal strengths of different production mechanisms and their ratio to ggF are measured**
- ❑ **First fiducial total and differential cross-sections measured with Run 1 data**
- ❑ **Exploration of different production mechanisms a very important role for Run 2**
- ❑ **First measurements of signal strengths and fiducial cross-sections with Run 2 measured**
 - ❑ **Looking forward to measurements with significantly larger Run 2 data sets**

Extra Slides

TABLE I. Summary of event generators and PDF sets used to model the signal and the main background processes. The SM cross sections σ for the Higgs production processes with $m_H = 125.4$ GeV are also given separately for $\sqrt{s} = 7$ TeV and $\sqrt{s} = 8$ TeV, together with the orders of the calculations.


Process	Generator	Showering	PDF set	Order of calculation	σ [pb]	σ [pb]
					$\sqrt{s} = 7$ TeV	$\sqrt{s} = 8$ TeV
ggF	POWHEG-BOX	PYTHIA8	CT10	NNLO(QCD) + NLO(EW)	15.04	19.15
VBF	POWHEG-BOX	PYTHIA8	CT10	NLO(QCD + EW) + app.NNLO(QCD)	1.22	1.57
WH	PYTHIA8	PYTHIA8	CTEQ6L1	NNLO(QCD) + NLO(EW)	0.57	0.70
ZH	PYTHIA8	PYTHIA8	CTEQ6L1	NNLO(QCD) + NLO(EW)	0.33	0.41
$t\bar{t}H$	POWHEL	PYTHIA8	CT10	NLO(QCD)	0.09	0.13
$tHbj$	MADGRAPH	PYTHIA8	CT10	NLO(QCD)	0.01	0.02
tHW	MADGRAPH5_AMC@NLO	HERWIG++	CT10	NLO(QCD)	< 0.01	< 0.01
$b\bar{b}H$	5FS(NNLO) + 4FS(NLO)	0.15	0.20
$\gamma\gamma$	SHERPA	SHERPA	CT10			
γ -jet	SHERPA	SHERPA	CT10			
jet jet	PYTHIA8	PYTHIA8	CTEQ6L1			

	σ_{68} [GeV]	σ_{90} [GeV]
Central low- p_{Tt}	1.60	2.72
Central high- p_{Tt}	1.39	2.36
Forward low- p_{Tt}	2.07	3.65
Forward high- p_{Tt}	1.84	3.22
VBF loose	1.68	2.96
VBF tight	1.53	2.73
VH hadronic loose	1.65	2.90
VH hadronic tight	1.49	2.61
VH E_T^{miss}	1.65	2.86
VH one-lepton	1.68	2.99
VH dilepton	1.73	2.99
$t\bar{t}H$ hadronic	1.60	2.87
$t\bar{t}H$ leptonic	1.65	2.85


Cross-section measurements

- First Higgs differential cross section results
- Follows closely the coupling analysis strategy
- Allows to probe kinematics of Higgs boson
- As model-independent as possible

JHEP09(2014)112
ATLAS-CONF-2016-067



- Extract signal at reconstruction level
- Derive unfolding factors to unfold to particle level
- All results presented at particle level
- Also provide NP and fiducial correction factors for theorist use



Statistical correlations between the measured cross sections in bins of the diphoton transverse momentum and jet multiplicity distributions.

