



Trigger & DAQ



SLAC Summer Institute

Wesley H. Smith

U. Wisconsin - Madison

August 18, 2016

Outline:

Introduction to LHC Trigger & DAQ

Challenges & Architecture


LHC Experiments Trigger & DAQ

The future of LHC Trigger & DAQ

Triggering

- **Task: inspect detector information and provide a first decision on whether to keep the event or throw it out**


The trigger is a function of :



Event data & Apparatus
Physics channels & Parameters

- Detector data not (all) promptly available
 - Selection function highly complex
- ⇒ $T(\dots)$ is evaluated by successive approximations, the
- TRIGGER LEVELS**
- (possibly with zero dead time)


LHC Overview



Proton-Proton	2835 bunch/beam
Protons/bunch	10^{11}
Beam energy	7 TeV (7×10^{12} eV)
Luminosity	$10^{34} \text{ cm}^{-2} \text{ s}^{-1}$
Crossing rate	40 MHz

with every bunch crossing
 ~25 Minimum Bias events
 with ~2000 particles produced

**Selection of 1 in
 10,000,000,000,000**





LHC Physics – Trigger Challenge



Electroweak Symmetry Breaking Scale

- Higgs discovery and higgs sector characterization
- Quark, lepton Yukawa couplings to higgs

Low ≈ 40 GeV

Low P_T γ, e, μ

Low P_T B, τ jets

New physics at TeV scale to stabilize higgs sector

- Spectroscopy of new resonances (SUSY or otherwise)
- Find dark matter candidate

Multiple low P_T objects

Missing E_T

Multi-TeV scale physics (loop effects)

- Indirect effects on flavor physics (mixing, FCNC, etc.)
 - B_s mixing and rare B decays
- Lepton flavor violation
 - Rare Z and higgs decays

~ Dedicated triggers using displaced vertices


Low P_T leptons

Planck scale physics

- Large extra dimensions to bring it closer to experiment
- New heavy bosons
- Blackhole production

High P_T leptons and photons
Multi particle and jet events


The LHC Plan




↑
 You are here!

	Design	2010	2011	2012
Beam Energy (TeV)	7	3.5	3.5	4
Bunches/Beam	2808	368	1380	1380
Proton/Bunch (10^{11})	1.15	1.3	1.5	1.7
Peak Lumi. ($10^{32} \text{ cm}^{-2} \text{ s}^{-1}$)	100	2	30	76
Integrated Lumi. (fb^{-1})	100/yr	0.036	6	20
Pile-Up	23	~1	10	20
Bunch Spacing	25 ns	50 ns	50 ns	50 ns

Pile-Up – the number of proton interactions occurring during each bunch crossing



	Design	2015	2016	2017/8 (*)
Beam Energy (TeV)	7	6.5	6.5	6.5/7
Bunches/Beam	2808	2244	2748	2808
Proton/Bunch (10^{11})	1.15	1.2	1.2	1.2
Peak Lumi. ($10^{32} \text{ cm}^{-2} \text{ s}^{-1}$)	100	51	100(*)	100
Integrated Lumi. (fb^{-1})	100/yr	4	30(*)	70(*)
Pile-Up	23	<20	40	40
Bunch Spacing	25 ns	50/25 ns	25 ns	25 ns



Pile-Up – the number of proton interactions occurring during each bunch crossing

*expected value


At design $L = 10^{34} \text{cm}^{-2}\text{s}^{-1}$

- 23 pp events/25 ns xing
 - ~ 1 GHz input rate
 - “Good” events contain ~ 20 bkg. events
- 1 kHz W events
- 10 Hz top events
- $< 10^4$ detectable Higgs decays/year

Can store ~ 1 kHz events




Select in stages

- Level-1 Triggers
 - 1 GHz to 100 kHz
- High Level Triggers
 - 100 kHz to 1 kHz






Collisions (p-p) at LHC



Event size: ~1 MByte
Processing Power: ~X TFlop

LHC Trigger & DAQ Challenges



Challenges:

1 GHz of Input Interactions

Beam-crossing every 25 ns with ~ 23 interactions produces over 1 MB of data




Archival Storage up to 1 kHz of 1 MB events

Challenges: Pile-up

- “In-time” pile-up: particles from the same crossing but from a different pp interaction


- Long detector response/pulse shapes:

- ◆ “Out-of-time” pile-up: left-over signals from interactions in previous crossings
- ◆ Need “bunch-crossing identification”




PileUp: major trigger problem

$Z \rightarrow \mu\mu$ event from 2012 data with 25 vertices



Now


Simulated Event Display at 140 PU (102 Vertices)



High
Luminosity
LHC: 2025

Challenges: Time of Flight




$c = 30 \text{ cm/ns} \rightarrow \text{in } 25 \text{ ns, } s = 7.5 \text{ m}$



ATLAS & CMS Trigger & Readout Structure

≈ 30 Collisions/25ns
(10^9 event/sec)

10^7 channels
(10^{16} bit/sec)



ATLAS Run 1 Trigger & DAQ

Event rates design
(2012 peak)

40 MHz
(20 MHz)

<2.5 μ s

75 kHz
(~65 kHz)

~40 ms
(~100 ms)

3 kHz
(~6.5 kHz)

~4 s
(~1 s)

~200 Hz
(~1000 Hz)

Trigger

3 Levels

DAQ


Data rates design
(2012 peak)

ATLAS Event
1.5 MB/25 ns
(1.6 MB/50ns)

~110 GB/s
(~105 GB/s)

~4.5 GB/s
(~10.5 GB/s)

~300 MB/s
(~1600 MB/s)





ATLAS Run 2 Trigger & DAQ: Merged Levels 2,3



Event rates
(peak)

Trigger

DAQ

Data rates
(peak)

40 MHz


100 kHz

~ 1.5 kHz

- LVL1 decision made with calorimeter data with coarse granularity and muon trigger chamber data

- HLT first uses Region of Interest data (ca. 2%) with full granularity and combines HLTSV information from all detectors; performs fast rejection.

HLT then refines the selection, can perform event reconstruction at full granularity using latest alignment and calibration data



New HLT Architecture

New use of Tracking Information at start of HLT: FTK




CMS Run 1,2 Trigger & DAQ: 2 Levels




Level-1 Trigger

Event Manager

Detector Frontend



Builder Networks



Computing Services

- Data
- Trigger
- Back Pressure

— The Lv1-Accept decision causes the readout system to respond with backpressure information.


Run Control

Interaction rate: 1 GHz
 Bunch Xing rate: 40 MHz
 Level 1 Output: 100 kHz
 Storage Output: ~ 400 Hz
 Avg Event Size: ~ 0.5 MB
 Data production ~ 1 TB/day

Lv1 decision is distributed to front-ends & readout via TTC system (red).
Readout buffers designed to accommodate Poisson fluctuations from 100 kHz Lv1 trigger rate.


Present ATLAS & CMS Level 1 Trigger Data

Use prompt data (calorimetry and muons) to identify:
High p_t electron, muon, jets, missing E_T




CALORIMETERS

Cluster finding and energy deposition evaluation



New data every 25 ns
Decision latency $\sim \mu\text{s}$

- Process reduced granularity data from calorimeter and muon detectors
- Trigger decision based on object multiplicities
- Generate L1A and send via TTC distribution to detector front-ends to initiate readout
- Maximum round-trip latency 2.5 us
 - Data stores in on-detector pipelines
- Identify regions-of-interest (RoI) to seed L2 trigger
- Custom built electronics
- Synchronous, pipelined processing system operating at the bunch crossing rate



LVL1 triggers on high p_T objects


- Calorimeter cells and muon chambers to find $e/\gamma/\tau$ -jet- μ candidates above thresholds

LVL2 uses Regions of Interest as identified by Level-1


- Local data reconstruction, analysis, and sub-detector matching of Rol data

The total amount of Rol data is minimal

- $\sim 2\%$ of the Level-1 throughput but it has to be extracted from the rest at 75 kHz



ATLAS Run 2 L1 Trigger





Run 1 CMS L1 Trigger System



Lv1 trigger is based on calorimeter & muon detectors.


At L1 trigger on:

- 4 highest E_t e^\pm/γ
- 4 highest E_t central jets
- 4 highest E_t forward jets
- 4 highest E_t tau-jets
- 4 highest P_t muons

For each of these objects rapidity, η , and ϕ are also transmitted to Global Trigger for topological cuts & so Higher Level Triggers can seed on them.

Also trigger on inclusive triggers:

- E_t , MET, H_t



Generate L1A and send via TTC distribution to detector front-ends to initiate readout

Maximum round-trip latency 4 μ s

Data stored in on-detector pipelines


Lv1 trigger is based on calorimeter & muon detectors.

Increased η and φ granularity of the objects

Larger object available to the GlobalTrigger algorithms

- 12 highest E_t e^\pm/γ
- 12 highest E_t jets
- 8 highest E_t tau-jets
- 8 highest P_t muons


Larger reach of topological cuts at GlobalTrigger & so Higher Level Triggers can seed on them



Generate L1A and send via TCDS distribution to detector front-ends to initiate readout


Maximum round-trip latency $4 \mu\text{s}$

Data stored in on-detector pipelines



Muon Trigger looking for coincidences in muon trigger chambers
 2 out of 3 (low- p_T ; >6 GeV) and
 3 out of 3 (high- p_T ; >20 GeV)

Trigger efficiency 99% (low- p_T) and
 98% (high- p_T)



Trigger towers ($\Delta\eta \times \Delta\phi = 0.1 \times 0.1$)



Calorimetry Trigger looking for $e/\gamma/\tau$ + jets


- Various combinations of cluster sums and isolation criteria
- $\Sigma E_{T,em,had}$, $E_{T,miss}$

Calo. Trig: Improvements: Feature Extractor Processors

- Higher granularity in eta
0.025
 - present $\eta \times \phi = 0.1 \times 0.1$
- Segmentation in depth
- Higher resolution
(E_T : 0.125 GeV/count, now
is 1 GeV/count)


Expected Improvement wrt. Run 1 System

- At Pileup of 80:



Muon. Trig: Improvements:

- New Small Wheel
- Rejects tracks not from IP:
 - B: creation within toroid
 - C: multiple scattering
- Matching θ btw. Big Wheel and NSW








Present Endcap fake rate is almost 90%

- Angular resolution of 1 mrad (trigger)
 - After phase-2 BW upgrade
 - Until LS3: NSW confirmation of BW tracks with angular cut of ± 7 mrad

CMS Electron/Photon Algorithm: Basics



Trigger Primitive Generator

Fine grain

Flag Max of ( ,  ,  , ) & Sum ET 

Regional Calorimeter Trigger

E_T cut





 + Max () > Threshold


Longitudinal cut (H/E)

 AND /  < 0.05

Isolation, Hadronic & EM

 < 2 GeV

AND
One of ( ,  ,  , ) < 1 GeV



ELECTRON or PHOTON



CMS Electron/Photon Algorithm: Run 2 Version



Trigger Primitive Generator


Fine grain

Flag Max of (, , , ) & Sum ET




EGamma Identification



Dynamic clustering around a seed trigger tower (ET>2GeV)
Shape identification: based on ET, eta and cluster shape



e/g like




jet like



**ELECTRON
OR PHOTON**

CMS τ / Jet Algorithm: Run 1



Jet or τ E_T

- 12x12 trigger tower E_T sums in 4x4 region steps with central region $>$ others

- Larger trigger towers in HF but \sim same jet region size, $1.5 \eta \times 1.0 \phi$

τ algorithm (isolated narrow energy deposits), within $-2.5 < \eta < 2.5$

- Redefine jet as τ jet if none of the nine 4x4 region τ -veto bits are on

Output

- Top 4 τ -jets and top 4 jets in central rapidity, and top 4 jets in forward rapidity





CMS Jet Algorithms: From 2016



Sliding-Window Algorithm, centred on a local maximum ET trigger tower

- 9x9 trigger towers considered - corresponding to anti-kt jets of $R=0.4$
- jet position from the central (local maximum) TT
- jet ET from the 9x9 TT sum
- inequality mask to avoid self veto & double counting




PileUp rejection based on a “donut” algorithm



- energy in the four 3x9 trigger towers blocks around the jet used to estimate pile-up energy density

Topology can be used to distinguish hadronically decaying taus from taus:

- Enhanced position resolution by increasing granularity
- Introduce isolation as a handle to control rate
- Better energy resolution with specific calibration sequences




A few of the possible Stage-1 Patterns:





CMS Muon Chambers (> 2014*)





CMS Run 1 Muon Trigger



Counting Room: USC55
Cavern: UXC55

$|\eta| < 1.2$

DT hits

$0.8 < |\eta|$

CSC hits

$|\eta| < 2.4$

RPC hits

$|\eta| < 2.1$
 $|\eta| < 1.6$ in 2007

local trigger track segments
 $(\phi, \delta\phi, \eta, \delta\eta)$


local trigger track segments
 $(\phi, \delta\phi, \eta, \delta\eta)$

PATtern Comparator Trigger
 ≤ 4 barrel +
 ≤ 4 endcap
muon candidates
 $(p_t, \eta, \phi, \text{quality})$

regional trigger Barrel Track Finder
 ≤ 4 muon candidates
 $(p_t, \eta, \phi, \text{quality})$

regional trigger Endcap Track Finder
 ≤ 4 muon candidates
 $(p_t, \eta, \phi, \text{quality})$

Global Muon Trigger
 ≤ 4 muons
 $(p_t, \eta, \phi, \text{quality})$



(possibility of an isolated muon trigger)

BARREL MUON TRACK FINDER:

- DT + RPC Hits
- $0.8 < |\eta|$
- Optical links from the fronted of the DTs to the track finder boards (MP7)

ENDCAP MUON TRACK FINDER:

- CSC+RPC Hits
- $1.25 < |\eta| < 2.5$
- Optical signals sent from the CSC and RPC to the trigger boards (MTF7)
- Will include GEM detectors in the future

OVERLAP MUON TRACK FINDER

- DT+RPC+CSC
- $1.25 < |\eta| < 2.5$

All track finders assign eta/phi/pt and quality

GLOBAL MUON TRIGGER

- Receives muons raking according to pt accuracy
- Sorts and sends the 8 highest ranking ones to the GT


RPC pattern recognition

- Pattern catalog
- Fast logic

Memory to store patterns


Fast logic for matching

FPGAs are ideal




DT and CSC track finding:

- Finds hit/segments
- Combines vectors
- Formats a track
- Assigns p_t value




Drift Tubes (DT)

Drift Tubes



Meantimers recognize tracks and form vector / quartet.




Correlator combines them into one vector / station.

Match with RPC

Improve efficiency and quality

Cathod Strip Chambers (CSC)


CSC



Comparators give 1/2-strip resol.



Hit strips of 6 layers form a vector.



Sort based on P_T ,
Quality - keep loc.


Combine at next level
- match

Sort again - Isolate?

Top 4 highest P_T and
quality muons with
location coord.



CMS Global Trigger Runs 1&2





L1Menu: list of all the operational GT algorithms for a particular moment of data taking

Basic algorithms: counting single or multiple particles with energy above a threshold in a pseudorapidity range (eg: SingleMu16; DoubleEG20_10)


Complex algorithms take into account topological correlations of the candidates (eg: $\Delta\eta$ or invariant mass)

Output of the GT: L1 Accept after the check of the different combinations

Particle Conditions





Logical Combinations



Flexible algorithms implemented in FPGAs
100s of possible algorithms can be reprogrammed

High Level Trigger Strategy



High level triggers. CPU farms

- Finer granularity precise measurement
- Clean particle signature (π^0 - γ , isolation, ...)
- Kinematics. Effective mass cuts and topology
- Track reco and matching, b, τ -jet tagging
- Full event reconstruction and analysis

Successive improvements :
background event filtering,
physics selection





Start with L1 Trigger Objects



Electrons, Photons, τ -jets, Jets, Missing E_T , Muons

- HLT refines L1 objects (no volunteers)

Goal

- Keep L1T thresholds for electro-weak symmetry breaking physics
- However, reduce the dominant QCD background
 - From 100 kHz down to 100 Hz nominally

QCD background reduction




- Fake reduction: e^\pm , γ , τ
- Improved resolution and isolation: μ
- Exploit event topology: Jets
- Association with other objects: Missing E_T
- Sophisticated algorithms necessary
 - Full reconstruction of the objects
 - Due to time constraints we avoid full reconstruction of the event - L1 seeded reconstruction of the objects only
 - Full reconstruction only for the HLT passed events

“Level-2” electron:

- Search for match to Level-1 trigger
 - Use 1-tower margin around 4x4-tower trigger region
- Bremsstrahlung recovery “super-clustering”
- Select highest E_T cluster

Bremsstrahlung recovery:


- Road along ϕ — in narrow η -window around seed
- Collect all sub-clusters in road \rightarrow “super-cluster”




Cluster shape discrimination and isolation techniques similar or identical to the ones used offline after full reconstruction:

- precise energy and position determination
- enhanced background rejection

Cluster Shape Distribution



Single Electron Turn On for 32 GeV Online Cut





Muon HLT





Similar reconstruction to the offline one: tracker and muon chambers hits available for a full fit to the trajectory of the muon

- “Standalone” track reconstruction in the muon chambers only
- “Combined” reconstruction uniting Muon+Tracker

Outside-In and Inside-Out track fitting; track reconstruction quality; and depth of penetration in the system used to reduce misidentification

Isolation around the muon direction can be used to reduce rate

Typically high efficiencies and robustness versus pileup





Tracking & B tagging @ HLT

Offline algorithms for track reconstruction are too slow $O(10s)$ to be used online

—> Iterative tracking algorithm used to achieve $O(100ms)$:

- Each step reconstructs a specific subset of tracks (prompt, low/high pt, displaced)...
- First reconstruct the most energetic tracks (high pt seeds)—> remove hits associated to found tracks —> repeat pattern recognition with looser criteria





Tracking & B tagging @ HLT



Offline algorithms for track reconstruction are too slow $O(10s)$ to be used online

—> Iterative tracking algorithm used to achieve $O(100ms)$:


- Each step reconstructs a specific subset of tracks (prompt, low/high pt, displaced)...
- First reconstruct the most energetic tracks (high pt seeds)—> remove hits associated to found tracks —> repeat pattern recognition with looser criteria

Good HLT track reconstruction —> Fast primary vertex identification

—> impact parameter of tracks can be identified and used to tag displaced vertices

—> identify inclusive secondary vertices to tag events with a b-quark decay

—> similar algorithms to offline





τ -jet tagging at HLT




τ -jet ($E_t^{\tau\text{-jet}} > 60 \text{ GeV}$) identification (mainly) in the tracker:


Hard track, $p_t^{\text{max}} > 40 \text{ GeV}$, within $\Delta R < 0.1$ around calorimeter jet axis

Isolation: no tracks, $p_t > 1 \text{ GeV}$, within $0.03 < \Delta R < 0.4$ around the hard track


For 3-prong selection 2 more tracks in the signal cone $\Delta r < 0.03$



QCD jet rejection from isolation and hard track cuts



Further reduction by ~ 5 expected for 3-prong QCD jets from τ vertex reconstruction (CMS full simulation)



Jets and Energy Sums



2015, 13 TeV

Again, the techniques are now very close to offline ones

Both simple calorimetric based and PF-based algorithms are available for Jets, missing energy and HTT

Jet clustering:

- anti-kt jet with a 0.4 cone as the default jet algorithm
- anti-kt jets with 0.8 cone to trigger on boosted topologies (top, W,Z,Higgs tagging)
- offline-like pile up subtraction



ParticleFlow: Comprehensive event reconstruction algorithm that aim to identify all the particles in the event. Heavily used in CMS to exploit the excellent track reconstruction of the detector




CMS HLT Time Distribution (example from early 2011)




Prescale set used: $2E32 \text{ Hz/cm}^2$

Sample: MinBias L1-skim $5E32 \text{ Hz/cm}^2$ with 10 Pile-up




Monte Carlo simulation, MinimumBias @ 13TeV

Black: 20 PU Blue: 40 PU






CMS DAQ for Run 1



CMS Run 1 2-Stage Event Builder



CMS Run 2 DAQ ~ All New



10 GbE replacing Myrinet


New μ TCA readouts interfaced thru BU AMC13

New CPUs Infiniband switches

DAQ is an order of magnitude smaller!

Filter Farm: CMSSW from files

Distributed file system for backend



- BU-FU appliance**
- 1 BU (256 GB RAM, 2TB magnetic disks)
 - 16 FU nodes
 - FU: Dual E5-2670 8 core (2 x1 GbE)
 - FU: Dual X5650 6 core (2 x1 GbE)

- BU-FU appliance**
- 1 BU (256 GB RAM, 2TB magnetic disks)
 - 8 FU nodes
 - FU: Dual Haswell with 14 cores (10 GbE)




ATLAS Run 2 v. 1 TDAQ



Run2 system:

- new L1CALO pre-processor and interface to Central trigger.
- new muon chambers and Tile calorimeter input to endcap L1MUON.
- Central Trigger with new Topology Trigger and Central Trigger Processor modules.
- initial deployment of Fast Tracker (FTK) in HLT (Next slide)



Beyond Run 2 (Pre HL-LHC):

- Cal Trig: Increased granularity for better isolation
- Mu Trig: Endcap suppresses fakes using New Small Wheel



ATLAS FastTracker (FTK)



For Phase 1:

Dedicated hardware processor completes GLOBAL track reconstruction by beginning of level-2 processing.


- Allows very rapid rejection of most background, which dominates the level-1 trigger rate.
- Frees up level-2 farm to carry out needed sophisticated event selection algorithms.

Addresses two time-consuming stages in tracking

- Pattern recognition – find track candidates with enough Si hits
 - 10^9 prestored patterns simultaneously see each silicon hit leaving the detector at full speed.
- Track fitting – precise helix parameter & χ^2 determination
 - Equations linear in local hit coordinates give near offline resolution

Maintain current physics sensitivity at HL-LHC challenging for trigger

- EWK, top (and Higgs) scale physics remain critical for HL-LHC
- Cannot fit same “interesting” physics events in trigger at 13-14 TeV, $5 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$




• Increasing p_T thresholds reduces signal efficiency

- Trigger on lepton daughters from $H \rightarrow ZZ$ at $p_T \sim 10\text{-}20 \text{ GeV}$
- Very easy to reach the worst case: thresholds increase beyond energy scale of interesting processes

• Backgrounds from HL-LHC pileup further reduces the ability to trigger on rare decay products

- Leptons, photons no longer appear isolated and are lost in QCD backgrounds
- Increased hadronic activity from pileup impacts jet p_T and MET measurements

CMS HL-LHC Evolution



HL-LHC: Lumi = 5 - 7 x 10³⁴

<PU> = 140 - 200 (increase × 6 - 8 v. run 1)


E = 13-14 TeV (increase ~ 2 v. run 1)

25 nsec bunch spacing (reduce × 2 v. run 1)

Integrated Luminosity > 250 fb⁻¹ per year



ATLAS Upgrade Evolution





ATLAS & CMS L1 Tracking Trigger



Reduces Leptonic Trigger Rate

- Validate calorimeter or muon trigger object, e.g. discriminating electrons from hadronic ($\pi^0 \rightarrow \gamma\gamma$) backgrounds in jets
- Addition of precise tracks to improve precision on p_T measurement, sharpening thresholds in muon trigger
- Degree of isolation of e , γ , μ or τ candidate
- Requires calorimeter trigger trigger at the finest granularity to reduce electron trigger rate


Other Triggers

- Primary z-vertex location within 30 cm luminous region derived from projecting tracks found in trigger layers,
- Provide discrimination against pileup events in multiple object triggers, e.g. in lepton plus jet triggers.

Example: CMS Track

Trigger design finds stubs with $p_T \geq 2$:

Thanks to
CMS 3.8 T
magnetic field!





ATLAS & CMS @ HL-LHC

An introductory Summary



ATLAS*:

- Divide L1 Trigger into L0, L1 of latency 6, 30 μ sec, rate ≤ 1 MHz, ≤ 400 kHz, HLT output rate of 5-10 kHz
 - Calorimeter readout at 40 MHz w/backend waveform processing (140 Tbps)
- L0 uses Cal. & μ Triggers, which generate track trigger seeds
- L1 uses Track Trigger and more fine-grained calorimeter trigger information.

CMS:

- L1 Trigger latency: 12.5 μ sec
- L1 Trigger rate: 500 kHz (PU=140), 750 kHz (PU=200)
- L1 uses Track Trigger, finer granularity μ & calo. Triggers
- HLT output rate of 5 kHz (PU=140), 7.5 kHz (PU=200)



CMS Level-1 Tracking Trigger



Require:

- Highest possible efficiency over all η for isolated high P_T tracks
- Good efficiency for tracks in jets for vertex identification
- $P_T > 2-3$ GeV (small difference within this range)
 - Expect ~ 115 charged tracks with $P_T > 2$ GeV at PU = 140
 - Design for 300 tracks per bunch crossing
- Vertex resolution ~ 1 mm

Use:

- Charged Lepton ID
- Improve P_T resolution of charged leptons
- Determine isolation of leptons and photons
- Determine vertex of charged leptons and jet objects
- Determine primary vertex and MET from L1 Tracks from this vertex

Pixel Trigger Option

- Under consideration for now, but need a strong physics case
- Challenging to meet $12.5 \mu\text{sec}$ latency



CMS Estimation of required HL-LHC HLT Capabilities



Observation so far

- Required HLT power scales linearly with pile-up
 - This has been observed for PU in the range of 10-40
 - Conservatively assume this continues – needs verification

Assuming

- Linear scaling with average PU up to 2000
- A factor 1.5 due to energy increase to 13 TeV
 - Also conservative – takes into account complexity of events selected by L1 Trigger scaling with energy
 - Operation after LS1 with 6.5 TeV per beam will quickly allow refining this estimate

	LHC Run-I 7-8 TeV	LHC Phase-I upgr. 13 TeV	HL-LHC Phase-II upgr. 13 TeV	
Energy				
Peak Pile Up (Av./crossing)	35	50	140	200
Level-1 accept rate (maximum)	100 kHz	100 kHz	500 kHz	750 kHz
Event size (design value)	1 MB	1.5 MB	4.5 MB	5.0 MB
HLT accept rate	1 kHz	1 kHz	5 kHz	7.5 kHz
HLT computing power	0.2 MHS06	0.4 MHS06	6 MHS06	13 MHS06
Storage throughput (design value)	2 GB/s	3 GB/s	27 GB/s	42 GB/s



ATLAS Estimation of required HL-LHC HLT capabilities



Processing time extrapolations to PU = 200

- Run 4 rejection requirements similar/better than in Run 1-2 → 1k/100k vs 10k/400k

A factor O(50) in HLT compute power needed wrt to Run 1

Moore's law on a ~10 years period predicts a factor 100 increase

- Compute power requirements within expected technology envelope → HLT farm of similar size wrt to Run 1

BUT

- Software will have evolve to be at least as efficient as today on future technologies (GPGPU, Many-cores, ARM64, ...)



Assume a similar packaging → ~50 racks

Network:

- 5MB@400kHz → ~20 Tbps
- Reasonable to assume
 - 100 Gbps per CPU socket (computing unit)
 - established (>)400 Gbps technology
 - Infiniband EDR x12 → 300 Gbps
- Total number of ports ~unchanged
- Network topology and link speeds mix & match depend on compute power packaging

25% performance improvement per year is not the same as doubling performance every 2 years (more like 3)

However also important to notice that this is a power law, so small changes in assumed %/year lead to big differences on 10-20 year timescale..





Trigger & DAQ Summary

Continuously Evolving



ATLAS and CMS Level-1 Trigger (pre-LS3)

- Select 100 kHz interactions from 1 GHz
- Processing is synchronous & pipelined
- Decision latency is 3 μ s
- Algorithms run on local, coarse cal & muon data
 - Use of ASICs & FPGAs

ATLAS Level-1 Trigger (post-LS3):

- Divide L1 Trigger into L0, L1 of latency 6, 30 μ sec, rate \leq 1 MHz, \leq 0.4 MHz
- L0 uses Cal. & μ Triggers, which generate track trigger seeds
- L1 uses Track Trigger & more muon detectors & more fine-grained calorimeter trigger information.

CMS Level-1 Trigger (post LS3):

- L1 Trigger latency, rate: 12.5 μ sec, .5 – .75 MHz (140 – 200 PU)
- L1 uses Track Trigger, finer granularity μ & calo. Triggers

Higher Level Triggers

- Depending on experiment, done in one or two steps
- If two steps, first is hardware region of interest
- Then run software/algorithms as close to offline as possible on dedicated farm of PCs
- Pre-LS3 output rate of $<$ 1 kHz.
- Post LS3 HLT output rate of 5 – 10 (ATLAS)/7.5 (CMS) kHz (140 – 200 PU)