



Contribution ID: 147

Type: Oral

Ab initio study on the effect of phenyl substitution on the binding of carbon dioxide to salcomine

Thursday 25 May 2017 16:05 (15 minutes)

Metal-organic complexes, such as metal-porphyrins, can function as homogeneous catalysts for electrochemical reduction. In this work we present ab initio electronic structure calculations for a reaction involving carbon dioxide and salcomine, a coordination complex consisting of a salen ligand and a cobalt (II) atom. Changes to the binding energy due to substitutions on the phenyl ring by different functional groups are also studied. Cases where a shrinking HOMO-LUMO gap decreases the overpotential and increases the catalytic efficiency of the complex are analyzed. This study provides a theoretical basis for possible salen complex candidates as electrocatalyst for the reduction of carbon dioxide.

Primary author: Mr CHIONG III, Meliton (Materials Science and Engineering Program, College of Science, University of the Philippines, Diliman, Quezon City, Philippines)

Co-author: Dr PARAAN, Francis (National Institute of Physics, University of the Philippines, Diliman, Quezon City, Philippines)

Presenter: Mr CHIONG III, Meliton (Materials Science and Engineering Program, College of Science, University of the Philippines, Diliman, Quezon City, Philippines)

Session Classification: A15: Atomics

Track Classification: Atomic Physics, Quantum Physics, Molecular and Chemical Physics