

# Transport coefficients of a two component hadronic gas consisting of pions and nucleons

*Thursday 18 February 2016 16:50 (20 minutes)*

We derive the transport coefficients of a two component Hadronic gas consisting of Pions and Nucleons. We solve the relativistic Boltzmann equation with binary collision ( $\pi\pi$ ,  $\pi N$  and  $NN$ ) employing the relaxation time approximation method. In medium effects are introduced through one loop self-energies in the propagator of the exchange of  $\rho$  and  $\sigma$  for the  $\pi\pi$  cross-section, and one loop self-energy in the propagator of the exchange of  $\Delta$  for the  $\pi N$  cross-section. We use phenomenological amplitude for  $NN$  collisions. The effect of early freeze-out in heavy collision is implemented through temperature dependent pion chemical potential. The temperature dependence of the transport coefficients are studied.

**Primary author:** GANGOPADHYAYA, Utsab (Variable Energy Cyclotron Centre, Kolkata)

**Co-authors:** GHOSH, Snigdha (Variable Energy Cyclotron Centre, Kolkata); SARKA, Sourav (Variable Energy Cyclotron Centre, Kolkata); MITRA, Sukanya (Variable Energy Cyclotron Centre, Kolkata)

**Presenter:** GANGOPADHYAYA, Utsab (Variable Energy Cyclotron Centre, Kolkata)

**Session Classification:** Session 20