

National HEP meeting in Morocco.

Friday, 27 October 2017 - Saturday, 28 October 2017

Faculté des Sciences et Techniques de Tanger.



Book of Abstracts

Contents

Welcome Speech	1
On High Energy Physics and Quantum Information	1
High Energie Activities in LPMR	1
High Energie Activities in LPTPM	1
Particle Physics activities in ESMAR	1
Quantum physics activities in ESMAR	1
Beni Mellal participation	1
Tanger participation	2
Open discussion	2
High Energy Physics Activities in Marrakech	2
Poster session	2
W+4gamma signature from light charged Higgs boson at the LHC Run-2.	2
Radiative Neutrino Masses Models	2
Search of Dark Matter at the LHC, recent results from the ATLAS detector and future prospects	2
Dark Matter in String Theory. Two Generic Predictions for Dark Matter.	3
Open discussion	3
prospect of creating a new framework in collaboration with ICTP	3
HEP Activities	3
HEP Activities in Tanger	3
HEP Activities in Casablanca	3
HEP Activities in Agadir	4
Probing anomalous W tb couplings at the LHC.	4
High Energy physics activities	4

High Energy physics activities	4
Study of the Track-Based systematic uncertainties of the Soft Term Missing Transverse Momentum (MET)	4

1

Welcome Speech

Corresponding Authors: farida.fassi@cern.ch, aarhrib@ictp.it, yahya.tayalati@cern.ch

2

On High Energy Physics and Quantum Information

Corresponding Author: belhaj@unizar.es

3

High Energie Activities in LPMR

Corresponding Authors: ouali1962@gmail.com, abdelilah_moussa@yahoo.fr

4

High Energie Activities in LPTPM

Corresponding Authors: mohamed.ouchrif@cern.ch, jamal.derkaoui@cern.ch

5

Particle Physics activities in ESMAR

Corresponding Authors: yahya.tayalati@cern.ch, farida.fassi@cern.ch

6

Quantum physics activities in ESMAR

Corresponding Author: yassine.hassouni@gmail.com

7

Beni Mellal participation

8

Tanger participation

9

Open discussion

10

High Energy Physics Activities in Marrakech

Corresponding Author: mchabab@uca.ma

11

Poster session

12

W+4gamma signature from light charged Higgs boson at the LHC Run-2.

Corresponding Author: r.benbrik@uca.ma

13

Radiative Neutrino Masses Models

14

Search of Dark Matter at the LHC, recent results from the ATLAS detector and future prospects

Corresponding Author: rachid.mazini@cern.ch

15

Dark Matter in String Theory. Two Generic Predictions for Dark Matter.

Corresponding Author: bobby.acharya@cern.ch

16

Open discussion

17

prospect of creating a new framework in collaboration with ICTP

19

HEP Activities

Corresponding Author: r.benbrik@uca.ma

20

HEP Activities in Tanger

Corresponding Author: aarhrib@ictp.it

21

HEP Activities in Casablanca

Corresponding Authors: yousef.khoulaki@cern.ch, driss.benchekroun@cern.ch

22

HEP Activities in Agadir

Corresponding Author: mgouigh@cern.ch

34

Probing anomalous W tb couplings at the LHC.

Corresponding Author: adil.jueid@cern.ch

36

High Energy physics activities

39

High Energy physics activities

46

Study of the Track-Based systematic uncertainties of the Soft Term Missing Transverse Momentum (MET)

Authors: Farida Fassi¹; Rachid Mazini²; Souad Batlamous³

¹ *Mohammed V university in Rabat*

² *Academia Sinica (TW)*

³ *Universite Mohammed V (MA)*

Corresponding Authors: rachid.mazini@cern.ch, farida.fassi@cern.ch, souad.batlamous@cern.ch

The large hadron collider (LHC) is unique among accelerators currently existing or under construction. It is operating at a center of mass energy of 13 TeV and an integrated luminosity of about 41.5 fb⁻¹ to enable vital discoveries. A very good measurement of the missing transverse energy, E_{miss} , is essential for many physics studies in ATLAS both for Standard Model channels and for discovering channels. Events with large E_{miss} are expected to be the key signature for new physics such as supersymmetry and extra dimensions. E_{miss} is defined as the event momentum imbalance in the plane

transverse to the beam axis, where momentum conservation is expected. Such an imbalance may signal the presence of undetectable particles, such as neutrinos or new stable, weakly-interacting particles. The vector momentum imbalance in the transverse plane is obtained from the negative vector sum of the momenta of all particles detected in a proton-proton collision. The reconstruction of the Emiss involves combining contributions from the fully reconstructed and calibrated hard objects in the event, most of which rely on energy measurements in the calorimeters to provide some or all of the information about the object. Additionally, contributions from particles that do not pass the hard object Pt thresholds form a “soft” term. This poster presents the performance study of the Emiss Track-Based Soft Term. Development of a new approach to evaluate the Track-Based systematic uncertainties of the Emiss Soft Term is also discussed.

Summary: