

$Z \rightarrow \tau\tau$ with Safe Cuts

Carl Gwilliam



UNIVERSITY OF
LIVERPOOL



16th April 2009

Tau Workshop, Copenhagen

“Safe” Tau ID

- The full tau ID uses many variables from both the calorimeter and the tracker.
 - Some of these are likely to be less well understood at the beginning.
- Consequently, there has been an effort (B. Gosdzik et al.) to define a set of well understood variables for use in first data.
 - <http://indico.cern.ch/getFile.py/access?contribId=3&resId=0&materialId=slides&confId=43438>
- This uses a cut-based method with a limited set of robust variables. Two approaches:
 - Calorimeter only or calorimeter + tracking
- Will compare the performance of these on $Z \rightarrow \tau\tau$
 - Use official variables by running v15 and redo TauCommonSetIsTau
 - Thanks to Bjoern and Stan!

“Safe” Tau ID: Variables


- Calo variables:

- R_{EM}
- Isolation Fraction
- $(\Delta\eta)^2$,
- $E_t(EM)/E_t(EM+Had)$

- Tracking variables:

- $E_t/P_t(\text{lead-track})$
- $E_t(EM)/\Sigma P_t(\text{track})$
- $E_t(Had)/\Sigma P_t(\text{track})$
- $\Sigma p_t(\text{track})/E_t(EM+Had)$
- 2-track width


- Cuts values are dependent on both the P_t and multiplicity of the tau




B. Gosdzik

“Safe” Tau ID: Performance

CaloOnly, 1-prong, 25-45 GeV




CaloOnly, 3-prong, 25-45 GeV




- R_{EM} provides the strongest separation
- Significantly worse than LH in 3-prong case

Calo+Track, 1-prong, 25-45 GeV



Calo+Track, 3-prong, 25-45 GeV




- Define 3 cut levels:
 - Loose (70%)
 - Medium (50%)
 - **Tight (30%)**

B. Godzik

Signal and Backgrounds

- We consider only lepton-hadron mode:
 - Isolated lepton for triggering
 - Narrow tau-jet
 - Missing E_t (MET)



- Signal & (non-QCD) backgrounds @ 10 TeV:

| Process | Dataset | NLO Cross section (pb) | Filter Efficiency |
|---------------------------|---------|--------------------------|-------------------|
| $Z \rightarrow \tau\tau$ | 106052 | 1357 | - |
| $Z \rightarrow ee$ | 106050 | 1357 | 0.96 |
| $Z \rightarrow \mu\mu$ | 106051 | 1357 | 0.96 |
| $W \rightarrow e\nu$ | 106020 | 13814 | 0.87 |
| $W \rightarrow \mu\nu$ | 106021 | 13814 | 0.87 |
| $W \rightarrow \tau\nu$ | 106022 | 13814×0.352 | 0.87 |
| $W \rightarrow \tau_h\nu$ | 106023 | $13814 \times (1-0.352)$ | - |
| $t\bar{t} (ll, lh)$ | 105200 | 401.60×0.543 | - |

New

Cross section significantly lower at 10 TeV

- QCD di-jet background not yet been considered.
- All results that shown are normalised to 100 pb^{-1}

Selection




Based on post-CSC note ATL-COM-PHYS-2008-127

- Trigger
 - e10_medium || mu_10
- 1 isolated lepton
 - $P_t^l > 15$ GeV and $|\eta^l| < 2.5$
 - Track+calo isolation:
 - $P_t^{\text{trk}} \& E_t^{\text{calo}} (\Delta R=0.2) < 5$ GeV
 - Electron: medium isEM and author != softe
 - Muon: High P_t combined
- Missing E_t
 - MET > 20 GeV
 - $M_t(\text{lep}, \text{MET}) < 30$ GeV
 - $\Sigma E_t < 400$ GeV
- Make all lepton- τ pairs with:
 - $P_t^\tau > 15$ GeV and $|\eta^\tau| < 2.5$
 - $1 < \Delta\phi_{l,\tau\text{-jet}} < 3.1$ rad.
 - Tau ID:
 - Tight calo, calo+track, LH
 - 1 or 3 tracks
 - Remove overlaps with e/ μ
- Separate OS (signal) and SS (background) events
- Reconstruct visible mass
 - $37 < M_{l,\tau\text{-jet}} < 77$ GeV
- Generally, there is only 1 combination remaining




Missing E_t

1 lepton
+ Tau ID




LH



Calo +
track







Calo
(different
scale)







Track Multiplicity





LH



Calo + track



Calo
(different
scale)






Results

LH




Calo+Track

Calo

OS



SS



Results (2)

- Event-level cut flow:

| Cut | $Z \rightarrow \tau\tau$ | $Z \rightarrow ee$ | $Z \rightarrow \mu\mu$ | $W \rightarrow e\nu$ | $W \rightarrow \mu\nu$ | $W \rightarrow \tau\nu$ | $W \rightarrow \tau_h\nu$ | $t\bar{t}$ |
|---------------------|--------------------------|--------------------|------------------------|----------------------|------------------------|-------------------------|---------------------------|------------|
| Total # events | 135700 ± 320 | 130285 ± 120 | 130332 ± 119 | 1210962 ± 1334 | 1210525 ± 796 | 425106 ± 553 | 895147 ± 2068 | 21807 ± 47 |
| Trigger | 24342 ± 136 | 97497 ± 104 | 103864 ± 106 | 716716 ± 1026 | 742887 ± 623 | 128240 ± 304 | 4290 ± 143 | 12826 ± 36 |
| 1 isol. lepton | 15026 ± 107 | 51417 ± 75 | 44587 ± 69 | 568003 ± 914 | 692746 ± 602 | 82629 ± 244 | 817 ± 62 | 9338 ± 31 |
| $MET > 20$ | 4551 ± 59 | 1663 ± 14 | 25593 ± 53 | 493619 ± 852 | 538171 ± 531 | 49685 ± 189 | 568 ± 52 | 8527 ± 29 |
| $M_t^{l, MET} < 30$ | 2275 ± 41 | 320 ± 6 | 433 ± 7 | 4937 ± 85 | 5924 ± 56 | 5153 ± 61 | 48 ± 15 | 1238 ± 11 |
| $\sum E_t < 400$ | 2182 ± 41 | 285 ± 6 | 420 ± 7 | 4626 ± 82 | 5710 ± 55 | 4930 ± 60 | 38 ± 14 | 516 ± 7 |
| At least 1 cand. | 1955 ± 38 | 164 ± 4 | 328 ± 6 | 3929 ± 76 | 4948 ± 51 | 4341 ± 56 | 38 ± 14 | 515 ± 7 |

- At this stage W is the largest (non -QCD) background.

- Efficiency of safe cuts for $Z \rightarrow \tau\tau$

- $P_t^\tau > 15$ GeV and $|\eta^\tau| < 2.5$
- Truth-matched ($\Delta R < 0.1$)

| Cut | Level | Efficiency |
|-------------------|--------|------------|
| Safe Calo | Loose | 53.1% |
| | Medium | 39.1% |
| | Tight | 25.5% |
| Safe Calo + Track | Loose | 41.2% |
| | Medium | 30.6% |
| | Tight | 18.5% |

- What is the effect of the various tau IDs on $Z \rightarrow \tau\tau$ cut flow ...

Results (3)

LH

| Cut | $Z \rightarrow \tau\tau$ | $Z \rightarrow ee$ | $Z \rightarrow \mu\mu$ | $W \rightarrow e\nu$ | $W \rightarrow \mu\nu$ | $W \rightarrow \tau_l\nu$ | $W \rightarrow \tau_h\nu$ | $t\bar{t}$ |
|------------------------|--------------------------|--------------------|------------------------|----------------------|------------------------|---------------------------|---------------------------|------------|
| # candidates | 3027 ± 48 | 244 ± 5 | 459 ± 7 | 5650 ± 91 | 7226 ± 61 | 6224 ± 67 | 48 ± 15 | 1817 ± 13 |
| $1 < \Delta\phi < 3.1$ | 2685 ± 45 | 189 ± 5 | 398 ± 7 | 5070 ± 86 | 6457 ± 58 | 5614 ± 64 | 43 ± 14 | 1534 ± 12 |
| Tau ID | 240 ± 13 | 22 ± 2 | 10 ± 1 | 25 ± 6 | 20 ± 3 | 11 ± 3 | 0 ± 0 | 18 ± 1 |
| OS | 236 ± 13 | 21 ± 2 | 10 ± 1 | 22 ± 6 | 16 ± 3 | 9 ± 3 | 0 ± 0 | 16 ± 1 |
| $M_{l,\tau-jet}$ | 213 ± 13 | 0 ± 0 | 7 ± 1 | 12 ± 4 | 4 ± 1 | 2 ± 1 | 0 ± 0 | 5 ± 1 |
| SS | 5 ± 2 | 1 ± 0 | 0 ± 0 | 3 ± 2 | 4 ± 1 | 2 ± 1 | 0 ± 0 | 2 ± 0 |
| $M_{l,\tau-jet}$ | 3 ± 2 | 0 ± 0 | 0 ± 0 | 0 ± 0 | 2 ± 1 | 0 ± 0 | 0 ± 0 | 1 ± 0 |

$$\frac{S}{\sqrt{B}} = 39$$

Calo +
track

| Cut | $Z \rightarrow \tau\tau$ | $Z \rightarrow ee$ | $Z \rightarrow \mu\mu$ | $W \rightarrow e\nu$ | $W \rightarrow \mu\nu$ | $W \rightarrow \tau_l\nu$ | $W \rightarrow \tau_h\nu$ | $t\bar{t}$ |
|------------------------|--------------------------|--------------------|------------------------|----------------------|------------------------|---------------------------|---------------------------|------------|
| # candidates | 3027 ± 48 | 244 ± 5 | 459 ± 7 | 5650 ± 91 | 7226 ± 61 | 6224 ± 67 | 48 ± 15 | 1817 ± 13 |
| $1 < \Delta\phi < 3.1$ | 2685 ± 45 | 189 ± 5 | 398 ± 7 | 5070 ± 86 | 6457 ± 58 | 5614 ± 64 | 43 ± 14 | 1534 ± 12 |
| Tau ID | 242 ± 14 | 32 ± 2 | 11 ± 1 | 50 ± 9 | 48 ± 5 | 41 ± 5 | 0 ± 0 | 23 ± 2 |
| OS | 234 ± 13 | 30 ± 2 | 9 ± 1 | 38 ± 7 | 32 ± 4 | 28 ± 4 | 0 ± 0 | 20 ± 1 |
| $M_{l,\tau-jet}$ | 203 ± 12 | 1 ± 0 | 6 ± 1 | 19 ± 5 | 16 ± 3 | 14 ± 3 | 0 ± 0 | 9 ± 1 |
| SS | 8 ± 2 | 2 ± 0 | 2 ± 0 | 12 ± 4 | 15 ± 3 | 13 ± 3 | 0 ± 0 | 3 ± 1 |
| $M_{l,\tau-jet}$ | 3 ± 2 | 1 ± 0 | 1 ± 0 | 9 ± 4 | 7 ± 2 | 8 ± 2 | 0 ± 0 | 1 ± 0 |

$$\frac{S}{\sqrt{B}} = 25$$

Calo

| Cut | $Z \rightarrow \tau\tau$ | $Z \rightarrow ee$ | $Z \rightarrow \mu\mu$ | $W \rightarrow e\nu$ | $W \rightarrow \mu\nu$ | $W \rightarrow \tau_l\nu$ | $W \rightarrow \tau_h\nu$ | $t\bar{t}$ |
|------------------------|--------------------------|--------------------|------------------------|----------------------|------------------------|---------------------------|---------------------------|------------|
| # candidates | 3027 ± 48 | 244 ± 5 | 459 ± 7 | 5650 ± 91 | 7226 ± 61 | 6224 ± 67 | 48 ± 15 | 1817 ± 13 |
| $1 < \Delta\phi < 3.1$ | 2685 ± 45 | 189 ± 5 | 398 ± 7 | 5070 ± 86 | 6457 ± 58 | 5614 ± 64 | 43 ± 14 | 1534 ± 12 |
| Tau ID | 286 ± 15 | 45 ± 2 | 14 ± 1 | 97 ± 12 | 117 ± 8 | 114 ± 9 | 0 ± 0 | 35 ± 2 |
| OS | 265 ± 14 | 42 ± 2 | 10 ± 1 | 57 ± 9 | 74 ± 6 | 76 ± 7 | 0 ± 0 | 28 ± 2 |
| $M_{l,\tau-jet}$ | 235 ± 13 | 1 ± 0 | 6 ± 1 | 24 ± 6 | 30 ± 4 | 36 ± 5 | 0 ± 0 | 11 ± 1 |
| SS | 20 ± 4 | 3 ± 1 | 4 ± 1 | 40 ± 8 | 43 ± 5 | 38 ± 5 | 0 ± 0 | 7 ± 1 |
| $M_{l,\tau-jet}$ | 14 ± 3 | 1 ± 0 | 2 ± 1 | 18 ± 5 | 19 ± 3 | 22 ± 4 | 0 ± 0 | 3 ± 1 |

Similar to
v14 results

$$\frac{S}{\sqrt{B}} = 23$$

Conclusion and Plans

- Prospects for $Z \rightarrow \tau\tau$ in 100 pb^{-1} look promising and robust
 - Depends on the being able to control the QCD backgrounds
- Safe tau ID performs reasonably well compared to full ID
 - Some reduction in S/\sqrt{B} , but still a clear signal
 - Calo only cuts let in significantly more background (but also more signal)
 - Need to see the effects on the QCD background!
 - Safe likelihood/NN?
- Future plans:
 - Add in QCD backgrounds.
 - Study and optimise selection criteria for first data.
 - Investigate data-driven techniques for background estimation.

Back-up Slides