Hard Probe 2016



Contribution ID: 172

Type: not specified

## Measurement of the Shared Momentum Fraction $z_g$ using Jet Reconstruction in p+p and Au+Au Collisions with STAR

Sunday 25 September 2016 11:00 (20 minutes)

Light quark and gluon jet quenching in heavy ion collisions is described within the overwhelming majority of models as resulting predominantly from pQCD-type gluon radiation, but details of the underlying mechanisms differ greatly. One key difference lies in the treatment of the Altarelli-Parisi, AP, splitting functions. While in some models, such as Q-PYTHIA, the splitting functions are directly modified in the medium, this core component remains unchanged in others (e.g. YaJEM). It has been shown that the shared momentum fraction  $z_g$  can be made a Sudakov-safe measurement [1]. This quantity measures the  $p_T$  ratio between the two dominant branches as determined by the SoftDrop grooming process. We will present the first study of  $z_g$  measurements at top RHIC energy in p+p and Au+Au. The focus will be on the specific di-jet selection introduced in our previous momentum imbalance measurement, i.e. jets geometrically matched to "hard core" jets found using only constituents above 2 GeV/c and with a high tower above 5.5 GeV [2].

Such di-jet pairs were found to be significantly imbalanced with respect to p+p, yet regained balance when all soft constituents were included.

Individual examination of the splitting behavior of leading and recoil jet adds a new dimension to this observation, and new input to energy loss models.

[1] A. J. Larkoski, S. Marzani and J. Thaler, Phys. Rev. D 91, no. 11, 111501 (2015) [arXiv:1502.01719 [hep-ph]].

[2] K. Kauder (STAR Collaboration), Nuclear and Particle Physics Proceedings, Hard Probes 2015

## Summary

## **Presentation type**

Oral

Author: KAUDER, Kolja (Wayne State University)

Presenter: KAUDER, Kolja (Wayne State University)

Session Classification: Parallel Session VI: Jet Modification in A+A (IV)